

Optimal Stopping for Non-linear Expectations

BY ERHAN BAYRAKTAR, SONG YAO

Department of Mathematics, University of Michigan, Ann Arbor, MI 48109;
email: erhan@umich.edu. E. Bayraktar is supported in part by the National Science Foundation under an applied mathematics research grant and a Careergrant, DMS-0906257 and DMS-0955463, respectively, and in part by the Susan M. Smith Professorship. .

Department of Mathematics, University of Michigan, Ann Arbor, MI 48109;
email: songyao@umich.edu.

Abstract

We develop a theory for solving continuous time optimal stopping problems for non-linear expectations. Our motivation is to consider problems in which the stopper uses risk measures to evaluate future rewards.

Keywords: Nonlinear expectations, Optimal stopping, Snell envelope, Stability, g -expectations.

1 Introduction

We solve continuous time optimal stopping problems in which the reward is evaluated using *non-linear* expectations. Our purpose is to use criteria other than the expected value to evaluate the present value of future rewards. Such criteria include *risk measures*, which are not necessarily linear. Given a filtered probability space $(\Omega, \mathcal{F}, P, \mathbf{F} = \{\mathcal{F}_t\}_{t \in [0, T]})$ satisfying the *usual assumptions*, we define a filtration-consistent non-linear expectation (\mathbf{F} -expectation for short) with domain Λ as a collection of operators $\left\{ \mathcal{E}[\cdot | \mathcal{F}_t]: \Lambda \mapsto \Lambda \stackrel{\Delta}{=} \Lambda \cap L^0(\mathcal{F}_t) \right\}_{t \in [0, T]}$ satisfying "Monotonicity", "Time-Consistency", "Zero-one Law" and "Translation-Invariance". This definition is similar to the one proposed in [\[citet|Pln\]](#). A notable example of an \mathbf{F} -expectation is the so-called *g-expectation*, introduced by [\[citet|Peng-97\]](#). A fairly large class of *convex risk measures* (see e.g. [\[citet|Follmer_Schied_2004\]](#) for the definition

of risk measures) are g -expectations (see [\[CHMP\]](#), [\[Pln\]](#), [\[MaYao_08\]](#) and [\[HMPY-07\]](#)).

We consider two optimal stopping problems. In the first one, the stopper aims to find an optimal stopping time when there are multiple priors and the *Nature* is in cooperation with the stopper; i.e., the stopper finds an optimal stopping time that attains

$$Z(0) \triangleq \sup_{(i, \rho) \in \mathcal{I} \times \mathcal{S}_{0,T}} \mathcal{E}_i[Y_\rho + H_\rho^i | \mathcal{F}_0], \quad (1.1)$$

in which $\mathcal{E} = \{\mathcal{E}_i\}_{i \in \mathcal{I}}$ is a *stable* class of \mathbf{F} -expectations, $\mathcal{S}_{0,T}$ is the set of stopping times that take value in $[0, T]$. The reward process Y is a right-continuous \mathbf{F} -adapted process and for any $\nu \in \mathcal{S}_{0,T}$, Y_ν belongs to $\Lambda^\# \triangleq \{\xi \in \Lambda \mid \xi \geq c, \text{ a.s. } c \in \mathbb{R}\}$, where Λ is the common domain of the elements in \mathcal{E} . On the other hand, the *model-dependent* reward processes $\{H^i\}_{i \in \mathcal{I}}$ is a family of right-continuous adapted processes with $H_0^i = 0$ that is *consistent* with \mathcal{E} . We will express the solution of this problem in terms of the \mathcal{E} -upper Snell envelope Z^0 of Y_t , the smallest RCLL \mathbf{F} -adapted process dominating Y such that $Z^{i,0} \triangleq \{Z_t^0 + H_t^i\}_{t \in [0,T]}$ is an $\tilde{\mathcal{E}}_i$ -supermartingale for each $i \in \mathcal{I}$.

The construction of the Snell envelope is not straightforward. First, for any $i \in \mathcal{I}$, the conditional expectation $\mathcal{E}_i[\xi | \mathcal{F}_\nu]$, $\xi \in \Lambda$ and $\nu \in \mathcal{S}_{0,T}$ may not be well defined. However, we show that $t \rightarrow \mathcal{E}_i[\xi | \mathcal{F}_t]$ admits a right-continuous modification $t \rightarrow \tilde{\mathcal{E}}_i[\xi | \mathcal{F}_t]$ for any $\xi \in \Lambda^\#$ and that $\tilde{\mathcal{E}}_i$ is itself an \mathbf{F} -expectation on $\Lambda^\#$ such that $\tilde{\mathcal{E}}_i[\cdot | \mathcal{F}_\nu]$ is well defined on $\Lambda^\#$ for any $\nu \in \mathcal{S}_{0,T}$. In terms of $\tilde{\mathcal{E}}_i$ we have that

$$Z(0) = \sup_{(i, \rho) \in \mathcal{I} \times \mathcal{S}_{0,T}} \tilde{\mathcal{E}}_i[Y_\rho + H_\rho^i | \mathcal{F}_0]. \quad (1.2)$$

Finding a RCLL modification requires the development of an upcrossing theorem. This theorem relies on the strict monotonicity of \mathcal{E}_i and other mild hypotheses, one of which is equivalent to having lower semi-continuity (i.e. Fatou's lemma). Thanks to the right continuity of $t \rightarrow \tilde{\mathcal{E}}_i[\xi | \mathcal{F}_t]$, we also have an optional sampling theorem for right-continuous $\tilde{\mathcal{E}}_i$ -supermartingales. Another important tool in finding an optimal stopping time, the dominated convergence theorem is also developed under another mild assumption.

The stability assumption we make on the family \mathcal{E} is another essential ingredient in the construction of the Snell envelope. It guarantees that the class \mathcal{E} is closed under *pasting*: for any $i, j \in \mathcal{I}$ and $\nu \in \mathcal{S}_{0,T}$ there exists a $k \in \mathcal{I}$ such that $\tilde{\mathcal{E}}_k[\xi | \mathcal{F}_\sigma] = \tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_j[\xi | \mathcal{F}_{\nu \vee \sigma}] | \mathcal{F}_\sigma]$, for any $\sigma \in \mathcal{S}_{0,T}$. Under this assumption it can then be seen, for example, that the collection of random variables $\{\tilde{\mathcal{E}}_i[X(\rho) + H_\rho^i - H_\nu^i | \mathcal{F}_\nu], (i, \rho) \in \mathcal{I} \times \mathcal{S}_{\nu,T}\}$ is directed upwards. When the constituents of \mathcal{E} are linear expectations, the notion

of stability of this collection is given by (, ?, Definition 6.44), who showed that pasting two probability measures equivalent to P at a stopping time one will result in another probability measure equivalent to P . Our result in Proposition 75 shows that we have the same pasting property for \mathbf{F} -expectations. As we shall see, the stability assumption is crucial in showing that the Snell envelope is a supermartingale. This property of the Snell envelope is a generalization of *time consistency*, i.e.,

$$\operatorname{esssup}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[\xi | \mathcal{F}_\nu] = \operatorname{esssup}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i \left[\operatorname{esssup}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[\xi | \mathcal{F}_\sigma] \middle| \mathcal{F}_\nu \right], \quad a.s., \quad \forall \nu, \sigma \in \mathcal{S}_{0,T} \text{ with } \nu \leq \sigma, \quad a.s. \quad (1.3)$$

(, ?, Theorem 12) showed in the linear expectations case that the time consistency (1.3) is equivalent to the stability.

When the reward $t \rightarrow Y_t + H_t^i$ is " \mathcal{E} -uniformly-left-continuous" and each non-linear expectation in \mathcal{E} is convex, we can find an optimal stopping time $\bar{\tau}(0)$ for (1.1) in terms of the Snell envelope. Then we can solve the problem

$$\sup_{\rho \in \mathcal{S}_{0,T}} \mathcal{E}_i[Y_\rho + H_\rho^i | \mathcal{F}_0], \quad (1.4)$$

when $\mathcal{E}_i[\cdot | \mathcal{F}_t]$ has among other properties strict monotonicity, lower semi-continuity, dominated convergence theorem and the upcrossing lemma. Note that although, $\operatorname{esssup}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[\cdot | \mathcal{F}_t]$ has similar properties to $\tilde{\mathcal{E}}_i[\cdot | \mathcal{F}_t]$ (and that might lead one to think that (1.1) can actually be considered as a special case of (1.4)), the former does not satisfy strict monotonicity, the upcrossing lemma, and the dominated convergence theorem. One motivation for considering optimal stopping with multiple priors is to solve optimal stopping problems for "non-linear expectations" which do not satisfy these properties.

We show that the collection of g -expectations with uniformly Lipschitz generators satisfy the uniform left continuity assumption. Moreover, a g -expectation satisfies all the assumptions we ask of each \mathcal{E}_i for the upcrossing theorem, Fatou's lemma and the dominated convergence theorem to hold; and pasting of g -expectations results in another g -expectation. As a result the case of g -expectations presents a non-conventional example in which we can determine an optimal stopping time for (1.1). In fact, in the g -expectation example we can even find an optimal *prior* $i^* \in \mathcal{I}$, i.e.,

$$Z(0) = \mathcal{E}_i[Y_{\bar{\tau}(0)} + H_{\bar{\tau}(0)}^i | \mathcal{F}_0]. \quad (1.5)$$

In the second problem, the *stopper* tries to find a robust optimal stopping time that attains

$$V(0) \triangleq \sup_{\rho \in \mathcal{S}_{0,T}} \inf_{i \in \mathcal{I}} \mathcal{E}_i[Y_\rho \neg + \neg \neg H_\rho^i | \mathcal{F}_0]. \quad (1.6)$$

Under the “ \mathcal{E} -uniform-right-continuity” assumption, we find an optimal stopping time in terms of the \mathcal{E} -lower Snell envelope. An immediate by-product is the following minimax theorem

$$V(0) = \inf_{i \in \mathcal{I}} \sup_{\rho \in \mathcal{S}_{0,T}} \mathcal{E}_i[Y_\rho \neg + \neg \neg H_\rho^i | \mathcal{F}_0]. \quad (1.7)$$

Our paper was inspired by [\[Kara_Zam_2006\]](#) and [\[Kara_Zam_2008\]](#), which developed a martingale approach to solving (1.1) and (1.6), when \mathcal{E} is a class of linear expectations. In particular, [\[Kara_Zam_2006\]](#) considered the *controller-stopper* problem

$$\sup_{\rho \in \mathcal{S}_{0,T}} \sup_{U \in \mathcal{U}} \mathbf{E}^u \left[g(X(\rho)) + \int_0^\rho h(s, X, U_s) ds \right], \quad (1.8)$$

where $X(t) = x + \int_0^t f(s, X, U_s) ds + \int_0^t \sigma(s, X) dW_s^U$. In this problem, the stability condition is automatically satisfied. Here, g and h are assumed to be bounded measurable functions. Our results on g -expectations extend the results of [\[Kara_Zam_2006\]](#) from bounded rewards to rewards satisfying linear growth. [\[Delbaen_2006\]](#), [\[Kara_Zam_2005\]](#) also considered (1.1) when the \mathcal{E}_i 's are linear expectations. The latter paper made a *convexity* assumption on the collection of equivalent probability measures instead of a stability assumption. On the other hand, the discrete time version of the robust optimization problem was analyzed by [\[Follmer_Schied_2004\]](#). Also see [\[?, Sections 5.2 and 5.3\]](#).

The rest of the paper is organized as follows: In Section 1.1 we will introduce some notations that will be used throughout the paper. In Section 7, we define what we mean by an \mathbf{F} -expectation \mathcal{E} , propose some basic hypotheses on \mathcal{E} and discuss their implications such as Fatou's lemma, dominated convergence theorem and upcrossing lemma. We show that $t \rightarrow \mathcal{E}[\cdot | \mathcal{F}_t]$ admits a right-continuous modification which is also an \mathbf{F} -expectation and satisfies Fatou's lemma and the dominated convergence theorem. This step is essential since $\mathcal{E}[\cdot | \mathcal{F}_\nu]$, $\nu \in \mathcal{S}_{0,T}$ may not be well defined. We also show that the optional sampling theorem holds. The results in Section 7 will be the backbone of our analysis in the later sections.

In Section 7 we introduce the stable class of \mathbf{F} -expectations and review the properties of essential extremum. In Section 4 we solve (1.2), and find an optimal stopping time in terms of the \mathcal{E} -upper Snell envelope. On the other hand, in Section 5 we solve the robust optimization problem (1.6) in terms of the \mathcal{E} -lower Snell envelope. In Section 6, we give some interpretations and remarks on our results in the previous sections. In Section 7 we consider the case when \mathcal{E} is a certain collection of g -expectations. We see that in this framework, our assumptions on each \mathcal{E}_i , the stability condition and the uniform left/right continuity conditions are naturally satisfied. We also determine an optimal prior $i^* \in \mathcal{I}$ satisfying (1.5). Moreover, we show how the controller and stopper problem of [\[Kara_Zam_2006\]](#) fits into our g -expectations framework. This lets us extend their result from bounded rewards to rewards satisfying linear growth. In this section, we also solve

optimal stopping problem for quadratic g -expectations. The proofs of our results are presented in Section 8.

1.1 Notation

Throughout this paper, we fix a finite time horizon $T > 0$ and consider a complete probability space (Ω, \mathcal{F}, P) equipped with a right continuous filtration $\mathbf{F} \triangleq \{\mathcal{F}_t\}_{t \in [0, T]}$, not necessarily Brownian one, such that \mathcal{F}_0 is generated by all P -null sets in \mathcal{F} (in fact, \mathcal{F}_0 collects all measurable sets with probability 0 or 1). Let $\mathcal{S}_{0, T}$ be the collection of all \mathbf{F} -stopping times ν such that $0 \leq \nu \leq T$, a.s. For any $\nu, \sigma \in \mathcal{S}_{0, T}$ with $\nu \leq \sigma$, a.s., we define $\mathcal{S}_{\nu, \sigma} \triangleq \{\rho \in \mathcal{S}_{0, T} \mid \nu \leq \rho \leq \sigma, \text{ a.s.}\}$ and let $\mathcal{S}_{\nu, \sigma}^F$ denote all finite-valued stopping times in $\mathcal{S}_{\nu, \sigma}$. We let $\mathcal{D} = \{k 2^{-n} \mid k \in \mathbb{Z}, n \in \mathbb{N}\}$ denote the set of all dyadic rational numbers and set $\mathcal{D}_T \triangleq ([0, T) \cap \mathcal{D}) \cup \{T\}$. For any $t \in [0, T]$ and $n \in \mathbb{N}$, we also define

$$q_n^+(t) \triangleq \frac{\lceil 2^n t \rceil}{2^n} \wedge T. \quad (1.9)$$

It is clear that $q_n^+(t) \in \mathcal{D}_T$.

In what follows we let \mathcal{F}' be a generic sub- σ -field of \mathcal{F} and let \mathbb{B} be a generic Banach space with norm $|\cdot|_{\mathbb{B}}$. The following spaces of functions will be used in the sequel.

(1) For $0 \leq p \leq \infty$, we define

- $L^p(\mathcal{F}'; \mathbb{B})$ to be the space of all \mathbb{B} -valued, \mathcal{F}' -measurable random variables ξ such that $E(|\xi|_{\mathbb{B}}^p) < \infty$. In particular, if $p = 0$, $L^0(\mathcal{F}'; \mathbb{B})$ stands for the space of all \mathbb{B} -valued, \mathcal{F}' -measurable random variables; and if $p = \infty$, $L^\infty(\mathcal{F}'; \mathbb{B})$ denotes the space of all \mathbb{B} -valued, \mathcal{F}' -measurable random variables ξ with $\|\xi\|_\infty \triangleq \operatorname{esssup}_{\omega \in \Omega} |\xi(\omega)|_{\mathbb{B}} < \infty$.
- $L_{\mathbf{F}}^p([0, T]; \mathbb{B})$ to be the space of all \mathbb{B} -valued, \mathbf{F} -adapted processes X such that $E \int_0^T |X_t|_{\mathbb{B}}^p dt < \infty$. In particular, if $p = 0$, $L_{\mathbf{F}}^0([0, T]; \mathbb{B})$ stands for the space of all \mathbb{B} -valued, \mathbf{F} -adapted processes; and if $p = \infty$, $L_{\mathbf{F}}^\infty([0, T]; \mathbb{B})$ denotes the space of all \mathbb{B} -valued, \mathbf{F} -adapted processes X with $\|X\|_\infty \triangleq \operatorname{esssup}_{(t, \omega) \in [0, T] \times \Omega} |X_t(\omega)|_{\mathbb{B}} < \infty$.
- $\mathbb{C}_{\mathbf{F}}^p([0, T]; \mathbb{B}) \triangleq \{X \in L_{\mathbf{F}}^p([0, T]; \mathbb{B}) : \langle \text{hbox}|X \text{ has continuous paths} \rangle\}$.
- $\mathcal{H}_{\mathbf{F}}^p([0, T]; \mathbb{B}) \triangleq \{X \in L_{\mathbf{F}}^p([0, T]; \mathbb{B}) : \langle \text{hbox}|X \text{ is predictably measurable} \rangle\}$.

(2) For $p \geq 1$, we define a Banach space

$$M_{\mathbf{F}}^p([0, T]; \mathbb{B}) = \left\{ X \in \mathcal{H}_{\mathbf{F}}^0([0, T]; \mathbb{B}) : \|X\|_{M^p} \triangleq \left\{ E \left[\left(\int_0^T |X_s|_{\mathbb{B}}^2 ds \right)^{p/2} \right] \right\}^{1/p} < \infty \right\},$$

and denote $M_{\mathbf{F}}([0, T]; \mathbb{B}) \stackrel{\Delta}{=} \bigcap_{p \geq 1} M_{\mathbf{F}}^p([0, T]; \mathbb{B})$.

(3) We further define

$$\begin{aligned} L^{-e}(\mathcal{F}; \mathbb{B}) &\stackrel{\Delta}{=} \{ \xi \in L^0(\mathcal{F}; \mathbb{B}) : E[e^{\lambda|\xi|_{\mathbb{B}}}] < \infty \text{ for all } \lambda > 0 \}, \\ \mathbb{C}_{\mathbf{F}}^e([0, T]; \mathbb{B}) &\stackrel{\Delta}{=} \left\{ X \in \mathbb{C}_{\mathbf{F}}^0([0, T]; \mathbb{B}) : E \left[\exp \left\{ \lambda \sup_{t \in [0, T]} |X_t|_{\mathbb{B}} \right\} \right] < \infty \text{ for all } \lambda > 0 \right\}. \end{aligned}$$

For any $p > 0$, we denote $\mathcal{M}^p(\mathbb{R}^d)$ to be the space of all \mathbb{R}^d -valued predictable processes X such that

$$\|X\|_{\mathcal{M}^p} \stackrel{\Delta}{=} \left(E \left(\int_0^T |X_s|^2 ds \right)^{p/2} \right)^{1 \wedge 1/p} < \infty. \quad (1.10)$$

We note that for $p \geq 1$, $\mathcal{M}^p(\mathbb{R}^d)$ is a Banach space with the norm $\|\cdot\|_{\mathcal{M}^p}$, and for $p \in (0, 1)$, $\mathcal{M}^p(\mathbb{R}^d)$ is a complete metric space with the distance defined through (1.10). [\(fi\)](#)

If $d = 1$, we shall drop $\mathbb{B} = \mathbb{R}$ from the above notations (e.g., $L_{\mathbf{F}}^p([0, T]) = L_{\mathbf{F}}^p([0, T]; \mathbb{R})$, $L^p(\mathcal{F}_T) = L^p(\mathcal{F}_T; \mathbb{R})$). In this paper, all \mathbf{F} -adapted processes are supposed to be real-valued unless specifying otherwise.

2 \mathbf{F} -expectations and Their Properties

We will define non-linear expectations on subspaces of $L^0(\mathcal{F}_T)$ satisfying certain algebraic properties, which are listed in the definition below.

Definition 1. Let \mathcal{D}_T denote the collection of all non-empty subsets Λ of $L^0(\mathcal{F}_T)$ satisfying:

- D1 $0, 1 \in \Lambda$;
- D2 Λ is closed under addition and under multiplication with indicator random variables. Namely, for any $\xi, \eta \in \Lambda$ and $A \in \mathcal{F}_T$, both $\xi + \eta$ and $11_A \xi$ belong to Λ ;
- D3 Λ is positively solid: For any $\xi, \eta \in L^0(\mathcal{F}_T)$ with $0 \leq \xi \leq \eta$, a.s., if $\eta \in \Lambda$, then $\xi \in \Lambda$ as well.

Remark 2. (1) Each $\Lambda \in \mathcal{D}_T$ is also closed under maximization " \vee " and under minimization " \wedge ": In fact, for any $\xi, \eta \in \Lambda$, since the set $\{\xi > \eta\} \in \mathcal{F}_T$, (D2) implies that $\xi \vee \eta = \xi 11_{\{\xi > \eta\}} + \eta 11_{\{\xi \leq \eta\}} \in \Lambda$. Similarly, $\xi \wedge \eta \in \Lambda$;

- (2) For each $\Lambda \in \mathcal{D}_T$, (D1)-(D3) imply that $c \in \Lambda$ for any $c \geq 0$;
- (3) \mathcal{D}_T is closed under intersections: If $\{\Lambda_i\}_{i \in \mathcal{I}}$ is a subset of \mathcal{D}_T , then $\bigcap_{i \in \mathcal{I}} \Lambda_i \in \mathcal{D}_T$; \mathcal{D}_T is closed under unions of increasing sequences: If $\{\Lambda_n\}_{n \in \mathbb{N}} \subset \mathcal{D}_T$ such that $\Lambda_n \subset \Lambda_{n+1}$ for any $n \in \mathbb{N}$, then $\bigcup_{n \in \mathbb{N}} \Lambda_n \in \mathcal{D}_T$;
- (4) It is clear that $L^p(\mathcal{F}_T) \in \mathcal{D}_T$ for all $0 \leq p \leq \infty$.

Definition 3. An **F**-consistent non-linear expectation **F** – expectation for short is a pair \mathcal{E}, Λ in which $\Lambda \in \mathcal{D}_T$ and \mathcal{E} denotes a family of operators $\{\mathcal{E}[\cdot|\mathcal{F}_t]: \Lambda \mapsto \Lambda_t \triangleq \Lambda \cap L^0(\mathcal{F}_t)\}_{t \in [0, T]}$ satisfying the following hypothesis for any $\xi, \eta \in \Lambda$ and $t \in [0, T]$:

- (A1) “Monotonicity (positively strict)”: $\mathcal{E}[\xi|\mathcal{F}_t] \leq \mathcal{E}[\eta|\mathcal{F}_t]$, a.s. if $\xi \leq \eta$, a.s.; Moreover, if $0 \leq \xi \leq \eta$ a.s. and $\mathcal{E}[\xi|\mathcal{F}_0] = \mathcal{E}[\eta|\mathcal{F}_0]$, then $\xi = \eta$, a.s.;
- (A2) “Time Consistency”: $\mathcal{E}[\mathcal{E}[\xi|\mathcal{F}_t]|\mathcal{F}_s] = \mathcal{E}[\xi|\mathcal{F}_s]$, a.s. for any $0 \leq s \leq t \leq T$;
- (A3) “Zero-one Law”: $\mathcal{E}[\mathbf{1}_A \xi|\mathcal{F}_t] = \mathbf{1}_A \mathcal{E}[\xi|\mathcal{F}_t]$, a.s. for any $A \in \mathcal{F}_t$;
- (A4) “Translation Invariance”: $\mathcal{E}[\xi + \eta|\mathcal{F}_t] = \mathcal{E}[\xi|\mathcal{F}_t] + \eta$, a.s. if $\eta \in \Lambda_t$.

We denote the domain Λ by $Dom(\mathcal{E})$ and define

$$Dom_\nu(\mathcal{E}) \triangleq Dom(\mathcal{E}) \cap L^0(\mathcal{F}_\nu), \quad \forall \nu \in \mathcal{S}_{0, T}.$$

For any $\xi, \eta \in Dom(\mathcal{E})$ with $\xi = \eta$, a.s., (A1) implies that $\mathcal{E}[\xi|\mathcal{F}_t] = \mathcal{E}[\eta|\mathcal{F}_t]$, a.s. for any $t \in [0, T]$, which shows that the **F**-expectation $(\mathcal{E}, Dom(\mathcal{E}))$ is well-defined. Moreover, since $Dom_0(\mathcal{E}) = Dom(\mathcal{E}) \cap L^0(\mathcal{F}_0) \subset L^0(\mathcal{F}_0) = \mathbb{R}$, $\mathcal{E}[\cdot|\mathcal{F}_0]$ is a real-valued function on $Dom(\mathcal{E})$. In the rest of the paper, we will substitute $\mathcal{E}[\cdot]$ for $\mathcal{E}[\cdot|\mathcal{F}_0]$.

Remark 4. Our definition of **F**-expectations is similar to that of \mathcal{F}_t^X -consistent non-linear expectations introduced in (Peng, 2004, page 4).

Example 5. The following pairs satisfy (A1)-(A4); thus they are **F**-expectations:

- (1) $(\{E[\cdot|\mathcal{F}_t]\}_{t \in [0, T]}, L^1(\mathcal{F}_T))$: the linear expectation E is a special **F**-expectation with domain $L^1(\mathcal{F}_T)$;
- (2) $(\{\mathcal{E}_g[\cdot|\mathcal{F}_t]\}_{t \in [0, T]}, L^2(\mathcal{F}_T))$: the g -expectation with generator $g(t, z)$ Lipschitz in z (see [\(Peng-97\)](#), [\(CHMP\)](#) or Subsection 7.3 of the present paper);
- (3) $(\{\mathcal{E}_g[\cdot|\mathcal{F}_t]\}_{t \in [0, T]}, L^{\infty}(\mathcal{F}_T))$: the g -expectation with generator $g(t, z)$ having quadratic growth in z (see Subsection 7.4 of this paper).

\mathbf{F} -expectations can alternatively be introduced in a more classical way:

Proposition 6. *Let $\mathcal{E}^\circ: \Lambda \mapsto \mathbb{R}$ be a mapping on some $\Lambda \in \mathcal{D}_T$ satisfying:*

a1 For any $\xi, \eta \in \Lambda$ with $\xi \leq \eta$, a.s., we have $\mathcal{E}^\circ[\xi] \leq \mathcal{E}^\circ[\eta]$. Moreover, if $\mathcal{E}^\circ[\xi] = \mathcal{E}^\circ[\eta]$, then $\xi = \eta$, a.s.;

a2 For any $\xi \in \Lambda$ and $t \in [0, T]$, there exists a unique random variable $\xi_t \in \Lambda_t$ such that $\mathcal{E}^\circ[\mathbf{1}_A \xi + \gamma] = \mathcal{E}^\circ[\mathbf{1}_A \xi_t + \gamma]$ holds for any $A \in \mathcal{F}_t$ and $\gamma \in \Lambda_t$.

Then $\{\mathcal{E}^\circ[\xi|\mathcal{F}_t] \stackrel{\Delta}{=} \xi_t, \xi \in \Lambda\}_{t \in [0, T]}$ defines an \mathbf{F} -expectation with domain Λ .

Remark 7. *For a mapping \mathcal{E}° on some $\Lambda \in \mathcal{D}_T$ satisfying (a1) and (a2), the implied operator $\mathcal{E}^\circ[\cdot|\mathcal{F}_0]$ is also from Λ to \mathbb{R} , which, however, may not equal to \mathcal{E}° . In fact, one can only deduce that $\mathcal{E}^\circ[\xi] = \mathcal{E}^\circ[\mathcal{E}^\circ[\xi|\mathcal{F}_0]]$ for any $\xi \in \Lambda$.*

From now on, when we say an \mathbf{F} -expectation \mathcal{E} , we will refer to the pair $(\mathcal{E}, \text{Dom}(\mathcal{E}))$. Besides (A1)-(A4), the \mathbf{F} -expectation \mathcal{E} has the following properties:

Proposition 8. *For any $\xi, \eta \in \text{Dom}(\mathcal{E})$ and $t \in [0, T]$, we have*

(1) "Local Property": $\mathcal{E}[\mathbf{1}_A \xi + \mathbf{1}_{A^c} \eta|\mathcal{F}_t] = \mathbf{1}_A \mathcal{E}[\xi|\mathcal{F}_t] + \mathbf{1}_{A^c} \mathcal{E}[\eta|\mathcal{F}_t]$, a.s. for any $A \in \mathcal{F}_t$;

(2) "Constant-Preserving": $\mathcal{E}[\xi|\mathcal{F}_t] = \xi$, a.s. if $\xi \in \text{Dom}_t(\mathcal{E})$;

(3) "Comparison": Let $\xi, \eta \in L^0(\mathcal{F}_\nu)$ for some $\nu \in \mathcal{S}_{0, T}$. If $\eta \geq c$, a.s. for some $c \in \mathbb{R}$, then $\xi \leq (\text{hbox|or}) \eta$, a.s. if and only if $\mathcal{E}[\mathbf{1}_A \xi] \leq (\text{hbox|or}) \mathcal{E}[\mathbf{1}_A \eta]$ for all $A \in \mathcal{F}_\nu$.

The following two subsets of $\text{Dom}(\mathcal{E})$ will be of interest:

$$\text{Dom}^+(\mathcal{E}) \stackrel{\Delta}{=} \{\xi \in \text{Dom}(\mathcal{E}): \xi \geq 0, \text{ a.s.}\}, \quad \text{Dom}^\#(\mathcal{E}) \stackrel{\Delta}{=} \{\xi \in \text{Dom}(\mathcal{E}): \xi \geq c, \text{ a.s. } (\text{hbox|for some}) c = c(\xi) \in \mathbb{R}\}. \quad (2.1)$$

Remark 9. *The restrictions of \mathcal{E} on $\text{Dom}^+(\mathcal{E})$ and on $\text{Dom}^\#(\mathcal{E})$, namely $(\mathcal{E}, \text{Dom}^+(\mathcal{E}))$ and $(\mathcal{E}, \text{Dom}^\#(\mathcal{E}))$ respectively, are both \mathbf{F} -expectations: To see this, first note that $\text{Dom}^+(\mathcal{E})$ and $\text{Dom}^\#(\mathcal{E})$ both belong to \mathcal{D}_T . For any $t \in [0, T]$, (A1) and Proposition 8 (2) imply that for any $\xi \in \text{Dom}^\#(\mathcal{E})$*

$$\mathcal{E}[\xi|\mathcal{F}_t] \geq \mathcal{E}[c(\xi)|\mathcal{F}_t] = c(\xi), \quad \text{a.s.}, \quad (\text{hbox|thus}) \quad \mathcal{E}[\xi|\mathcal{F}_t] \in \text{Dom}^\#(\mathcal{E}),$$

which shows that $\mathcal{E}[\cdot|\mathcal{F}_t]$ maps $\text{Dom}^\#(\mathcal{E})$ into $\text{Dom}^\#(\mathcal{E}) \cap L^0(\mathcal{F}_t)$. Then it is easy to check that the restriction of $\mathcal{E} = \{\mathcal{E}[\cdot|\mathcal{F}_t]\}_{t \in [0, T]}$ on $\text{Dom}^\#(\mathcal{E})$ satisfies (A1) to (A4), thus it is an \mathbf{F} -expectation. Similarly, $(\mathcal{E}, \text{Dom}^+(\mathcal{E}))$

is also an \mathbf{F} -expectation. Moreover, the restrictions of \mathcal{E} on $\text{Dom}^-(\mathcal{E}) \stackrel{\Delta}{=} \{\xi \in \text{Dom}(\mathcal{E}) \mid \xi \leq 0, \text{ a.s.}\}$ and on $\text{Dom}_\flat(\mathcal{E}) \stackrel{\Delta}{=} \{\xi \in \text{Dom}(\mathcal{E}) \mid \xi \leq c, \text{ a.s. } (\text{hbox|for some}) c = c(\xi) \in \mathbb{R}\}$ are \mathbf{F} -expectations as well. (fi)

We should remark that restricting \mathcal{E} on any subset Λ' of $\text{Dom}(\mathcal{E})$, with $\Lambda' \in \mathcal{D}_T$, may not result in an \mathbf{F} -expectation, i.e. (\mathcal{E}, Λ') may not be an \mathbf{F} -expectation.

Definition 10. (1) An F -adapted process $X = \{X_t\}_{t \in [0, T]}$ is called an “ \mathcal{E} -process” if $X_t \in \text{Dom}(\mathcal{E})$ for any $t \in [0, T]$;
 (2) An \mathcal{E} -process X is said to be an \mathcal{E} -supermartingale (resp. \mathcal{E} -martingale, \mathcal{E} -submartingale) if for any $0 \leq s < t \leq T$, $\mathcal{E}[X_t | \mathcal{F}_s] \leq (\text{hbox|resp.} =, \geq) X_s$, a.s.

Given a $\nu \in \mathcal{S}_{0, T}^F$ taking values in a finite set $\{t_1 < \dots < t_n\}$, if X is an \mathcal{E} -process, (D2) implies that $X_\nu = \sum_{i=1}^n \mathbf{1}_{\{\nu=t_i\}} X_{t_i} \in \text{Dom}(\mathcal{E})$, thus $X_\nu \in \text{Dom}_\nu(\mathcal{E})$. Since $\{X_t^\xi \triangleq \mathcal{E}[\xi | \mathcal{F}_t]\}_{t \in [0, T]}$ is an \mathcal{E} -process for any $\xi \in \text{Dom}(\mathcal{E})$, we can define an operator $\mathcal{E}[\cdot | \mathcal{F}_\nu]$ from $\text{Dom}(\mathcal{E})$ to $\text{Dom}_\nu(\mathcal{E})$ by

$$\mathcal{E}[\xi | \mathcal{F}_\nu] \triangleq X_\nu^\xi, \quad \langle \text{hbox|for any} \rangle \xi \in \text{Dom}(\mathcal{E}),$$

which allows us to state a basic Optional Sampling Theorem for \mathcal{E} .

Proposition 11. (Optional Sampling Theorem) Let X be an \mathcal{E} -supermartingale (resp. \mathcal{E} -martingale, \mathcal{E} -submartingale). Then for any $\nu, \sigma \in \mathcal{S}_{0, T}^F$, $\mathcal{E}[X_\nu | \mathcal{F}_\sigma] \leq (\text{hbox|resp.} =, \geq) X_{\nu \wedge \sigma}$, a.s.

In particular, applying Proposition 11 to each \mathcal{E} -martingale $\{\mathcal{E}[\xi | \mathcal{F}_t]\}_{t \in [0, T]}$, in which $\xi \in \text{Dom}(\mathcal{E})$, yields the following result.

Corollary 12. For any $\xi \in \text{Dom}(\mathcal{E})$ and $\nu, \sigma \in \mathcal{S}_{0, T}^F$, $\mathcal{E}[\mathcal{E}[\xi | \mathcal{F}_\nu] | \mathcal{F}_\sigma] = \mathcal{E}[\xi | \mathcal{F}_{\nu \wedge \sigma}]$, a.s.

Remark 13. Corollary 12 extends the “Time-Consistency” (A2) to the case of finite-valued stopping times.

$\mathcal{E}[\cdot | \mathcal{F}_\nu]$ inherits other properties of $\mathcal{E}[\cdot | \mathcal{F}_t]$ as well:

Proposition 14. For any $\xi, \eta \in \text{Dom}(\mathcal{E})$ and $\nu \in \mathcal{S}_{0, T}^F$, it holds that

- (1) “Monotonicity (positively strict)”: $\mathcal{E}[\xi | \mathcal{F}_\nu] \leq \mathcal{E}[\eta | \mathcal{F}_\nu]$, a.s. if $\xi \leq \eta$, a.s.; Moreover, if $0 \leq \xi \leq \eta$, a.s. and $\mathcal{E}[\xi | \mathcal{F}_\sigma] = \mathcal{E}[\eta | \mathcal{F}_\sigma]$, a.s. for some $\sigma \in \mathcal{S}_{0, T}^F$, then $\xi = \eta$, a.s.;
- (2) “Zero-one Law”: $\mathcal{E}[\mathbf{1}_A \xi | \mathcal{F}_\nu] = \mathbf{1}_A \mathcal{E}[\xi | \mathcal{F}_\nu]$, a.s. for any $A \in \mathcal{F}_\nu$;
- (3) “Translation Invariance”: $\mathcal{E}[\xi + \eta | \mathcal{F}_\nu] = \mathcal{E}[\xi | \mathcal{F}_\nu] + \eta$, a.s. if $\eta \in \text{Dom}_\nu(\mathcal{E})$;
- (4) “Local Property”: $\mathcal{E}[\mathbf{1}_A \xi + \mathbf{1}_{A^c} \eta | \mathcal{F}_\nu] = \mathbf{1}_A \mathcal{E}[\xi | \mathcal{F}_\nu] + \mathbf{1}_{A^c} \mathcal{E}[\eta | \mathcal{F}_\nu]$, a.s. for any $A \in \mathcal{F}_\nu$;
- (5) “Constant-Preserving”: $\mathcal{E}[\xi | \mathcal{F}_\nu] = \xi$, a.s., $\langle \text{hbox|if} \rangle \xi \in \text{Dom}_\nu(\mathcal{E})$.

We make the following basic hypotheses on the F -expectation \mathcal{E} . These hypotheses will be essential in developing Fatou’s lemma, the Dominated Convergence Theorem and the Upcrossing Theorem.

Hypotheses

- (H0) For any $A \in \mathcal{F}_T$ with $P(A) > 0$, we have $\lim_{n \rightarrow \infty} \mathcal{E}[n \mathbf{1}_A] = \infty$;
- (H1) For any $\xi \in \text{Dom}^+(\mathcal{E})$ and any $\{A_n\}_{n \in \mathbb{N}} \subset \mathcal{F}_T$ with $\lim_{n \rightarrow \infty} \neg \uparrow \mathbf{1}_{A_n} = 1$, a.s., we have $\lim_{n \rightarrow \infty} \neg \uparrow \mathcal{E}[\mathbf{1}_{A_n} \xi] = \mathcal{E}[\xi]$;
- (H2) For any $\xi, \eta \in \text{Dom}^+(\mathcal{E})$ and any $\{A_n\}_{n \in \mathbb{N}} \subset \mathcal{F}_T$ with $\lim_{n \rightarrow \infty} \neg \downarrow \mathbf{1}_{A_n} = 0$, a.s., we have $\lim_{n \rightarrow \infty} \neg \downarrow \mathcal{E}[\xi + \mathbf{1}_{A_n} \eta] = \mathcal{E}[\xi]$.

Remark 15. The linear expectation E on $L^1(\mathcal{F}_T)$ clearly satisfies (H0)-(H2). We will show that Lipschitz and quadratic g -expectations also satisfy (H0)-(H2) in Propositions 72 and 81 respectively.

The \mathbf{F} -expectation \mathcal{E} satisfies the following Fatou's Lemma and Dominated Convergence Theorem.

Theorem 16. (Fatou's Lemma) (H1) is equivalent to the lower semi-continuity of \mathcal{E} : If a sequence $\{\xi_n\}_{n \in \mathbb{N}} \subset \text{Dom}^+(\mathcal{E})$ converges a.s. to some $\xi \in \text{Dom}^+(\mathcal{E})$, then for any $\nu \in \mathcal{S}_{0,T}^F$, we have

$$\mathcal{E}[\xi | \mathcal{F}_\nu] \leq \liminf_{n \rightarrow \infty} \mathcal{E}[\xi_n | \mathcal{F}_\nu], \quad \text{a.s.}, \quad (2.2)$$

where the right hand side of (2.2) could be equal to infinity with non-zero probability.

Remark 17. In the case of the linear expectation E , a converse to (2.2) holds: For any non-negative sequence $\{\xi_n\}_{n \in \mathbb{N}} \subset L^1(\mathcal{F}_T)$ that converges a.s. to some $\xi \in L^0(\mathcal{F}_T)$, if $\liminf_{n \rightarrow \infty} E[\xi_n] < \infty$, then $\xi \in L^1(\mathcal{F}_T)$. However, this statement may not be the case for an arbitrary \mathbf{F} -expectation. That is, $\liminf_{n \rightarrow \infty} \mathcal{E}[\xi_n] < \infty$ may not imply that $\xi \in \text{Dom}^+(\mathcal{E})$ given that $\{\xi_n\}_{n \in \mathbb{N}} \subset \text{Dom}^+(\mathcal{E})$ is a sequence convergent a.s. to some $\xi \in L^0(\mathcal{F}_T)$. See Example 74 for a counterexample in the case of a Lipschitz g -expectation.

Theorem 18. (Dominated Convergence Theorem) Assume (H1) and (H2) hold. Let $\{\xi_n\}_{n \in \mathbb{N}}$ be a sequence in $\text{Dom}^+(\mathcal{E})$ that converges a.s. If there is an $\eta \in \text{Dom}^+(\mathcal{E})$ such that $\xi_n \leq \eta$ a.s. for any $n \in \mathbb{N}$, then the limit ξ of $\{\xi_n\}_{n \in \mathbb{N}}$ belongs to $\text{Dom}^+(\mathcal{E})$, and for any $\nu \in \mathcal{S}_{0,T}^F$, we have

$$\lim_{n \rightarrow \infty} \mathcal{E}[\xi_n | \mathcal{F}_\nu] = \mathcal{E}[\xi | \mathcal{F}_\nu], \quad \text{a.s.}$$

Next, we will derive an Upcrossing Theorem for \mathcal{E} -supermartingales, which is crucial in obtaining an RCLL (right-continuous, with limits from the left) modification for the process $\{\mathcal{E}[\xi | \mathcal{F}_t]\}_{t \in [0,T]}$ as long as $\xi \in \text{Dom}(\mathcal{E})$ is bounded from below. Obtaining a right continuous modification is crucial, since otherwise the conditional expectation $\mathcal{E}[\xi | \mathcal{F}_\nu]$ may not be well defined for any $\nu \in \mathcal{S}_{0,T}$.

Let us first recall what the “number of upcrossings” is: Given a real-valued process $\{X_t\}_{t \in [0, T]}$ and two real numbers $a < b$, for any finite subset F of $[0, T]$, we can define the “number of upcrossings” $U_F(a, b; X(\omega))$ of the interval $[a, b]$ by the sample path $\{X_t(\omega)\}_{t \in F}$ as follows: Set $\nu_0 = -1$, and for any $j = 1, 2, \dots$ we recursively define

$$\begin{aligned}\nu_{2j-1}(\omega) &\triangleq \min \{t \in F: t > \nu_{2j-2}(\omega), X_t(\omega) < a\} \wedge T \in \mathcal{S}_{0, T}^F, \\ \nu_{2j}(\omega) &\triangleq \min \{t \in F: t > \nu_{2j-1}(\omega), X_t(\omega) > b\} \wedge T \in \mathcal{S}_{0, T}^F,\end{aligned}$$

with the convention that $\min \emptyset = \infty$. Then $U_F(a, b; X(\omega))$ is defined to be the largest integer j for which $\nu_{2j}(\omega) < T$. If $I \subset [0, T]$ is not a finite set, we define

$$U_I(a, b; X(\omega)) \triangleq \sup \{U_F(a, b; X(\omega)): F \text{ is a finite subset of } I\}.$$

It will be convenient to introduce a subcollection of \mathcal{S}_T

$$\tilde{\mathcal{S}}_T \triangleq \{\Lambda \in \mathcal{S}_T: \mathbb{R} \subset \Lambda\}.$$

Clearly, $\tilde{\mathcal{S}}_T$ contains all $L^p(\mathcal{F}_T)$, $0 \leq p \leq \infty$. In particular, $L^\infty(\mathcal{F}_T)$ is the smallest element of $\tilde{\mathcal{S}}_T$ in the following sense:

Lemma 19. *For each $\Lambda \in \tilde{\mathcal{S}}_T$, $L^\infty(\mathcal{F}_T) \subset \Lambda$.*

Proof: For any $\xi \in L^\infty(\mathcal{F}_T)$, we have $-\|\xi\|_\infty, 2\|\xi\|_\infty \in \mathbb{R} \subset \Lambda$. Since $0 \leq \xi + \|\xi\|_\infty \leq 2\|\xi\|_\infty$, a.s., (D3) implies that $\xi + \|\xi\|_\infty \in \Lambda$. Then we can deduce from (D2) that $\xi = (\xi + \|\xi\|_\infty) + (-\|\xi\|_\infty) \in \Lambda$. \square

For any \mathbf{F} -adapted process X , we define its right-limit process by

$$X_t^+ \triangleq \liminf_{n \rightarrow \infty} X_{q_n^+(t)}, \quad \text{for any } t \in [0, T],$$

where $q_n^+(t)$ is defined in (1.9). Since the filtration \mathbf{F} is right-continuous, we see that X^+ is an \mathbf{F} -adapted process.

It is now the time to present our Upcrossing Theorem for \mathcal{E} -supermartingales.

Theorem 20. (*Upcrossing Theorem*) *Assume that (H0), (H1) hold and that $\text{Dom}(\mathcal{E}) \in \tilde{\mathcal{S}}_T$. For any \mathcal{E} -supermartingale X , we assume either that $X_T \geq c$, a.s. for some $c \in \mathbb{R}$ or that the operator $\mathcal{E}[\cdot]$ is concave: For any $\xi, \eta \in \text{Dom}(\mathcal{E})$*

$$\mathcal{E}[\lambda\xi + (1-\lambda)\eta] \geq \lambda\mathcal{E}[\xi] + (1-\lambda)\mathcal{E}[\eta], \quad \forall \lambda \in (0, 1). \quad (2.3)$$

Then for any two real numbers $a < b$, it holds that $P(U_{\mathcal{D}_T}(a, b; X) < \infty) = 1$. Thus we have

$$P\left(X_t^+ = \lim_{n \rightarrow \infty} X_{q_n^+(t)} \text{ for any } t \in [0, T]\right) = 1. \quad (2.4)$$

As a result, X^+ is an RCLL process.

In the rest of this section, we assume that the \mathbf{F} -expectation \mathcal{E} satisfies (H0)-(H2) and that $Dom(\mathcal{E}) \in \tilde{\mathcal{D}}_T$. The following proposition will play a fundamental role throughout this paper.

Proposition 21. *Let X be a non-negative \mathcal{E} -supermartingale. (1) Assume either that $\text{esssup}_{t \in \mathcal{D}_T} X_t \in Dom^+(\mathcal{E})$ or that for any sequence $\{\xi_n\}_{n \in \mathbb{N}} \subset Dom^+(\mathcal{E})$ convergent a.s. to some $\xi \in L^0(\mathcal{F}_T)$,*

$$\langle \liminf_{n \rightarrow \infty} \mathcal{E}[\xi_n] < \infty \text{ implies } \xi \in Dom^+(\mathcal{E}). \rangle \quad (2.5)$$

Then for any $\nu \in \mathcal{S}_{0,T}$, X_ν^+ belongs to $Dom^+(\mathcal{E})$;

(2) If $X_t^+ \in Dom^+(\mathcal{E})$ for any $t \in [0, T]$, then X^+ is an RCLL \mathcal{E} -supermartingale such that for any $t \in [0, T]$, $X_t^+ \leq X_t$, a.s.;

(3) Moreover, if the function $t \mapsto \mathcal{E}[X_t]$ from $[0, T]$ to \mathbb{R} is right continuous, then X^+ is an RCLL modification of X . Conversely, if X has a right-continuous modification, then the function $t \mapsto \mathcal{E}[X_t]$ is right continuous.

Now we add one more hypothesis to the \mathbf{F} -expectation \mathcal{E} :

(H3) For any $\xi \in Dom^+(\mathcal{E})$ and $\nu \in \mathcal{S}_{0,T}$, $X_\nu^{\xi,+} \in Dom^+(\mathcal{E})$.

In light of Proposition 21 (1), (H3) holds if $\text{esssup}_{t \in \mathcal{D}_T} \mathcal{E}[\xi | \mathcal{F}_t] \in Dom^+(\mathcal{E})$ or if \mathcal{E} satisfies (2.5).

For each $\xi \in Dom^\#(\mathcal{E})$, we define $\xi' \triangleq \xi - c(\xi) \in Dom^+(\mathcal{E})$. Clearly $X^{\xi'} \triangleq \{\mathcal{E}[\xi' | \mathcal{F}_t]\}_{t \in [0, T]}$ is a non-negative \mathcal{E} -martingale. By (A2), $\mathcal{E}[X_t^{\xi'}] = \mathcal{E}[\mathcal{E}[\xi' | \mathcal{F}_t]] = \mathcal{E}[\xi']$ for any $t \in [0, T]$, which means that $t \mapsto \mathcal{E}[X_t^{\xi'}]$ is continuous function on $[0, T]$. Thanks to Proposition 21 (2) and (H3), the process $X_t^{\xi',+} \triangleq \liminf_{n \rightarrow \infty} X_{q_n^+(t)}^{\xi'}$, $t \in [0, T]$ is an RCLL modification of $X^{\xi'}$. Then for any $\nu \in \mathcal{S}_{0,T}$, we define

$$\tilde{\mathcal{E}}[\xi | \mathcal{F}_\nu] \triangleq X_\nu^{\xi',+} - c(\xi) \quad (2.6)$$

as the conditional \mathbf{F} -expectation of ξ at the stopping time $\nu \in \mathcal{S}_{0,T}$. Since we have assumed $Dom(\mathcal{E}) \in \tilde{\mathcal{D}}_T$, Lemma 19, (H3), (D2) as well as the non-negativity of $X_\nu^{\xi',+}$ imply that

$$\tilde{\mathcal{E}}[\xi | \mathcal{F}_\nu] \in Dom^\#(\mathcal{E}), \quad (2.7)$$

which shows that $\tilde{\mathcal{E}}[\cdot | \mathcal{F}_\nu]$ is an operator from $Dom^\#(\mathcal{E})$ to $Dom^\#(\mathcal{E}) \triangleq Dom^\#(\mathcal{E}) \cap L^0(\mathcal{F}_\nu)$. In fact, $\{\tilde{\mathcal{E}}[\cdot | \mathcal{F}_t]\}_{t \in [0, T]}$ defines a \mathbf{F} -expectation on $Dom^\#(\mathcal{E})$, as the next result shows.

Proposition 22. *For any $\xi \in Dom^\#(\mathcal{E})$, $\tilde{\mathcal{E}}[\xi | \mathcal{F}_\cdot]$ is an RCLL modification of $\mathcal{E}[\xi | \mathcal{F}_\cdot]$. $\{\tilde{\mathcal{E}}[\cdot | \mathcal{F}_t]\}_{t \in [0, T]}$ is an \mathbf{F} -expectation with domain $Dom(\tilde{\mathcal{E}}) = Dom^\#(\mathcal{E}) \in \tilde{\mathcal{D}}_T$ and satisfying (H0)-(H2); thus all preceding results are applicable to $\tilde{\mathcal{E}}$.*

Proof: As $Dom(\mathcal{E}) \in \tilde{\mathcal{D}}_T$ is assumed, we see that $Dom^\#(\mathcal{E})$ also belongs to $\tilde{\mathcal{D}}_T$. Fix $\xi \in Dom^\#(\mathcal{E})$. Since $X^{\xi',+}$ is an RCLL modification of $X^{\xi'}$, (A4) implies that for any $t \in [0, T]$

$$\tilde{\mathcal{E}}[\xi|\mathcal{F}_t] = X_t^{\xi',+} \neg + c(\xi) = \mathcal{E}[\xi'|\mathcal{F}_t] + c(\xi) = \mathcal{E}[\xi' + c(\xi)|\mathcal{F}_t] = \mathcal{E}[\xi|\mathcal{F}_t], \quad a.s. \quad (2.8)$$

Thus $\tilde{\mathcal{E}}[\xi|\mathcal{F}]$ is actually an RCLL modification of $\mathcal{E}[\xi|\mathcal{F}]$. Then it is easy to show that the pair $(\tilde{\mathcal{E}}, Dom^\#(\mathcal{E}))$ satisfies (A1)-(A4) and (H0)-(H2); thus it is an **F**-expectation. \square

We restate Proposition 22 with respect to $\tilde{\mathcal{E}}$ for future use.

Corollary 23. *Let X be an $\tilde{\mathcal{E}}$ -supermartingale such that $\text{essinf}_{t \in [0, T]} X_t \geq c$, a.s. for some $c \in \mathbb{R}$.*

(1) *If $\text{esssup}_{t \in \mathcal{D}_T} X_t \in Dom^\#(\mathcal{E})$ or if (2.5) holds, then X_ν^+ belongs to $Dom^\#(\mathcal{E})$ for any $\nu \in \mathcal{S}_{0, T}$;*

(2) *If $X_t^+ \in Dom^\#(\mathcal{E})$ for any $t \in [0, T]$, then X^+ is an RCLL $\tilde{\mathcal{E}}$ -supermartingale such that for any $t \in [0, T]$, $X_t^+ \leq X_t$, a.s.*

(3) *Moreover, if the function $t \mapsto \tilde{\mathcal{E}}[X_t]$ from $[0, T]$ to \mathbb{R} is right continuous, then X^+ is an RCLL modification of X . Conversely, if X has a right-continuous modification, then the function $t \mapsto \tilde{\mathcal{E}}[X_t]$ is right continuous.*

The next result is the Optional Sampling Theorem of $\tilde{\mathcal{E}}$ for the stopping times in $\mathcal{S}_{0, T}$.

Theorem 24. (Optional Sampling Theorem 2) *Let X be a right-continuous $\tilde{\mathcal{E}}$ -supermartingale (resp. $\tilde{\mathcal{E}}$ -martingale, $\tilde{\mathcal{E}}$ -submartingale) such that $\text{essinf}_{t \in \mathcal{D}_T} X_t \geq c$, a.s. for some $c \in \mathbb{R}$. If $X_\nu \in Dom^\#(\mathcal{E})$ for any $\nu \in \mathcal{S}_{0, T}$, then for any $\nu, \sigma \in \mathcal{S}_{0, T}$, we have*

$$\tilde{\mathcal{E}}[X_\nu|\mathcal{F}_\sigma] \leq (\text{hbox|resp.}) =, \geq) X_{\nu \wedge \sigma}, \quad a.s.$$

Using the Optional Sampling Theorem, we are able to extend Corollary 23 and Proposition 14 to the operators $\tilde{\mathcal{E}}[\cdot|\mathcal{F}_\nu]$, $\nu \in \mathcal{S}_{0, T}$.

Corollary 25. *For any $\xi \in Dom^\#(\mathcal{E})$ and $\nu, \sigma \in \mathcal{S}_{0, T}$, we have*

$$\tilde{\mathcal{E}}[\tilde{\mathcal{E}}[\xi|\mathcal{F}_\nu]|\mathcal{F}_\sigma] = \tilde{\mathcal{E}}[\xi|\mathcal{F}_{\nu \wedge \sigma}], \quad a.s. \quad (2.9)$$

Proof: Since $(\tilde{\mathcal{E}}, Dom^\#(\mathcal{E}))$ is an **F**-expectation by Proposition 22, for any $\xi \in Dom^\#(\mathcal{E})$, (A2) implies that the RCLL process $\tilde{X}^\xi \triangleq \{\tilde{\mathcal{E}}[\xi|\mathcal{F}_t]\}_{t \in [0, T]}$ is an $\tilde{\mathcal{E}}$ -martingale. For any $t \in [0, T]$, (2.8) and Proposition 8 (2) show that

$$\tilde{X}_t^\xi = \tilde{\mathcal{E}}[\xi|\mathcal{F}_t] \geq \tilde{\mathcal{E}}[c(\xi)|\mathcal{F}_t] = \mathcal{E}[c(\xi)|\mathcal{F}_t] = c(\xi), \quad a.s.,$$

which implies that $\operatorname{ess\,inf}_{t \in [0, T]} \tilde{X}_t^\xi \geq c(\xi)$, a.s. Then (??) and Theorem 24 give rise to (??). \square

Proposition 26. *For any $\xi, \eta \in \operatorname{Dom}^\#(\mathcal{E})$ and $\nu \in \mathcal{S}_{0, T}$, it holds that*

- (1) “Strict Monotonicity”: $\tilde{\mathcal{E}}[\xi|\mathcal{F}_\nu] \leq \tilde{\mathcal{E}}[\eta|\mathcal{F}_\nu]$, a.s. if $\xi \leq \eta$, a.s.; Moreover, if $\tilde{\mathcal{E}}[\xi|\mathcal{F}_\sigma] = \tilde{\mathcal{E}}[\eta|\mathcal{F}_\sigma]$, a.s. for some $\sigma \in \mathcal{S}_{0, T}$, then $\xi = \eta$, a.s.;
- (2) “Zero-one Law”: $\tilde{\mathcal{E}}[\mathbf{1}_A \xi|\mathcal{F}_\nu] = \mathbf{1}_A \tilde{\mathcal{E}}[\xi|\mathcal{F}_\nu]$, a.s. for any $A \in \mathcal{F}_\nu$;
- (3) “Translation Invariance”: $\tilde{\mathcal{E}}[\xi + \eta|\mathcal{F}_\nu] = \tilde{\mathcal{E}}[\xi|\mathcal{F}_\nu] + \eta$, a.s. if $\eta \in \operatorname{Dom}_\nu^\#(\mathcal{E})$;
- (4) “Local Property”: $\tilde{\mathcal{E}}[\mathbf{1}_A \xi + \mathbf{1}_{A^c} \eta|\mathcal{F}_\nu] = \mathbf{1}_A \tilde{\mathcal{E}}[\xi|\mathcal{F}_\nu] + \mathbf{1}_{A^c} \tilde{\mathcal{E}}[\eta|\mathcal{F}_\nu]$, a.s. for any $A \in \mathcal{F}_\nu$;
- (5) “Constant-Preserving”: $\tilde{\mathcal{E}}[\xi|\mathcal{F}_\nu] = \xi$, a.s., if $\xi \in \operatorname{Dom}_\nu^\#(\mathcal{E})$.

Remark 27. Corollary 25, Proposition 26 (2) and (2.8) imply that for any $\xi \in \operatorname{Dom}^\#(\mathcal{E})$ and $\nu \in \mathcal{S}_{0, T}$,

$$\mathcal{E}[\mathbf{1}_A \xi] = \tilde{\mathcal{E}}[\mathbf{1}_A \xi] = \tilde{\mathcal{E}}[\tilde{\mathcal{E}}[\mathbf{1}_A \xi|\mathcal{F}_\nu]] = \tilde{\mathcal{E}}[\mathbf{1}_A \tilde{\mathcal{E}}[\xi|\mathcal{F}_\nu]] = \mathcal{E}[\mathbf{1}_A \tilde{\mathcal{E}}[\xi|\mathcal{F}_\nu]], \quad \forall A \in \mathcal{F}_\nu. \quad (2.10)$$

In light of Proposition 8 (3), $\tilde{\mathcal{E}}[\xi|\mathcal{F}_\nu]$ is the unique element (up to a P -null set) in $\operatorname{Dom}_\nu^\#(\mathcal{E})$ that makes (2.10) hold. Therefore, we see that the random variable $\tilde{\mathcal{E}}[\xi|\mathcal{F}_\nu]$ defined by (??) is exactly the conditional \mathbf{F} -expectation of ξ at the stopping time ν in the classical sense.

In light of Corollary 25 and Proposition 26, we can generalize Fatou’s Lemma (Theorem 16) and the Dominated Convergence Theorem (Theorem 18) to the conditional \mathbf{F} -expectation $\tilde{\mathcal{E}}[\cdot|\mathcal{F}_\nu]$, $\nu \in \mathcal{S}_{0, T}$.

Proposition 28. (Fatou’s Lemma 2) *Let $\{\xi_n\}_{n \in \mathbb{N}}$ be a sequence in $\operatorname{Dom}^\#(\mathcal{E})$ that converges a.s. to some $\xi \in \operatorname{Dom}^\#(\mathcal{E})$ and satisfies $\operatorname{ess\,inf}_{n \in \mathbb{N}} \xi_n \geq c$, a.s. for some $c \in \mathbb{R}$, then for any $\nu \in \mathcal{S}_{0, T}$, we have*

$$\tilde{\mathcal{E}}[\xi|\mathcal{F}_\nu] \leq \liminf_{n \rightarrow \infty} \tilde{\mathcal{E}}[\xi_n|\mathcal{F}_\nu], \quad \text{a.s.}, \quad (2.11)$$

where the right hand side of (2.11) could be equal to infinity with non-zero probability.

Proposition 29. (Dominated Convergence Theorem 2) *Let $\{\xi_n\}_{n \in \mathbb{N}}$ be a sequence in $\operatorname{Dom}^\#(\mathcal{E})$ that converges a.s. and that satisfies $\operatorname{ess\,inf}_{n \in \mathbb{N}} \xi_n \geq c$, a.s. for some $c \in \mathbb{R}$. If there is an $\eta \in \operatorname{Dom}^\#(\mathcal{E})$ such that $\xi_n \leq \eta$ a.s. for any $n \in \mathbb{N}$, then the limit ξ of $\{\xi_n\}_{n \in \mathbb{N}}$ belongs to $\operatorname{Dom}^\#(\mathcal{E})$ and for any $\nu \in \mathcal{S}_{0, T}$, we have*

$$\lim_{n \rightarrow \infty} \tilde{\mathcal{E}}[\xi_n|\mathcal{F}_\nu] = \tilde{\mathcal{E}}[\xi|\mathcal{F}_\nu], \quad \text{a.s.} \quad (2.12)$$

Proof of Propositions 28 and 29: In the proofs of Theorem 16 and Theorem 18, we only need to replace $\{\xi_n\}_{n \in \mathbb{N}}$ and $\mathcal{E}[\cdot|\mathcal{F}_t]$ by $\{\xi_n - c\}_{n \in \mathbb{N}}$ and $\tilde{\mathcal{E}}[\cdot|\mathcal{F}_\nu]$ respectively. Instead of (A1), (A3) and (A4), we apply Proposition 26 (1)-(3). Moreover, since (A2) is only used on $Dom^+(\mathcal{E})$ in the proofs of Theorem 16 and Theorem 18, we can substitute Corollary 25 for it. Eventually, a simple application of Proposition 26 (3) yields (2.11) and (2.12). \square

3 Collections of \mathbf{F} -Expectations

In this section, we will show that *pasting* of two \mathbf{F} -expectations at a given stopping time is itself an \mathbf{F} -expectation. Moreover, pasting preserves (H1) and (H2). We will then introduce the concept of a *stable* class of \mathbf{F} -expectations, which are collections closed under pasting. We will solve the optimal stopping problems introduced in (1.1) and (1.6) over this class of \mathbf{F} -expectations. Before we show the pasting property of \mathbf{F} -expectations, we introduce the concept of convexity for an \mathbf{F} -expectation and give one of the consequences of having convexity:

Definition 30. An \mathbf{F} -expectation \mathcal{E} is called “positively-convex” if for any $\xi, \eta \in Dom^+(\mathcal{E})$, $\lambda \in (0, 1)$ and $t \in [0, T]$

$$\mathcal{E}[\lambda\xi + (1 - \lambda)\eta|\mathcal{F}_t] \leq \lambda\mathcal{E}[\xi|\mathcal{F}_t] + (1 - \lambda)\mathcal{E}[\eta|\mathcal{F}_t], \quad a. s.$$

Lemma 31. Any positively-convex \mathbf{F} -expectation satisfies (H0). Moreover, an \mathbf{F} -expectation \mathcal{E} is positively-convex if and only if the implied \mathbf{F} -expectation $(\tilde{\mathcal{E}}, Dom^\#(\mathcal{E}))$ is convex, i.e., for any $\xi, \eta \in Dom^\#(\mathcal{E})$, $\lambda \in (0, 1)$ and $t \in [0, T]$

$$\tilde{\mathcal{E}}[\lambda\xi + (1 - \lambda)\eta|\mathcal{F}_t] \leq \lambda\tilde{\mathcal{E}}[\xi|\mathcal{F}_t] + (1 - \lambda)\tilde{\mathcal{E}}[\eta|\mathcal{F}_t], \quad a. s. \quad (3.1)$$

Proposition 32. Let $\mathcal{E}_i, \mathcal{E}_j$ be two \mathbf{F} -expectations with the same domain $\Lambda \in \tilde{\mathcal{D}}_T$ and satisfying (H1)-(H3). For any $\nu \in \mathcal{S}_{0,T}$, we define the pasting of $\mathcal{E}_i, \mathcal{E}_j$ at the stopping time ν to be the following RCLL \mathbf{F} -adapted process

$$\mathcal{E}_{i,j}^\nu[\xi|\mathcal{F}_t] \triangleq \mathbf{1}_{\{\nu \leq t\}} \tilde{\mathcal{E}}_j[\xi|\mathcal{F}_t] + \mathbf{1}_{\{\nu > t\}} \tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_j[\xi|\mathcal{F}_\nu]|\mathcal{F}_t], \quad \forall t \in [0, T] \quad (3.2)$$

for any $\xi \in \Lambda^\# = \{\xi \in \Lambda : \xi \geq c, a. s. \text{ for some } c = c(\xi) \in \mathbb{R}\}$. Then $\mathcal{E}_{i,j}^\nu$ is an \mathbf{F} -expectation with domain $\Lambda^\# \in \tilde{\mathcal{D}}_T$ and satisfying (H1) and (H2). Moreover, if \mathcal{E}_i and \mathcal{E}_j are both positively-convex, $\mathcal{E}_{i,j}^\nu$ is convex in the sense of (?).

In particular, for any $\sigma \in \mathcal{S}_{0,T}$, applying Proposition 26 (4) and (5), we obtain

$$\begin{aligned} \mathcal{E}_{i,j}^\nu[\xi|\mathcal{F}_\sigma] &\neg\neg = \neg\neg \mathbf{1}_{\{\nu \leq \sigma\}} \tilde{\mathcal{E}}_j[\xi|\mathcal{F}_\sigma] + \mathbf{1}_{\{\nu > \sigma\}} \tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_j[\xi|\mathcal{F}_\nu]|\mathcal{F}_\sigma] = \\ &\mathbf{1}_{\{\nu \leq \sigma\}} \tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_j[\xi|\mathcal{F}_\sigma]|\mathcal{F}_\sigma] + \mathbf{1}_{\{\nu > \sigma\}} \tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_j[\xi|\mathcal{F}_\nu]|\mathcal{F}_\sigma] \\ &\neg\neg = \neg\neg \tilde{\mathcal{E}}_i[\mathbf{1}_{\{\nu \leq \sigma\}} \tilde{\mathcal{E}}_j[\xi|\mathcal{F}_\sigma] + \mathbf{1}_{\{\nu > \sigma\}} \tilde{\mathcal{E}}_j[\xi|\mathcal{F}_\nu]|\mathcal{F}_\sigma] = \\ &\tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_j[\xi|\mathcal{F}_{\nu \vee \sigma}]|\mathcal{F}_\sigma], \quad a.s., \end{aligned} \quad (3.3)$$

where we used the fact that $\{\nu > \sigma\} \in \mathcal{F}_{\nu \wedge \sigma}$ thanks to (, ?, Lemma 1.2.16).

Remark 33. *Pasting may not preserve (H0). From now on, we will replace the (H0) assumption by the positive convexity, which implies the former and which is an invariant property under pasting thanks to the previous two results. Positive convexity is also important in constructing an optimal stopping time of (1.1) (see Theorem 57).*

All of the ingredients are in place to introduce what we mean by a stable class of \mathbf{F} -expectations. As we will see in Lemma 31, stability assures that the essential supremum or infimum over the class can be approximated by an increasing or decreasing sequence in the class.

Definition 34. *A class $\mathcal{E} = \{\mathcal{E}_i\}_{i \in \mathcal{I}}$ of \mathbf{F} -expectations is said to be “stable” if*

- (1) *All \mathcal{E}_i , $i \in \mathcal{I}$ are positively-convex \mathbf{F} -expectations with the same domain $\Lambda \in \tilde{\mathcal{H}}_T$ and they satisfy H 1-H 3;*

- (2) *\mathcal{E} is closed under pasting: namely, for any $i, j \in \mathcal{I}$, $\nu \in \mathcal{S}_{0,T}$, there exists a $k = k(i, j, \nu) \in \mathcal{I}$ such that $\mathcal{E}_{i,j}^\nu$ coincides with $\tilde{\mathcal{E}}_k$ on $\Lambda^\#$.*

We shall denote $\text{Dom}(\mathcal{E}) \stackrel{\Delta}{=} \Lambda^\#$, thus $\text{Dom}(\mathcal{E}) = \text{Dom}^\#(\mathcal{E}_i) \in \tilde{\mathcal{H}}_T$ for any $i \in \mathcal{I}$. Moreover, if $\mathcal{E}' = \{\mathcal{E}_i\}_{i \in \mathcal{I}'}$ satisfies (2) for some non-empty subset \mathcal{I}' of \mathcal{I} , then we call \mathcal{E}' a stable subclass of \mathcal{E} , clearly $\text{Dom}(\mathcal{E}') = \text{Dom}(\mathcal{E})$.

Remark 35. *The notion of “pasting” for linear expectations was given by (, ?, Definition 6.41). The counterpart of Proposition 3.3 for the linear expectations, which states that pasting two probability measures equivalent to P results in another probability measure equivalent to P , is given by (, ?, Lemma 6.43). Note that in the case of linear expectations, (H1), (H2) and the convexity are trivially preserved because pasting in that case gives us a linear expectation. On the other hand, the notion of stability for linear expectations was given by (, ?, Definition 6.44). The stability is also referred to as “fork convexity” in stochastic control theory, “m-stability” in stochastic analysis or “rectangularity” in decision theory (see the introduction of [\[citet|Delbaen_2006\]](#) and [\[citet|Bion_2009\]](#) for details).*

Example 36. (1) *Let \mathcal{P} be the set of all probability measures equivalent to P , then $\mathcal{E}_{\mathcal{P}} \stackrel{\Delta}{=} \{E_Q\}_{Q \in \mathcal{P}}$ is a stable class of linear expectations; see (, ?, Proposition 6.45).*

(2) Consider a collection \mathfrak{U} of admissible control processes. For any $U \in \mathfrak{U}$, let P^U be the equivalent probability measure defined via (, ?, (5)) (or (, ?, (2.5))), then $\mathcal{E}_{\mathfrak{U}} \triangleq \{E_{P^U}\}_{U \in \mathfrak{U}}$ is a stable class of linear expectations; see Subsection 7.3 of the present paper.

(3) For any $M > 0$, a family \mathcal{E}_M of convex Lipschitz g -expectations with Lipschitz coefficient $K_g \leq M$ is an example of stable class of non-linear expectations; see Subsection 7.3 of this paper.

The following Lemma gives us a tool for checking whether a random variable is inside the domain $Dom(\mathcal{E})$ of a stable class \mathcal{E} .

Lemma 37. *Given a stable class \mathcal{E} of \mathbf{F} -expectations, a random variable ξ belongs to $Dom(\mathcal{E})$ if and only if $c \leq \xi \leq \eta$, a.s. for some $c \in \mathbb{R}$ and $\eta \in Dom(\mathcal{E})$.*

Proof: Consider a random variable ξ . If $\xi \in Dom(\mathcal{E})$, since $Dom(\mathcal{E}) = Dom^\#(\mathcal{E}_i)$ for any $i \in \mathcal{I}$, we know that there exists a $c = c(\xi) \in \mathbb{R}$ such that $\xi \geq c(\xi)$, a.s.

On the other hand, if $c \leq \xi \leq \eta$, a.s. for some $c \in \mathbb{R}$ and $\eta \in Dom(\mathcal{E})$, it follows that $0 \leq \xi - c \leq \eta - c$, a.s. Since $Dom(\mathcal{E}) \in \tilde{\mathcal{D}}_T$, we see that $-c, c \in \mathbb{R} \subset Dom(\mathcal{E})$. Then (D2) shows that $\eta - c \in Dom(\mathcal{E})$ and thus (D3) implies that $\xi - c \in Dom(\mathcal{E})$, which further leads to that $\xi = (\xi - c) + c \in Dom(\mathcal{E})$ thanks to (D2) again. \square Moreover, for any $\xi \in L^\infty(\mathcal{F}_T)$, since $-\|\xi\|_\infty \leq \xi \leq \|\xi\|_\infty$, a.s. and since $\|\xi\|_\infty \in \mathbb{R} \subset Dom(\mathcal{E})$, we see from the above argument that $\xi \in Dom(\mathcal{E})$. Hence $L^\infty(\mathcal{F}_T) \subset Dom(\mathcal{E})$, proving the lemma. \square (fi)

We end this section by reviewing some basic properties of the essential supremum and essential infimum (for their definitions, see e.g. (, ?, Proposition VI-11-1), or (, ?, Theorem A.32)).

Lemma 38. *Let $\{\xi_j\}_{j \in \mathcal{J}}$ and $\{\eta_j\}_{j \in \mathcal{J}}$ be two families of random variables of $L^0(\mathcal{F})$ with the same index set \mathcal{J} .*

(1) *If $\xi_j \leq (=) \eta_j$, a.s. for any $j \in \mathcal{J}$, then $\operatorname{esssup}_{j \in \mathcal{J}} \xi_j \leq (=) \operatorname{esssup}_{j \in \mathcal{J}} \eta_j$, a.s.*

(2) *For any $A \in \mathcal{F}$, it holds a.s. that $\operatorname{esssup}_{j \in \mathcal{J}} (11_A \xi_j + 11_{A^c} \eta_j) = 11_A \operatorname{esssup}_{j \in \mathcal{J}} \xi_j + 11_{A^c} \operatorname{esssup}_{j \in \mathcal{J}} \eta_j$; In particular, $\operatorname{esssup}_{j \in \mathcal{J}} (11_A \xi_j) = 11_A \operatorname{esssup}_{j \in \mathcal{J}} \xi_j$, a.s.*

(3) *For any random variable $\gamma \in L^0(\mathcal{F})$ and any $\alpha > 0$, we have $\operatorname{esssup}_{j \in \mathcal{J}} (\alpha \xi_j + \gamma) = \alpha \operatorname{esssup}_{j \in \mathcal{J}} \xi_j + \gamma$, a.s.*

Moreover, (1)-(3) hold when we replace $\operatorname{esssup}_{j \in \mathcal{J}}$ by $\operatorname{essinf}_{j \in \mathcal{J}}$.

4 Optimal Stopping with Multiple Priors

In this section, we will solve an optimal stopping problem in which the objective of the stopper is to determine an optimal stopping time τ^* that satisfies

$$\sup_{(i,\rho) \in \mathcal{I} \times \mathcal{S}_{0,T}} \mathcal{E}_i[Y_\rho + H_\rho^i] = \sup_{i \in \mathcal{I}} \mathcal{E}_i[Y_\tau + H_\tau^i], \quad (4.1)$$

where $\mathcal{E} = \{\mathcal{E}_i\}_{i \in \mathcal{I}}$ is a stable class of \mathbf{F} -expectations, Y is a primary reward process and H^i is a *model*-dependent cumulative reward process. (We will outline the assumptions on the reward processes below.) To find an optimal stopping time, we shall build a so-called “ \mathcal{E} -upper Snell envelope”, which we will denote by Z^0 , of the reward process Y . Namely, Z^0 is the smallest RCLL \mathbf{F} -adapted process dominating Y such that $Z^0 + H^i$ is an $\tilde{\mathcal{E}}_i$ -supermartingale for any $i \in \mathcal{I}$. We will show under certain assumptions that the first time Z^0 meets Y is an optimal stopping time for (4.1).

We start by making some assumptions on the reward processes: Let $\mathcal{E} = \{\mathcal{E}_i\}_{i \in \mathcal{I}}$ be a stable class of \mathbf{F} -expectations accompanied by a family $\mathcal{H} \triangleq \{H^i\}_{i \in \mathcal{I}}$ of right-continuous \mathbf{F} -adapted processes that satisfies

(S1) For any $i \in \mathcal{I}$, $H_0^i = 0$, a.s. and

$$H_{\nu,\rho}^i \triangleq H_\rho^i - H_\nu^i \in \text{Dom}(\mathcal{E}), \quad \forall \nu, \rho \in \mathcal{S}_{0,T} \text{ (hbox|with } h\nu \leq \rho, \text{ a.s.)} \quad (4.2)$$

Moreover, if no member of \mathcal{E} satisfies (2.5), we assume that there exists a $j \in \mathcal{I}$ such that

$$\zeta^j \triangleq \text{esssup}_{s,t \in \mathcal{D}_T; s < t} H_{s,t}^j \in \text{Dom}(\mathcal{E}). \quad (4.3)$$

(S2) There exists a $C_H < 0$ such that for any $i \in \mathcal{I}$, $\text{essinf}_{s,t \in \mathcal{D}_T; s < t} H_{s,t}^i \geq C_H$, a.s.

(S3) For any $\nu \in \mathcal{S}_{0,T}$ and $i, j \in \mathcal{I}$, it holds for any $0 \leq s < t \leq T$ that $H_{s,t}^k = H_{\nu \wedge s, \nu \wedge t}^i + H_{\nu \vee s, \nu \vee t}^j$, a.s., where $k = k(i, j, \nu) \in \mathcal{I}$ is the index defined in Definition 34 (2).

Remark 39. (1) For any $i \in \mathcal{I}$, (S2) and the right-continuity of H^i imply that except on a null set $N(i)$

$$H_{s,t}^i \geq C_H, \quad \text{(hbox|for any } 0 \leq s < t \leq T, \text{ (hbox|thus } H_{\nu,\rho}^i \geq C_H, \\ \forall \nu, \rho \in \mathcal{S}_{0,T} \text{ (hbox|with } h\nu \leq \rho, \text{ a.s.)} \quad (4.4)$$

(2) If (?) is assumed for some $j \in \mathcal{I}$, we can deduce from the right-continuity of H^j and (?) that except on a null set N

$$C_H \leq H_{s,t}^j \leq \zeta^j, \quad \text{(hbox|for any } 0 \leq s < t \leq T, \text{ (hbox|thus } C_H \leq H_{\nu,\rho}^j \leq \zeta^j, \\ \forall \nu, \rho \in \mathcal{S}_{0,T} \text{ (hbox|with } h\nu \leq \rho, \text{ a.s.)}$$

Then Lemma 38 implies that (4.2) holds for j . Hence we see that (?) is a stronger condition than (4.2).

(3) Since H^i , H^j and H^k are all right-continuous processes, (S3) is equivalent to the statement that a.s.

$$H_{s,t}^k = H_{\nu \wedge s, \nu \wedge t}^i + H_{\nu \vee s, \nu \vee t}^j, \quad \forall 0 \leq s < t \leq T. \quad (4.5)$$

Now we give an example of \mathcal{H} .

Lemma 40. Let $\{h^i\}_{i \in \mathcal{I}}$ be a family of progressive processes satisfying the following assumptions:

(h1) For any $i \in \mathcal{I}$ and $\nu, \rho \in \mathcal{S}_{0,T}$ with $\nu \leq \rho$, a.s., $\int_\nu^\rho h_t^i dt \in \text{Dom}(\mathcal{E})$. Moreover, if no member of \mathcal{E} satisfies

(4.3), we assume that there exists a $j \in \mathcal{I}$ such that $\int_0^T |h_t^j| dt \in \text{Dom}(\mathcal{E})$.

(h2) There exists a $c < 0$ such that for any $i \in \mathcal{I}$, $h_t^i \geq c$, $dt \times dP$ -a.s.

(h3) For any $\nu \in \mathcal{S}_{0,T}$ and $i, j \in \mathcal{I}$, it holds for any $t \in [0, T]$ that $h_t^k = \mathbf{1}_{\{\nu \leq t\}} h_t^j + \mathbf{1}_{\{\nu > t\}} h_t^i$, $dt \times dP$ -a.s., where

$k = k(i, j, \nu) \in \mathcal{I}$ is the index defined in Definition 34 (2).

Then $\left\{ H_t^i \triangleq \int_0^t h_s^i ds, t \in [0, T] \right\}_{i \in \mathcal{I}}$ is a family of right-continuous \mathbf{F} -adapted processes satisfying (S1)-(S3).

Standing assumptions on Y in this section.

Let Y be a right-continuous \mathbf{F} -adapted process that satisfies

(Y1) For any $\nu \in \mathcal{S}_{0,T}$, $Y_\nu \in \text{Dom}(\mathcal{E})$.

(Y2) $\sup_{(i, \rho) \in \mathcal{I} \times \mathcal{S}_{0,T}} \mathcal{E}_i[Y_\rho^i] < \infty$, where $Y^i \triangleq \{Y_t + H_t^i\}_{t \in [0, T]}$. Moreover, if no member of \mathcal{E} satisfies (4.3), then we

assume that

$$\zeta_Y \triangleq \text{esssup}_{(i, \rho, t) \in \mathcal{I} \times \mathcal{S}_{0,T} \times \mathcal{D}_T} \tilde{\mathcal{E}}_i[Y_\rho^i | \mathcal{F}_t] \in \text{Dom}(\mathcal{E}). \quad (4.6)$$

(Y3) $\text{essinf}_{t \in \mathcal{D}_T} Y_t \geq C_Y$, a.s. for some $C_Y < 0$.

Remark 41. (1) For any $i \in \mathcal{I}$, (A4) and (2.8) imply that \mathcal{E}_i satisfies (4.3) if and only if $\tilde{\mathcal{E}}_i$ satisfies the following: Let $\{\xi_n\}_{n \in \mathbb{N}} \subset \text{Dom}(\mathcal{E})$ be a sequence converging a.s. to some $\xi \in L^0(\mathcal{F}_T)$. If $\inf_{n \in \mathbb{N}} \xi_n \geq c$, a.s. for some $c \in \mathbb{R}$, then $\liminf_{n \rightarrow \infty} \tilde{\mathcal{E}}_i[\xi_n] < \infty$ implies $\xi \in \text{Dom}(\mathcal{E})$. The proof of this equivalence is similar to that of Corollary 25.

(2) It is clear that (4.6) implies $\sup_{(i, \rho) \in \mathcal{I} \times \mathcal{S}_{0,T}} \mathcal{E}_i[Y_\rho^i] < \infty$.

(3) In light of (Y3) and the right-continuity of Y , it holds except on a null set N that

$$Y_t \geq C_Y, \quad \forall t \in [0, T], \quad \langle \text{hbox} | t h u s \rangle \quad Y_\nu \geq C_Y, \quad \forall \nu \in \mathcal{S}_{0,T}. \quad (4.7)$$

Then for any $i \in \mathcal{I}$, Remark 41 (1) implies that except on a null set $\tilde{N}(i)$

$$Y_\nu^i = Y_\nu + H_\nu^i \geq C \triangleq C_Y + C_H, \quad \forall \nu \in \mathcal{S}_{0,T}. \quad (4.8)$$

The following lemma states that the supremum or infimum over a stable class of \mathbf{F} -expectations can be approached by an increasing or decreasing sequence in the class.

Lemma 42. Let $\nu \in \mathcal{S}_{0,T}$ and \mathcal{U} be a non-empty subset of $\mathcal{S}_{\nu,T}$ such that

$$\rho_1 \mathbf{1}_A + \rho_2 \mathbf{1}_{A^c} \in \mathcal{U}, \quad \forall \rho_1, \rho_2 \in \mathcal{U}, \quad \forall A \in \mathcal{F}_\nu.$$

Let $\{X(\rho)\}_{\rho \in \mathcal{U}} \subset \text{Dom}(\mathcal{E})$ be a family of random variables, indexed by ρ , such that for any $\nu, \sigma \in \mathcal{U}$, $\mathbf{1}_{\{\nu=\sigma\}} X(\nu) = \mathbf{1}_{\{\nu=\sigma\}} X(\sigma)$, a.s., then for any stable subclass $\mathcal{E}' = \{\mathcal{E}_i\}_{i \in \mathcal{I}'}$ of \mathcal{E} , there exist two sequences $\{(i_n, \rho_n)\}_{n \in \mathbb{N}}$ and $\{(i'_n, \rho'_n)\}_{n \in \mathbb{N}}$ in $\mathcal{I}' \times \mathcal{U}$ such that

$$\text{esssup}_{(i, \rho) \in \mathcal{I}' \times \mathcal{U}} \tilde{\mathcal{E}}_i[X(\rho) + H_{\nu, \rho}^i | \mathcal{F}_\nu] \quad \text{---} \quad \lim_{n \rightarrow \infty} \uparrow \tilde{\mathcal{E}}_{i_n}[X(\rho_n) + H_{\nu, \rho_n}^{i_n} | \mathcal{F}_\nu], \quad \text{a. s.}, \quad (4.9)$$

$$\text{essinf}_{(i, \rho) \in \mathcal{I}' \times \mathcal{U}} \tilde{\mathcal{E}}_i[X(\rho) + H_{\nu, \rho}^i | \mathcal{F}_\nu] \quad \text{---} \quad \lim_{n \rightarrow \infty} \downarrow \tilde{\mathcal{E}}_{i'_n}[X(\rho'_n) + H_{\nu, \rho'_n}^{i'_n} | \mathcal{F}_\nu], \quad \text{a. s.} \quad (4.10)$$

For any $\nu \in \mathcal{S}_{0,T}$ and $i \in \mathcal{I}$, let us define

$$Z(\nu) \triangleq \text{esssup}_{(i, \rho) \in \mathcal{I} \times \mathcal{S}_{\nu,T}} \tilde{\mathcal{E}}_i[Y_\nu + H_{\nu, \rho}^i | \mathcal{F}_\nu] \in \mathcal{F}_\nu \quad \langle \text{hbox} | a n d \rangle \quad Z^i(\nu) \triangleq Z(\nu) + H_\nu^i.$$

Clearly, taking $\rho = \nu$ above yields that

$$Y_\nu \leq Z(\nu), \quad \text{a. s.} \quad (4.11)$$

The following two Lemmas give the bounds on $Z(\nu)$, $Z^i(\nu)$, $i \in \mathcal{I}$, and show that they all belong to $\text{Dom}(\mathcal{E})$.

Lemma 43. For any $\nu \in \mathcal{S}_{0,T}$ and $i \in \mathcal{I}$

$$Z(\nu) \geq C \quad \langle \text{hbox} | a n d \rangle \quad Z^i(\nu) \geq C_Y + 2C_H, \quad \text{a. s.} \quad (4.12)$$

Moreover, if no member of \mathcal{E} satisfies (4.3), then we further have

$$Z(\nu) \leq \zeta_Y - C_H \quad \langle \text{hbox} | a n d \rangle \quad Z^i(\nu) \leq \zeta_Y - C_H + H_\nu^i, \quad \text{a. s.}, \quad (4.13)$$

where $\zeta_Y - C_H$ and $\zeta_Y - C_H + H_\nu^i$ both belong to $\text{Dom}(\mathcal{E})$.

Lemma 44. For any $\nu \in \mathcal{S}_{0,T}$ and $i \in \mathcal{I}$, both $Z(\nu)$ and $Z^i(\nu)$ belong to $\text{Dom}(\mathcal{E})$.

In the next two propositions, we will see that the \mathbf{F} -adapted process $\{Z(t)\}_{t \in [0, T]}$ has an RCLL modification Z^0 , and that both $\{Z^i(t)\}_{t \in [0, T]}$ and $Z^{i,0} \triangleq \{Z_t^0 + H_t^i\}_{t \in [0, T]}$ are $\tilde{\mathcal{E}}_i$ -supermartingales for any $i \in \mathcal{I}$.

Proposition 45. *For any $\nu, \sigma \in \mathcal{S}_{0, T}$ and $\gamma \in \mathcal{S}_{\nu, T}$, we have*

$$Z(\nu) = Z(\sigma), \quad a.s. \text{ on } \{\nu = \sigma\}, \quad (4.14)$$

$$\operatorname{esssup}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[Z(\gamma) + H_{\nu, \gamma}^i | \mathcal{F}_\nu] = \bigwedge_{(i, \rho) \in \mathcal{I} \times \mathcal{S}_{\gamma, T}} \operatorname{esssup} \tilde{\mathcal{E}}_i[Y_\rho + H_{\nu, \rho}^i | \mathcal{F}_\nu] \leq Z(\nu), \quad a.s. \quad (4.15)$$

Proposition 46. *Given $i \in \mathcal{I}$, for any $\nu, \rho \in \mathcal{S}_{0, T}$ with $\nu \leq \rho$, a.s., we have*

$$\tilde{\mathcal{E}}_i[Z^i(\rho) | \mathcal{F}_\nu] \leq Z^i(\nu), \quad a.s. \quad (4.16)$$

In particular, $\{Z^i(t)\}_{t \in [0, T]}$ is an $\tilde{\mathcal{E}}_i$ -supermartingale. Moreover, the process $\{Z(t)\}_{t \in [0, T]}$ admits an RCLL modification Z^0 . The process $Z^{i,0} \triangleq \{Z_t^0 + H_t^i\}_{t \in [0, T]}$ is also an $\tilde{\mathcal{E}}_i$ -supermartingale.

We call Z^0 the “ \mathcal{E} -upper Snell envelope” of the reward process Y . From (4.11) and their right-continuity, we see that Z^0 dominates Y in the following sense:

Definition 47. *We say that process X “dominates” process X' if $P(X_t \geq X'_t, \forall t \in [0, T]) = 1$.*

Remark 48. (1) *If X dominates X' , then $X_\nu \geq X'_\nu$, a.s. for any $\nu \in \mathcal{S}_{0, T}$.*

(2) *Let X and X' be two right-continuous \mathbf{F} -adapted processes. If $P(X_t \geq X'_t) = 1$ holds for all t in a countable dense subset of $[0, T]$, then X dominates X' .*

The following Proposition demonstrates that Z^0 is the smallest RCLL \mathbf{F} -adapted process dominating Y such that $Z^{i,0}$ is an $\tilde{\mathcal{E}}_i$ -supermartingale for any $i \in \mathcal{I}$.

Proposition 49. *The process Z^0 dominates the process Y . Moreover, for any $\nu \in \mathcal{S}_{0, T}$ and $i \in \mathcal{I}$, we have $Z_\nu^0, Z_\nu^{i,0} \in \operatorname{Dom}(\mathcal{E})$ and*

$$Z_\nu^0 = Z(\nu), \quad Z_\nu^{i,0} = Z^i(\nu), \quad a.s. \quad (4.17)$$

Furthermore, if X is another RCLL \mathbf{F} -adapted process dominating Y such that $X^{i,0} \triangleq \{X_t + H_t^i\}_{t \in [0, T]}$ is an $\tilde{\mathcal{E}}_i$ -supermartingale for any $i \in \mathcal{I}$, then X also dominates Z^0 .

As a consequence of Proposition 49 and (?), we have for any $\nu \in \mathcal{S}_{0, T}$ and $i \in \mathcal{I}$ that

$$Z_\nu^0 \geq C, \quad Z_\nu^{i,0} \geq C_Y + 2C_H, \quad a.s. \quad (4.18)$$

Remark 50. If $\{X_t\}_{t \in [0, T]}$ is a càdlàg \mathcal{E} -supermartingale satisfying

$$P\{X_t \geq Y_t\} = 1, \quad \forall t \in [0, T],$$

then it is still true that X dominates Z^0 .

Our main concern in this section is to find an optimal stopping time for the cooperative game.

Definition 51. A stopping time $\tau^* \in \mathcal{S}_{0, T}$ is called an “ \mathcal{E} -optimal” stopping time for the cooperative game if

$$\sup_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[Y_{\tau^*}] = Z(0) = \sup_{(i, \rho) \in \mathcal{I} \times \mathcal{S}_{0, T}} \neg \neg \tilde{\mathcal{E}}_i[Y_{\rho}].$$

⟨fi⟩

In what follows, we first give *approximately* optimal stopping times. This family of stopping times will be crucial in finding an optimal stopping time for (4.1).

Definition 52. For any $\delta \in (0, 1)$ and $\nu \in \mathcal{S}_{0, T}$, we define

$$\begin{aligned} \tau_{\delta}(\nu) &\stackrel{\Delta}{=} \inf \{t \in [\nu, T]: Y_t \geq \delta Z_t^0 + (1 - \delta)(C_Y + 2C_H)\} \wedge T \in \mathcal{S}_{\nu, T} \\ \langle \text{error} | \text{compound hbox} \rangle \quad J_{\delta}(\nu) &\stackrel{\Delta}{=} \text{esssup}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[Z_{\tau_{\delta}(\nu)}^0 + H_{\nu, \tau_{\delta}(\nu)}^i | \mathcal{F}_{\nu}]. \end{aligned}$$

Remark 53. (1) For any $\delta \in (0, 1)$ and $\nu \in \mathcal{S}_{0, T}$, the right-continuity of Y and Z^0 implies that $\{\tau_{\delta}(t)\}_{t \in [0, T]}$ is also a right-continuous process. Moreover, since $Z_T^0 = Z(T) = Y_T$, a.s., we can deduce from (Y3) that $Y_T > \delta Z_T^0 + (1 - \delta)(C_Y + 2C_H)$. Then the right-continuity of processes Y and Z^0 implies that

$$Y_{\tau_{\delta}(\nu)} \geq \delta Z_{\tau_{\delta}(\nu)}^0 + (1 - \delta)(C_Y + 2C_H), \quad \text{a.s.}$$

(2) For any $\nu \in \mathcal{S}_{0, T}$, we can deduce from (4.17) and (4.15) that

$$\begin{aligned} J_{\delta}(\nu) &= \text{esssup}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[Z_{\tau_{\delta}(\nu)}^0 + H_{\nu, \tau_{\delta}(\nu)}^i | \mathcal{F}_{\nu}] = \text{esssup}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[Z(\tau_{\delta}(\nu)) + H_{\nu, \tau_{\delta}(\nu)}^i | \mathcal{F}_{\nu}] \leq \\ Z(\nu) &= Z_{\nu}^0, \quad \text{a.s.} \end{aligned} \tag{4.19}$$

The following two results show that for any $\delta \in (0, 1)$, the family $\{J_{\delta}(\nu)\}_{\nu \in \mathcal{S}_{0, T}}$ possesses similar properties to $\{Z(\nu)\}_{\nu \in \mathcal{S}_{0, T}}$.

Lemma 54. For any $\delta \in (0, 1)$ and $\nu \in \mathcal{S}_{0, T}$, we have $J_{\delta}(\nu) \in \text{Dom}(\mathcal{E})$. And for any $\sigma \in \mathcal{S}_{0, T}$, $J_{\delta}(\nu) = J_{\delta}(\sigma)$, a.s. on $\{\nu = \sigma\}$.

Proposition 55. Given $\delta \in (0, 1)$, the followings statements hold:

- (1) For any $i \in \mathcal{I}$, $\{J_\delta^i(t) \stackrel{\Delta}{=} J_\delta(t) + H_t^i\}_{t \in [0, T]}$ is an $\tilde{\mathcal{E}}_i$ -supermartingale;
- (2) $\{J_\delta(t)\}_{t \in [0, T]}$ admits an RCLL modification $J^{\delta, 0}$ such that the process $J^{\delta, 0} \stackrel{\Delta}{=} \{J_t^{\delta, 0} + H_t^i\}_{t \in [0, T]}$ is an $\tilde{\mathcal{E}}_i$ -supermartingale for each $i \in \mathcal{I}$;
- (3) For any $\nu \in \mathcal{S}_{0, T}$, $J_\nu^{\delta, 0} \in \text{Dom}(\mathcal{E})$ and $J_\nu^{\delta, 0} = J_\delta(\nu)$, a.s.

Fix $\nu \in \mathcal{S}_{0, T}$. The right continuity of Z^0 and (?) imply that the stopping times $\tau_\delta(\nu)$ are increasing in δ . Therefore, we can define the limiting stopping time

$$\bar{\tau}(\nu) \stackrel{\Delta}{=} \lim_{\delta \nearrow 1} \tau_\delta(\nu). \quad (4.20)$$

To show that $\bar{\tau}(0) \in \mathcal{S}_{0, T}$ is an optimal stopping time for (4.1), we need the family of processes $\{Y^i\}_{i \in \mathcal{I}}$ to be uniformly continuous from the left over the stable class \mathcal{E} .

Definition 56. The family $\{Y^i\}_{i \in \mathcal{I}}$ is called “ \mathcal{E} -uniformly-left-continuous” if for any $\nu, \rho \in \mathcal{S}_{0, T}$ with $\nu \leq \rho$, a.s. and for any sequence $\{\rho_n\}_{n \in \mathbb{N}} \subset \mathcal{S}_{\nu, T}$ increasing a.s. to ρ , we can find a subsequence $\{n_k\}_{k \in \mathbb{N}}$ of \mathbb{N} such that

$$\lim_{k \rightarrow \infty} \text{esssup}_{i \in \mathcal{I}} \left| \tilde{\mathcal{E}}_i \left[\langle \text{hbox} | \frac{n_k}{n_k - 1} \rangle Y_{\rho_{n_k}} + H_{\rho_{n_k}}^i \middle| \mathcal{F}_\nu \right] - \tilde{\mathcal{E}}_i[Y_\rho^i | \mathcal{F}_\nu] \right| = 0, \quad \text{a.s.} \quad (4.21)$$

The next theorem shows that $\bar{\tau}(\nu)$ is not only the first time when process Z^0 meets the process Y after ν , but it is also an optimal stopping time after ν . The assumption that the elements of the stable class \mathcal{E} are convex (see (?)) becomes crucial in the proof of this result.

Theorem 57. Assume that $\{Y^i\}_{i \in \mathcal{I}}$ is “ \mathcal{E} -uniformly-left-continuous”. Then for each $\nu \in \mathcal{S}_{0, T}$, the stopping time $\bar{\tau}(\nu)$ defined by (4.20) satisfies

$$\begin{aligned} Z(\nu) &= \text{esssup}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[\neg[Y_{\bar{\tau}(\nu)} \neg + \neg H_{\nu, \bar{\tau}(\nu)}^i \neg] | \mathcal{F}_\nu] \neg = \text{esssup}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[\neg[Z(\bar{\tau}(\nu)) \neg + \neg H_{\nu, \bar{\tau}(\nu)}^i \neg] | \mathcal{F}_\nu] \neg \\ &= \text{esssup}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[\neg[Z(\rho) \neg + \neg H_{\nu, \rho}^i \neg] | \mathcal{F}_\nu], \quad \text{a.s.} \end{aligned} \quad (4.22)$$

for any $\rho \in \mathcal{S}_{\nu, \bar{\tau}(\nu)}$ and $\bar{\tau}(\nu) = \tau_1(\nu) \stackrel{\Delta}{=} \inf \{t \in [\nu, T]: Z_t^0 = Y_t\}$, a.s.

In particular, taking $\nu = 0$ in (4.22), one can deduce from (2.8) that $\bar{\tau}(0) = \inf \{t \in [0, T]: Z_t^0 = Y_t\}$ satisfies

$$\begin{aligned} \sup_{(i, \rho) \in \mathcal{I} \times \mathcal{S}_{0, T}} \mathcal{E}_i[Y_\rho + H_\rho^i] &= \sup_{(i, \rho) \in \mathcal{I} \times \mathcal{S}_{0, T}} \tilde{\mathcal{E}}_i[Y_\rho + H_\rho^i] = Z(0) = \sup_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[\neg[Y_{\bar{\tau}(0)} + H_{\bar{\tau}(0)}^i \neg] \\ &= \sup_{i \in \mathcal{I}} \mathcal{E}_i[\neg[Y_{\bar{\tau}(0)} + H_{\bar{\tau}(0)}^i \neg]. \end{aligned}$$

Therefore, we see that $\bar{\tau}(0)$, the first time the Snell envelope Z^0 meets the process Y after time $t = 0$, is an optimal stopping time for (4.1).

5 Robust Optimal Stopping

In this section we analyze the *robust* optimal stopping problem in which the stopper aims to find an optimal stopping time τ_* that satisfies

$$\sup_{\rho \in \mathcal{S}_{0,T}} \inf_{i \in \mathcal{I}} \mathcal{E}_i[Y_\rho^i] = \inf_{i \in \mathcal{I}} \mathcal{E}_i[Y_\tau^i], \quad (5.1)$$

where $\mathcal{E} = \{\mathcal{E}_i\}_{i \in \mathcal{I}}$ is a stable class of \mathbf{F} -expectations and $Y^i = Y + H^i$, $i \in \mathcal{I}$ is the model-dependent reward process introduced in (5.1). (We will modify the assumptions on the reward processes shortly). In order to find an optimal stopping time we construct the lower and the upper values of the optimal stopping problem at any stopping time $\nu \in \mathcal{S}_{0,T}$, i.e.,

$$V(\nu) \triangleq \operatorname{esssup}_{\rho \in \mathcal{S}_{\nu,T}} \left(\operatorname{essinf}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[Y_\rho \neg + H_{\nu,\rho}^i | \mathcal{F}_\nu] \right), \quad \bar{V}(\nu) \triangleq \operatorname{essinf}_{i \in \mathcal{I}} \left(\operatorname{esssup}_{\rho \in \mathcal{S}_{\nu,T}} \tilde{\mathcal{E}}_i[Y_\rho \neg + H_{\nu,\rho}^i | \mathcal{F}_\nu] \right).$$

With the so-called “ \mathcal{E} -uniform-right-continuity” condition on $\{Y^i\}_{i \in \mathcal{I}}$, we can show that at any $\nu \in \mathcal{S}_{0,T}$, $V(\nu)$ and $\bar{V}(\nu)$ coincide with each other (see Theorem 68). We denote the common value, the *value* of the robust optimal stopping problem, as $V(\nu)$ at ν . We will show that up to a stopping time $\tau(0)$ (see Lemma 62), at which we have $V(\tau(0)) = Y_{\tau(0)}$, a.s., the stopped value process $\{V(\tau(0) \wedge t)\}_{t \in [0,T]}$ admits an RCLL modification V^0 . The main result in this section, Theorem 68, shows that the first time V^0 meets Y is an optimal stopping time for (5.1).

Standing assumptions on \mathcal{H} and Y in this section. Let us introduce

$$R^i(\nu) \triangleq \operatorname{esssup}_{\rho \in \mathcal{S}_{\nu,T}} \tilde{\mathcal{E}}_i[Y_\rho \neg + H_{\nu,\rho}^i | \mathcal{F}_\nu], \quad \langle \text{hbox} | \text{for any} \rangle i \in \mathcal{I} \langle \text{hbox} | \text{and} \rangle \nu \in \mathcal{S}_{0,T}.$$

To adapt the results we obtained for the family $\{Z(\nu)\}_{\nu \in \mathcal{S}_{0,T}}$ to each family $\{R^i(\nu)\}_{\nu \in \mathcal{S}_{0,T}}$, $i \in \mathcal{I}$, we assume that $\mathcal{H} = \{H^i\}_{i \in \mathcal{I}}$ is a family of right-continuous \mathbf{F} -adapted processes satisfying (S2), (S3) and,

- (S1') For any $i \in \mathcal{I}$, $H_0^i = 0$, a.s. and (4.2) holds. If \mathcal{E}_i does not satisfy (4.3), then we assume that $\zeta^i = \operatorname{esssup}_{s,t \in \mathcal{D}_T; s < t} H_{s,t}^i \in \operatorname{Dom}(\mathcal{E})$.

On the other hand, we assume that Y is a right-continuous \mathbf{F} -adapted process that satisfies (Y1), (Y3) and

- (Y2') For any $i \in \mathcal{I}$, $\sup_{\rho \in \mathcal{S}_{0,T}} \tilde{\mathcal{E}}_i[Y_\rho^i] < \infty$. If \mathcal{E}_i does not satisfy (4.3), then we assume that $\operatorname{esssup}_{(\rho,t) \in \mathcal{S}_{0,T} \times \mathcal{D}_T} \tilde{\mathcal{E}}_i[Y_\rho^i | \mathcal{F}_t] \in \operatorname{Dom}(\mathcal{E})$.

We also assume that for any $i \in \mathcal{I}$, Y^i is “quasi-left-continuous” under $\tilde{\mathcal{E}}_i$: for any $\nu, \rho \in \mathcal{S}_{0,T}$ with $\nu \leq \rho$, a.s. and for any sequence $\{\rho_n\}_{n \in \mathbb{N}} \subset \mathcal{S}_{\nu,T}$ increasing a.s. to ρ , we can find a subsequence $\{n_k = n_k^{(i)}\}_{k \in \mathbb{N}}$ of \mathbb{N} such that

$$\lim_{n \rightarrow \infty} \tilde{\mathcal{E}}_i \left[\left(\text{hbox} \left| \frac{n_k}{n_k - 1} \right. \right) Y_{\rho_{n_k}} + H_{\rho_{n_k}}^i \middle| \mathcal{F}_\nu \right] = \tilde{\mathcal{E}}_i[Y_\rho^i | \mathcal{F}_\nu], \quad a.s. \quad (5.2)$$

Remark 58. $(S1')$ and $(Y2')$ are analogous to $(S1)$ and $(Y2)$ respectively while the quasi-left-continuity (5.2) is the counterpart of the \mathcal{E} -uniform-left-continuity (4.21). It is obvious that $(S1')$ implies $(S1)$ and that (4.21) gives rise to (5.2). Moreover, (4.6) implies $(Y2')$: In fact, for any $i \in \mathcal{I}$, one can deduce from (?) that

$$C \leq \operatorname{esssup}_{(\rho,t) \in \mathcal{S}_{0,T} \times \mathcal{D}_T} \tilde{\mathcal{E}}_i[Y_\rho^i | \mathcal{F}_t] \leq \operatorname{esssup}_{(i,\rho,t) \in \mathcal{I} \times \mathcal{S}_{0,T} \times \mathcal{D}_T} \tilde{\mathcal{E}}_i[Y_\rho^i | \mathcal{F}_t], \quad a.s.$$

Then Lemma 54 implies that $\operatorname{esssup}_{(\rho,t) \in \mathcal{S}_{0,T} \times \mathcal{D}_T} \tilde{\mathcal{E}}_i[Y_\rho^i | \mathcal{F}_t] \in \operatorname{Dom}(\mathcal{E})$, and it follows that $\sup_{\rho \in \mathcal{S}_{0,T}} \tilde{\mathcal{E}}_i[Y_\rho^i] < \infty$. \square

Fix $i \in \mathcal{I}$. Applying Lemma 54, (?), (?), Proposition 46, Proposition 4.18 and Theorem 57 to the family $\{R^i(\nu)\}_{\nu \in \mathcal{S}_{0,T}}$, we obtain

Proposition 59. (1) For any $\nu \in \mathcal{S}_{0,T}$, $R^i(\nu)$ belongs to $\operatorname{Dom}(\mathcal{E})$ and satisfies

$$C_Y \leq Y_\nu \leq \operatorname{esssup}_{\rho \in \mathcal{S}_{\nu,T}} \tilde{\mathcal{E}}_i[Y_\rho + H_{\nu,\rho}^i | \mathcal{F}_\nu] = R^i(\nu), \quad a.s., \quad \left(\text{hbox} \left| t h u s \right. \right) C \leq Y_\nu^i, \quad a.s. \quad (5.3)$$

(2) For any $\nu, \sigma \in \mathcal{S}_{0,T}$ and $\gamma \in \mathcal{S}_{\nu,T}$, we have

$$R^i(\nu) = R^i(\sigma), \quad a.s. \left(\text{hbox} \left| o n \right. \right) \{\nu = \sigma\}, \quad (5.4)$$

$$\tilde{\mathcal{E}}_i[R^i(\gamma) + H_{\nu,\gamma}^i | \mathcal{F}_\nu] = \neg \neg \neg \operatorname{esssup}_{\rho \in \mathcal{S}_{\gamma,T}} \tilde{\mathcal{E}}_i[Y_\rho + H_{\nu,\rho}^i | \mathcal{F}_\nu] \leq R^i(\nu), \quad a.s. \quad (5.5)$$

(3) The process $\{R^i(t)\}_{t \in [0,T]}$ admits an RCLL modification $R^{i,0}$, called “ \mathcal{E}_i Snell envelope”, such that $\{R_t^{i,0} + H_t^i\}_{t \in [0,T]}$ is an $\tilde{\mathcal{E}}_i$ -supermartingale and that for any $\nu \in \mathcal{S}_{0,T}$

$$R_\nu^{i,0} = R^i(\nu), \quad a.s. \quad (5.6)$$

(4) For any $\nu \in \mathcal{S}_{0,T}$, $\tau^i(\nu) \triangleq \inf \{t \in [\nu, T] : R_t^{i,0} = Y_t\}$ is an optimal stopping time with respect to \mathcal{E}^i after time ν , i.e., for any $\gamma \in \mathcal{S}_{\nu, \tau^i(\nu)}$,

$$R^i(\nu) = \tilde{\mathcal{E}}_i[Y_{\tau^i(\nu)} + H_{\nu, \tau^i(\nu)}^i | \mathcal{F}_\nu] = \tilde{\mathcal{E}}_i[R^i(\tau^i(\nu)) + H_{\nu, \tau^i(\nu)}^i | \mathcal{F}_\nu] = \tilde{\mathcal{E}}_i[R^i(\gamma) + H_{\nu, \gamma}^i | \mathcal{F}_\nu], \quad a.s. \quad (5.7)$$

Corollary 60. For any $\nu \in \mathcal{S}_{0,T}$, both $V(\nu)$ and $\bar{V}(\nu)$ belong to $\operatorname{Dom}(\mathcal{E})$.

Proof: Fix $(l, \rho) \in \mathcal{I} \times \mathcal{S}_{\nu, T}$, for any $i \in \mathcal{I}$, (??), (??) and Proposition 26 (5) imply that

$$\tilde{\mathcal{E}}_i[Y_\rho \neg + H_{\nu, \rho}^i | \mathcal{F}_\nu] \geq \tilde{\mathcal{E}}_i[C_Y + C_H | \mathcal{F}_\nu] = C, \quad a. s.$$

Taking the essential infimum over $i \in \mathcal{I}$ on the left-hand-side yields that

$$C \leq \operatorname{essinf}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[Y_\rho \neg + H_{\nu, \rho}^i | \mathcal{F}_\nu] \leq \operatorname{esssup}_{\rho \in \mathcal{S}_{\nu, T}} \left(\operatorname{essinf}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[Y_\rho \neg + H_{\nu, \rho}^i | \mathcal{F}_\nu] \right) \neg = \bar{V}(\nu) \leq \bar{V}(\nu) = \operatorname{essinf}_{i \in \mathcal{I}} R^i(\nu) \leq R^l(\nu), \quad a. s.$$

Since $R^l(\nu) \in \operatorname{Dom}(\mathcal{E})$ by Proposition 59 (1), a simple application of Lemma 54 proves the corollary. \square

As we will see in the next lemma since the stable class \mathcal{E} is closed under pasting (see Definition 51 (2)), $\bar{V}(\nu)$ can be approximated by a decreasing sequence that belongs to the family $\{R^i(\nu)\}_{i \in \mathcal{I}}$.

Lemma 61. *For any $\nu \in \mathcal{S}_{0, T}$, there exists a sequence $\{i_n\}_{n \in \mathbb{N}} \subset \mathcal{I}$ such that*

$$\bar{V}(\nu) = \operatorname{essinf}_{i \in \mathcal{I}} R^i(\nu) = \lim_{n \rightarrow \infty} \neg \neg \downarrow R^{i_n}(\nu), \quad a. s. \quad (5.8)$$

Thanks again to the stability of \mathcal{E} under pasting, the infimum of the family $\{\tau^i(\nu)\}_{i \in \mathcal{I}}$ of optimal stopping times can be approached by a decreasing sequence in the family. As a result the infimum is also a stopping time.

Lemma 62. *For any $\nu \in \mathcal{S}_{0, T}$, there exists a sequence $\{i_n\}_{n \in \mathbb{N}} \subset \mathcal{I}$ such that*

$$\tau(\nu) \stackrel{\Delta}{=} \operatorname{essinf}_{i \in \mathcal{I}} \tau^i(\nu) = \lim_{n \rightarrow \infty} \neg \neg \downarrow \tau^{i_n}(\nu), \quad a. s., \quad \langle \text{hbox} | t h u s \rangle \tau(\nu) \in \mathcal{S}_{\nu, T}.$$

The family of stopping times $\{\tau(\nu)\}_{\nu \in \mathcal{S}_{0, T}}$ will play a critical role in this section. The next lemma basically shows that if $\tilde{\mathcal{E}}_j$ and $\tilde{\mathcal{E}}_k$ behave the same after some stopping time ν , then $R^{j, 0}$ and $R^{k, 0}$ are identical after ν :

Lemma 63. *For any $i, j \in \mathcal{I}$ and $\nu \in \mathcal{S}_{0, T}$, let $k = k(i, j, \nu) \in \mathcal{I}$ as in Definition 51. For any $\sigma \in \mathcal{S}_{\nu, T}$, we have $R_\sigma^{k, 0} = R^k(\sigma) = R^j(\sigma) = R_\sigma^{j, 0}$, a.s.*

Proof: For any $\rho \in \mathcal{S}_{\sigma, T}$, applying Proposition 26 (5) to $\tilde{\mathcal{E}}_i$, we can deduce from (4.5) and (3.3) that

$$\begin{aligned} \tilde{\mathcal{E}}_k[Y_\rho + H_{\sigma, \rho}^k | \mathcal{F}_\sigma] &= \tilde{\mathcal{E}}_k[Y_\rho + H_{\sigma, \rho}^j | \mathcal{F}_\sigma] = \mathcal{E}_{i, j}^\nu[Y_\rho + H_{\sigma, \rho}^j | \mathcal{F}_\sigma] = \tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_j[Y_\rho + H_{\sigma, \rho}^j | \mathcal{F}_{\nu \vee \sigma}] | \mathcal{F}_\sigma] \\ &= \tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_j[Y_\rho + H_{\sigma, \rho}^j | \mathcal{F}_\sigma] | \mathcal{F}_\sigma] = \tilde{\mathcal{E}}_j[Y_\rho + H_{\sigma, \rho}^j | \mathcal{F}_\sigma], \quad a. s. \end{aligned}$$

Then (5.6) implies that

$$R_\sigma^{k, 0} = R^k(\sigma) = \operatorname{esssup}_{\rho \in \mathcal{S}_{\sigma, T}} \tilde{\mathcal{E}}_k[Y_\rho + H_{\sigma, \rho}^k | \mathcal{F}_\sigma] = \operatorname{esssup}_{\rho \in \mathcal{S}_{\sigma, T}} \tilde{\mathcal{E}}_j[Y_\rho + H_{\sigma, \rho}^j | \mathcal{F}_\sigma] = R^j(\sigma) = R_\sigma^{j, 0}, \quad a. s.,$$

which proves the lemma. \square

We now introduce the notion of the uniform right continuity of the family $\{Y^i\}_{i \in \mathcal{I}}$ over the stable class \mathcal{E} . With this assumption on the reward processes, we can show that at any $\nu \in \mathcal{S}_{0,T}$, $V(\nu) = \bar{V}(\nu)$, a.s., thus the robust optimal stopping problem has a value $V(\nu)$ at ν .

Definition 64. *The family $\{Y^i\}_{i \in \mathcal{I}}$ is called “ \mathcal{E} -uniformly-right-continuous” if for any $\nu \in \mathcal{S}_{0,T}$ and for any sequence $\{\nu_n\}_{n \in \mathbb{N}} \subset \mathcal{S}_{\nu,T}$ decreasing a.s. to ν , we can find a subsequence of $\{\nu_n\}_{n \in \mathbb{N}}$ (we still denote it by $\{\nu_n\}_{n \in \mathbb{N}}$) such that $\lim_{n \rightarrow \infty} \operatorname{esssup}_{i \in \mathcal{I}} \left| \tilde{\mathcal{E}}_i[Y_{\nu_n}^i | \mathcal{F}_\nu] - Y_\nu^i \right| = 0$, a.s.*

Theorem 65. *Suppose that $\{Y^i\}_{i \in \mathcal{I}}$ is “ \mathcal{E} -uniformly-right-continuous”. Then for any $\nu \in \mathcal{S}_{0,T}$, we have*

$$\underline{V}(\nu) = \operatorname{essinf}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[Y_{\tau(\nu)} + \neg H_{\nu, \tau(\nu)}^i | \mathcal{F}_\nu] = \bar{V}(\nu) \geq Y_\nu, \quad \text{a.s.} \quad (5.9)$$

We will denote the common value by $V(\nu) (= \underline{V}(\nu) = \bar{V}(\nu))$. Observe that $\tau(0)$ is optimal for the robust optimal stopping problem in (5.1).

Standing assumption on Y for the rest of this section. We assume that the family of processes $\{Y^i\}_{i \in \mathcal{I}}$ is “ \mathcal{E} -uniformly-right-continuous”.

Proposition 66. *For any $\nu \in \mathcal{S}_{0,T}$, we have $V(\tau(\nu)) = Y_{\tau(\nu)}$, a.s.*

Note that $\tau(\nu)$ may not be the first time after ν when the value of the robust optimal stopping problem is equal to the primary reward. Actually, since the process $\{V(t)\}_{t \in [0,T]}$ is not necessarily right-continuous, $\inf\{t \in [\nu, T] \mid V(t) = Y_t\}$ may not even be a stopping time. We will address this issue in the next two results.

Proposition 67. *Given $i \in \mathcal{I}$, for any $\nu, \rho \in \mathcal{S}_{0,T}$ with $\nu \leq \rho$, a.s., we have*

$$\operatorname{essinf}_{k \in \mathcal{I}} \tilde{\mathcal{E}}_k[V^k(\rho) | \mathcal{F}_\nu] \leq V^i(\nu), \quad \text{a.s.}, \quad (5.10)$$

where $V^i(\nu) \triangleq V(\nu) + H_\nu^i \in \operatorname{Dom}(\mathcal{E})$. Moreover if $\rho \leq \tau(\nu)$, a.s., then

$$\tilde{\mathcal{E}}_i[V^i(\rho) | \mathcal{F}_\nu] \geq V^i(\nu), \quad \text{a.s.} \quad (5.11)$$

In particular, the stopped process $\{V^i(\tau(0) \wedge t)\}_{t \in [0,T]}$ is an $\tilde{\mathcal{E}}_i$ -submartingale.

Now we show that the stopped value process $\{V(\tau(0) \wedge t)\}_{t \in [0,T]}$ admits an RCLL modification V^0 . As a result, the first time when the process V^0 meets the process Y after time $t=0$ is an optimal stopping time of the robust optimal stopping problem.

Theorem 68. Assume that for some $i' \in \mathcal{I}$, $\zeta^{i'} = \operatorname{esssup}_{s, t \in \mathcal{D}_T; s < t} H_{s, t}^{i'} \in \operatorname{Dom}(\mathcal{E})$ and that there exists a concave \mathbf{F} -expectation \mathcal{E}' (not necessarily in \mathcal{E}) satisfying (H0) and (H1) such that

$$\operatorname{Dom}(\mathcal{E}') \supset \{-\xi: \xi \in \operatorname{Dom}(\mathcal{E})\} \quad \langle \text{and for every } \tilde{\mathcal{E}}_{i'}\text{-submartingale } X, \\ -X \text{ is an } \mathcal{E}'\text{-supermartingale.} \rangle \quad (5.12)$$

We also assume that for any $\rho \in \mathcal{S}_{0, T}$, there exists a $j = j(\rho) \in \mathcal{I}$ such that $\operatorname{esssup}_{t \in \mathcal{D}_T} \tilde{\mathcal{E}}_j[Y_\rho^j | \mathcal{F}_t] \in \operatorname{Dom}(\mathcal{E})$.

(1) Then the stopped value process $\{V(\tau(0) \wedge t)\}_{t \in [0, T]}$ admits an RCLL modification V^0 (called “ \mathcal{E} -lower Snell envelope” of Y) such that for any $\nu \in \mathcal{S}_{0, T}$

$$V_\nu^0 = V(\tau(0) \wedge \nu), \quad a. s. \quad (5.13)$$

(2) Consequently,

$$\tau_V \triangleq \inf\{t \in [0, T]: V_t^0 = Y_t\} \quad (5.14)$$

is a stopping time. In fact, it is an optimal stopping time of (5.1). i.e.

$$\inf_{i \in \mathcal{I}} \left(\sup_{\rho \in \mathcal{S}_{0, T}} \mathcal{E}_i[Y_\rho^i] \right) = \sup_{\rho \in \mathcal{S}_{0, T}} \left(\inf_{i \in \mathcal{I}} \mathcal{E}_i[Y_\rho^i] \right) = \inf_{i \in \mathcal{I}} \mathcal{E}_i[Y_{\tau_V}^i].$$

$\langle fi \rangle$

6 Remarks on Sections 4 & 5

Remark 1.

Let $\mathcal{E} = \{\mathcal{E}_i\}_{i \in \mathcal{I}}$ be a stable class of \mathbf{F} -expectations. For any $\xi \in \operatorname{Dom}(\mathcal{E})$ and $\nu \in \mathcal{S}_{0, T}$, we define

$$\bar{\mathcal{E}}[\xi | \mathcal{F}_\nu] \triangleq \operatorname{esssup}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[\xi | \mathcal{F}_\nu] \quad \langle \text{and} \rangle \quad \underline{\mathcal{E}}[\xi | \mathcal{F}_\nu] \triangleq \operatorname{essinf}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[\xi | \mathcal{F}_\nu]$$

as the maximal and minimal expectation of ξ over \mathcal{E} at the stopping time ν . It is worth pointing out that $\bar{\mathcal{E}}$ is not an \mathcal{F} -expectation on $\operatorname{Dom}(\mathcal{E})$ simply because $\bar{\mathcal{E}}[\xi | \mathcal{F}_t]$ may not belong to $\operatorname{Dom}(\mathcal{E})$ for some $\xi \in \operatorname{Dom}(\mathcal{E})$ and $t \in [0, T]$. On the other hand, we will see in Example 36 that neither $\bar{\mathcal{E}}$ nor $\underline{\mathcal{E}}$ satisfy strict monotonicity. Moreover, as we shall see in the same example, $\bar{\mathcal{E}}$ does not satisfy (H2) while $\underline{\mathcal{E}}$ does not satisfy (H1); thus we do not have a dominated convergence theorem for either $\bar{\mathcal{E}}$ or $\underline{\mathcal{E}}$. Note also that $\underline{\mathcal{E}}$ may not even be convex.

Our results in Sections 4 and 5 can be interpreted as a first step in extending the results for the optimal stopping problem $\sup_{\rho \in \mathcal{S}_{0,T}} \mathcal{E}_i[Y_\rho]$, in which \mathcal{E}_i ($i \in \mathcal{I}$) is an \mathbf{F} -expectation satisfying positive convexity and the assumptions (H1)-(H3), to optimal stopping problems for other non-linear expectations, such as $\bar{\mathcal{E}}$ and $\underline{\mathcal{E}}$, which may fail to satisfy these assumptions.

Example 69. Consider a probability space $([0, \infty), \mathcal{B}[0, \infty), \mathbf{F} = \{\mathcal{F}_t\}_{t \geq 0}, P)$ be a filtered probability space in which P is defined by

$$P(A) \triangleq \int_A e^{-x} dx, \quad \forall A \in \mathcal{B}[0, \infty).$$

We assume that the filtration \mathbf{F} satisfies the usual hypothesis. Let \mathcal{P} denote the set of all probability measures equivalent to P . For any $n \in \mathbb{N}$, we define a $P_n \in \mathcal{P}$ by

$$P_n(A) \triangleq n \int_A e^{-nx} dx, \quad \forall A \in \mathcal{B}[0, \infty).$$

As discussed in Example 69, $\mathcal{E} = \{E_Q\}_{Q \in \mathcal{P}}$ is a stable class. For any $h > 0$, one can deduce that

$$\begin{aligned} 1 &= \neg \sup_{Q \in \mathcal{P}} E_Q[1] \neg \geq \neg \bar{\mathcal{E}}[\mathbf{11}_{[0,h]}] \neg = \neg \sup_{Q \in \mathcal{P}} E_Q[\mathbf{11}_{[0,h]}] \neg \geq \sup_{n \in \mathbb{N}} E_{P_n}[\mathbf{11}_{[0,h]}] \neg = \\ \sup_{n \in \mathbb{N}} P_n[0, h] &= \lim_{n \in \mathbb{N}} (1 - e^{-nh}) \neg = 1, \end{aligned}$$

where we used the fact that $\tilde{E}_Q = E_Q$ for any $Q \in \mathcal{P}$ since $E_Q[\xi|\mathcal{F}]$ is an RCLL process for any $\xi \in L^1([0, \infty), \mathcal{B}[0, \infty), P)$. Hence, we have

$$\bar{\mathcal{E}}[\mathbf{11}_{[0,h]}] = 1, \quad \forall h > 0,$$

which implies that $\bar{\mathcal{E}}$ does not satisfy strict monotonicity.

Moreover, $\bar{\mathcal{E}}$ does not satisfy (H2). For $\xi = 0$, $\eta = 1$ and $A_n = [0, \frac{1}{n}]$, $n \in \mathbb{N}$, it follows that

$$\lim_{n \rightarrow \infty} \neg \downarrow \bar{\mathcal{E}}[\xi + \mathbf{11}_{A_n} \eta] = \lim_{n \rightarrow \infty} \bar{\mathcal{E}}[\mathbf{11}_{[0, \frac{1}{n}]}] = 1 \neq 0 = \sup_{Q \in \mathcal{P}} \tilde{E}_Q[0] = \bar{\mathcal{E}}[0] = \bar{\mathcal{E}}[\xi].$$

On the other hand, it is simple to see that $\underline{\mathcal{E}}[\mathbf{11}_{[h, \infty)}] = 0$ for any $h > 0$, which means that $\underline{\mathcal{E}}$ does not satisfy strict monotonicity either. Furthermore, $\underline{\mathcal{E}}$ does not satisfy (H1). For $\xi = 1$ and $A_n = [\frac{1}{n}, \infty)$, $n \in \mathbb{N}$, we have that

$$\begin{aligned} \lim_{n \rightarrow \infty} \neg \uparrow \underline{\mathcal{E}}[\mathbf{11}_{A_n} \xi] &= \lim_{n \rightarrow \infty} \underline{\mathcal{E}}[\mathbf{11}_{[\frac{1}{n}, \infty)}] = 0 \neq 1 = \\ \inf_{Q \in \mathcal{P}} \tilde{E}_Q[1] &= \underline{\mathcal{E}}[1] = \underline{\mathcal{E}}[\xi]. \end{aligned} \quad \langle \text{hbox} | \quad \square \rangle$$

Although it does not satisfy strict monotonicity, $\underline{\mathcal{E}}$ is almost an \mathbf{F} -expectations on $\text{Dom}(\underline{\mathcal{E}})$ as we will see next.

Proposition 70. For any $t \in [0, T]$, $\mathcal{E}[\cdot|\mathcal{F}_t]$ is an operator from $Dom(\mathcal{E})$ to $Dom_t(\mathcal{E}) \triangleq Dom(\mathcal{E}) \cap L^0(\mathcal{F}_t)$. Moreover, the family of operators $\{\mathcal{E}[\cdot|\mathcal{F}_t]\}_{t \in [0, T]}$ satisfies (A2)-(A4) as well as

$$\mathcal{E}[\xi|\mathcal{F}_t] \leq \mathcal{E}[\eta|\mathcal{F}_t], \quad a.s. \quad \langle \text{hbox} | \text{for any} \rangle \xi, \eta \in Dom(\mathcal{E}) \langle \text{hbox} | \text{with} \rangle \xi \leq \eta, \quad a.s. \quad (6.1)$$

Remark 2.

We have found that the first time $\bar{\tau}(0)$ when the Snell envelope Z^0 meets the process Y is an optimal stopping time for (4.1) while the first time τ_V when the process V^0 meets the process Y is an optimal stopping time for (5.1). It is natural to ask whether $\bar{\tau}(0)$ (resp. τ_V) is the minimal optimal stopping time of (4.1) (resp. (5.1)). This answer is affirmative when \mathcal{E} is a singleton. Let \mathcal{E} be a positively-convex \mathbf{F} -expectation satisfying (H1)-(H3) and let Y be a right-continuous \mathbf{F} -adapted process satisfying (Y1), (Y3) and the following

$$\sup_{\rho \in \mathcal{S}_{0,T}} \mathcal{E}[Y_\rho] < \infty; \quad \langle \text{hbox} \quad | \quad i \text{ f } \mathcal{E} \text{ does not satisfy (5.12),} \rangle$$

then $\text{esssup}_{(\rho,t) \in \mathcal{S}_{0,T} \times \mathcal{D}_T} \tilde{\mathcal{E}}[Y_\rho|\mathcal{F}_t] \in Dom^\#(\mathcal{E}).$

(Note that we have here merged the cumulative reward process H into the primary reward process Y .) If $\tau \in \mathcal{S}_{0,T}$ is an optimal stopping time for (4.1), i.e. $\sup_{\rho \in \mathcal{S}_{0,T}} \mathcal{E}[Y_\rho] = \mathcal{E}[Y_\tau]$, Proposition 46 and (4.17) imply that

$$\sup_{\rho \in \mathcal{S}_{0,T}} \mathcal{E}[Y_\rho] = \sup_{\rho \in \mathcal{S}_{0,T}} \tilde{\mathcal{E}}[Y_\rho] = Z(0) \geq \tilde{\mathcal{E}}[Z(\tau)] = \tilde{\mathcal{E}}[Z_\tau^0] = \mathcal{E}[Z_\tau^0] \geq \mathcal{E}[Y_\tau] = \sup_{\rho \in \mathcal{S}_{0,T}} \mathcal{E}[Y_\rho],$$

thus $\mathcal{E}[Z_\tau^0] = \mathcal{E}[Y_\tau]$. The second part of (A1) then implies that $Z_\tau^0 = Y_\tau$, a.s. Hence $\bar{\tau}(0) \leq \tau$, a.s., which means that $\bar{\tau}(0)$ is the minimal stopping time for (4.1).

However, this is not the case in general. Let $\mathcal{E} = \{\mathcal{E}_i\}_{i \in \mathcal{I}}$ be a stable class of \mathbf{F} -expectations and let Y be a right-continuous \mathbf{F} -adapted process satisfying (Y1)-(Y3). We take $H^i \equiv 0$ for any $i \in \mathcal{I}$. If $\tau \in \mathcal{S}_{0,T}$ is an optimal stopping time for (4.1), i.e. $\sup_{(i,\rho) \in \mathcal{I} \times \mathcal{S}_{0,T}} \mathcal{E}_i[Y_\rho] = \sup_{i \in \mathcal{I}} \mathcal{E}_i[Y_\tau]$, (?) and (4.17) then imply that

$$\begin{aligned} \sup_{(i,\rho) \in \mathcal{I} \times \mathcal{S}_{0,T}} \mathcal{E}_i[Y_\rho] &= \sup_{(i,\rho) \in \mathcal{I} \times \mathcal{S}_{0,T}} \tilde{\mathcal{E}}_i[Y_\rho] = Z(0) \geq \sup_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[Z(\tau)] = \sup_{i \in \mathcal{I}} \mathcal{E}_i[Z(\tau)] \\ &= \sup_{i \in \mathcal{I}} \mathcal{E}_i[Z_\tau^0] \geq \sup_{i \in \mathcal{I}} \mathcal{E}_i[Y_\tau] = \sup_{(i,\rho) \in \mathcal{I} \times \mathcal{S}_{0,T}} \mathcal{E}_i[Y_\rho], \end{aligned}$$

thus $\bar{\mathcal{E}}[Z_\tau^0] = \sup_{i \in \mathcal{I}} \mathcal{E}_i[Z_\tau^0] = \sup_{i \in \mathcal{I}} \mathcal{E}_i[Y_\tau] = \bar{\mathcal{E}}[Y_\tau]$. However, this may not imply that $Z_\tau^0 = Y_\tau$, a.s. since $\bar{\mathcal{E}}$ does not satisfy strict monotonicity as we have seen in Example 69.

Now we further assume that Y satisfies (Y2'), if $\tau' \in \mathcal{S}_{0,T}$ is an optimal stopping time for (5.1), i.e. $\sup_{\rho \in \mathcal{S}_{0,T}} \inf_{i \in \mathcal{I}} \mathcal{E}_i[Y_\rho] = \inf_{i \in \mathcal{I}} \mathcal{E}_i[Y_{\tau'}]$, (5.10) and Theorem 68 imply that

$$\begin{aligned} \sup_{\rho \in \mathcal{S}_{0,T}} \inf_{i \in \mathcal{I}} \mathcal{E}_i[Y_\rho] &= \sup_{\rho \in \mathcal{S}_{0,T}} \inf_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[Y_\rho] = V(0) = V(0) \geq \inf_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[V(\tau')] = \\ &\quad \inf_{i \in \mathcal{I}} \mathcal{E}_i[V(\tau')] \\ &\geq \inf_{i \in \mathcal{I}} \mathcal{E}_i[Y_{\tau'}] = \sup_{\rho \in \mathcal{S}_{0,T}} \inf_{i \in \mathcal{I}} \mathcal{E}_i[Y_\rho], \end{aligned}$$

thus $\mathcal{E}[V(\tau')] = \inf_{i \in \mathcal{I}} \mathcal{E}_i[V(\tau')] = \inf_{i \in \mathcal{I}} \mathcal{E}_i[Y_{\tau'}] = \mathcal{E}[Y_{\tau'}]$. However, this may not imply that $V(\tau') = Y_{\tau'}$, a.s. since \mathcal{E} does not satisfy strict monotonicity, which we have also seen in Example 69. (If $V(\tau')$ were a.s. equal to $Y_{\tau'}$, for any $i \in \mathcal{I}$, applying (?) to singleton $\{\mathcal{E}_i\}$, we would deduce from (5.13) and Lemma 63 that

$$\begin{aligned} V_{\tau' \wedge \tau_V}^0 &= V(\tau' \wedge \tau_V) = \bar{V}(\tau' \wedge \tau_V) = \operatorname{ess\,inf}_{i \in \mathcal{I}} R^i(\tau' \wedge \tau_V) = \\ &\quad \operatorname{ess\,inf}_{i \in \mathcal{I}} (\mathbf{1}_{\{\tau' \leq \tau_V\}} R^i(\tau') + \mathbf{1}_{\{\tau' > \tau_V\}} R^i(\tau_V)) \\ &= \mathbf{1}_{\{\tau' \leq \tau_V\}} \operatorname{ess\,inf}_{i \in \mathcal{I}} R^i(\tau') + \mathbf{1}_{\{\tau' > \tau_V\}} \operatorname{ess\,inf}_{i \in \mathcal{I}} R^i(\tau_V) = \mathbf{1}_{\{\tau' \leq \tau_V\}} \bar{V}(\tau') \\ &\quad + \mathbf{1}_{\{\tau' > \tau_V\}} \bar{V}(\tau_V) \\ &= \mathbf{1}_{\{\tau' \leq \tau_V\}} V(\tau') + \mathbf{1}_{\{\tau' > \tau_V\}} V(\tau_V) = \mathbf{1}_{\{\tau' \leq \tau_V\}} V(\tau') + \\ &\quad \mathbf{1}_{\{\tau' > \tau_V\}} V_{\tau_V}^0 \\ &= \mathbf{1}_{\{\tau' \leq \tau_V\}} Y_{\tau'} + \mathbf{1}_{\{\tau' > \tau_V\}} Y_{\tau_V} = Y_{\tau' \wedge \tau_V}, \quad a.s., \end{aligned}$$

which would further imply that $\tau_V = \tau' \wedge \tau_V$, a.s., thus $\tau_V \leq \tau'$, a.s.)

7 Applications

In this section, we take a d -dimensional Brownian motion B on the probability space (Ω, \mathcal{F}, P) and consider the Brownian filtration generated by it:

$$\mathbf{F} = \left\{ \mathcal{F}_t \triangleq \sigma(\sigma(B_s; s \in [0, t]) \cup \mathcal{N}) \right\}_{t \in [0, T]}, \quad (\text{where } \mathcal{N} \text{ collects all } P\text{-null sets in } \mathcal{F}). \quad (7.1)$$

We also let \mathcal{P} denote the predictable σ -algebra with respect to \mathbf{F} .

7.1 Lipschitz g -Expectations

Suppose that a “generator” function $g = g(t, \omega, z): [0, T] \times \Omega \times \mathbb{R}^d \mapsto \mathbb{R}$ satisfies

$$\left\{ \begin{array}{l} (i) \quad g(t, \omega, 0) = 0, \quad dt \times dP - a.s. \\ (ii) \quad g \text{ is Lipschitz in } z \text{ for some } K_g > 0: \text{ it holds } dt \times dP - a.s. \text{ that} \\ |g(t, \omega, z_1) - g(t, \omega, z_2)| \leq K_g |z_1 - z_2|, \quad \forall z_1, z_2 \in \mathbb{R}^d. \end{array} \right. \quad (7.2)$$

For any $\xi \in L^2(\mathcal{F}_T)$, it is well known from [\[PP-90\]](#) that the backward stochastic differential equation (BSDE)

$$\Gamma_t = \xi + \int_t^T g(s, \Theta_s) ds - \int_t^T \Theta_s dB_s, \quad t \in [0, T] \quad (7.3)$$

admits a unique solution $(\Gamma^{\xi, g}, \Theta^{\xi, g}) \in \mathbb{C}_F^2([0, T]) \times \mathcal{H}_F^2([0, T]; \mathbb{R}^d)$ (for convenience, we will denote (7.3) by BSDE(ξ, g) in the sequel), based on which [\[Peng-97\]](#) introduced the so-called “ g -expectation” of ξ by

$$\mathcal{E}_g[\xi | \mathcal{F}_t] \triangleq \Gamma_t^{\xi, g}, \quad t \in [0, T]. \quad (7.4)$$

To show that the g -expectation \mathcal{E}_g is an \mathbf{F} -expectation with domain $Dom(\mathcal{E}_g) = L^2(\mathcal{F}_T)$, we first note that $L^2(\mathcal{F}_T) \in \tilde{\mathcal{H}}_T$. The (strict) Comparison Theorem for BSDEs (see e.g. [\[Theorem 35.3\]](#)) then shows that (A1) holds for the family of operators $\{\mathcal{E}_g[\cdot | \mathcal{F}_t]: L^2(\mathcal{F}_T) \mapsto L^2(\mathcal{F}_t)\}_{t \in [0, T]}$, while the uniqueness of the solution $(\Gamma^{\xi, g}, \Theta^{\xi, g})$ to the BSDE(ξ, g) implies that the family $\{\mathcal{E}_g[\cdot | \mathcal{F}_t]\}_{t \in [0, T]}$ satisfies (A2)-(A4) (see e.g. [\[Lemma 36.6\]](#) and [\[Lemma 2.1\]](#)). Therefore, \mathcal{E}_g is an \mathbf{F} -expectation with domain $Dom(\mathcal{E}_g) = L^2(\mathcal{F}_T)$.

Moreover, the generator g characterizes \mathcal{E}_g in the following ways:

(1) Theorem 3.2 of [\[Jiang_2008\]](#) (see also Proposition 10 of [\[Eman\]](#)) shows that $\mathcal{E}_g[\cdot | \mathcal{F}_t]$ is a convex (resp. concave) operator on $L^2(\mathcal{F}_T)$ for any $t \in [0, T]$ if and only if the generator g is *convex* (resp. *concave*) in z , i.e., it holds $dt \times dP - a.s.$ that

$$g(t, \lambda z_1 + (1 - \lambda) z_2) \leq (\text{resp. } \geq) \lambda g(t, z_1) + (1 - \lambda) g(t, z_2), \quad \forall \lambda \in (0, 1), \quad \forall z_1, z_2 \in \mathbb{R}^d. \quad (7.5)$$

(2) Let \tilde{g} be another generator satisfying (7.2). If it holds $dt \times dP - a.s.$ that $g(t, z) \geq \tilde{g}(t, z)$, $\forall z \in \mathbb{R}^d$,

the Comparison Theorem for BSDEs (see e.g. [\[EPQ-97\]](#) or [\[Proposition 5.1\]](#)) shows that for any $\xi \in L^2(\mathcal{F}_T)$ and $t \in [0, T]$

$$\mathcal{E}_g[\xi | \mathcal{F}_t] \geq \mathcal{E}_{\tilde{g}}[\xi | \mathcal{F}_t], \quad a.s. \quad (7.6)$$

Thanks to Theorem 4.1 of [\[BCHMP\]](#), the reverse statement also holds given that almost surely, the mapping $t \mapsto g(t, z)$ is continuous for any $z \in \mathbb{R}^d$.

(3) $g^-(t, \omega, z) \triangleq -g(t, \omega, -z)$, $(t, \omega, z) \in [0, T] \times \Omega \times \mathbb{R}^d$ also satisfies (7.2). Its corresponding g -expectation \mathcal{E}_{g^-} relates to \mathcal{E}_g in that for any $\xi \in L^2(\mathcal{F}_T)$ and $t \in [0, T]$

$$\mathcal{E}_{g^-}[\xi | \mathcal{F}_t] = -\mathcal{E}_g[-\xi | \mathcal{F}_t], \quad a.s. \quad (7.7)$$

(In fact, multiplying both sides of BSDE $(-\xi, g)$ by -1 , we see that the pair $(-\Gamma^{-\xi, g}, -\Theta^{-\xi, g})$ solves the BSDE (ξ, g^-) .)

To show that the g -expectation \mathcal{E}_g satisfies (H0)-(H3), we need two basic inequalities it satisfies.

Lemma 71. *Let g be a generator satisfying (7.2).*

(1) *For any $\xi \in L^2(\mathcal{F}_T)$, we have*

$$\left| \sup_{t \in [0, T]} \mathcal{E}_g[\xi | \mathcal{F}_t] \right|_{L^2(\mathcal{F}_T)} + |\Theta^{\xi, g}|_{L^2_{\mathbb{F}}([0, T]; \mathbb{R}^d)} \leq C e^{(K_g + K_g^2)T} \|\xi\|_{L^2(\mathcal{F}_T)},$$

where C is a universal constant independent of ξ and g .

(2) *For any $\mu \geq K_g$ and $\xi, \eta \in L^2(\mathcal{F}_T)$, it holds a.s. that*

$$|\mathcal{E}_g[\xi | \mathcal{F}_t] - \mathcal{E}_g[\eta | \mathcal{F}_t]| \leq \mathcal{E}_{g_\mu}[|\xi - \eta| | \mathcal{F}_t], \quad \forall t \in [0, T],$$

where the generator g_μ is defined by $g_\mu(z) \triangleq \mu|z|$, $z \in \mathbb{R}^d$.

Proof: A simple application of (7.2, Proposition 2.2) yields (1). On the other hand, (2) is a mere generalization of (7.2, Proposition 3.7, inequality (60)) by taking into account the continuity of processes $\mathcal{E}_g[\xi | \mathcal{F}_\cdot]$ and $\mathcal{E}_{g_\mu}[\xi | \mathcal{F}_\cdot]$ for any $\xi \in L^2(\mathcal{F}_T)$. \square

Proposition 72. *Let g be a generator satisfying (7.2). Then \mathcal{E}_g satisfies (H0)-(H3).*

Remark 73. *Since $\mathcal{E}_g[\xi | \mathcal{F}_\cdot]$ is a continuous process for any $\xi \in L^2(\mathcal{F}_T)$, we see from (7.2) that $\tilde{\mathcal{E}}_g[\cdot | \mathcal{F}_\nu]$ is just a restriction of $\mathcal{E}_g[\cdot | \mathcal{F}_\nu]$ to $L^{2, \#}(\mathcal{F}_T) \triangleq \{\xi \in L^2(\mathcal{F}_T) : \xi \geq c, \text{ a.s. } \langle \text{hbox} | \text{for some} \rangle c = c(\xi) \in \mathbb{R}\}$ for any $\nu \in \mathcal{S}_{0, T}$.*

Thanks to Proposition 72, all results on \mathbf{F} -expectations \mathcal{E} and $\tilde{\mathcal{E}}$ in Section 7 are applicable to g -expectations. In the following example we deliver the promise we made in Remark 73. This example indicates that for some g -expectations, $\liminf_{n \rightarrow \infty} \mathcal{E}_g[\xi_n] < \infty$ is not a sufficient condition for $\lim_{n \rightarrow \infty} \xi_n \neg \in \neg$

$Dom^+(\mathcal{E}_g) = L^{2, +}(\mathcal{F}_T) \triangleq \neg\{\xi \in L^2(\mathcal{F}_T) : \xi \geq 0, \text{ a.s.}\}$ given that $\{\xi_n\}_{n \in \mathbb{N}}$ is an a.s. convergent sequence in $Dom^+(\mathcal{E}_g)$.

Example 74. *Consider a probability space $([0, 1], \mathcal{B}[0, 1], \lambda)$, where λ is the Lebesgue measure on $[0, 1]$. We define a generator $\tilde{g}(z) \triangleq -|z|$, $z \in \mathbb{R}^d$. For any $n \in \mathbb{N}$, it is clear that the random variable $\left\{ \xi_n(\omega) \triangleq \omega^{-\frac{1}{2} + \frac{1}{n+2}} \right\}_{\omega \in [0, 1]} \in L^{2, +}(\mathcal{F}_T) = Dom^+(\tilde{g})$. Proposition 8 (2) then implies that*

$$0 = \mathcal{E}_{\tilde{g}}[0] \leq \mathcal{E}_{\tilde{g}}[\xi_n] = \Gamma_0^{\xi_n \neg, \tilde{g}} = \xi_n - \int_0^T |\Theta_s^{\xi_n \neg, \tilde{g}}| ds - \int_0^T \Theta_s^{\xi_n \neg, \tilde{g}} dB_s \leq \xi_n - \int_0^T \Theta_s^{\xi_n \neg, \tilde{g}} dB_s.$$

Taking the expected value of the above inequality yields that

$$0 \leq \mathcal{E}_{\tilde{g}}[\xi_n] \leq E \left[\xi_n - \langle \text{hbox} \mid \int_0^T \Theta_s^{\xi_n, \tilde{g}} d B_s \rangle \right] = E[\xi_n] = \int_0^1 \omega^{-\frac{1}{2} + \frac{1}{n+2}} d\omega = \frac{1}{\frac{1}{2} + \frac{1}{n+2}} < 2. \quad (7.8)$$

Since $\{\xi_n\}_{n \in \mathbb{N}}$ is an increasing sequence, we can deduce from (A1) and (7.8) that $0 \leq \lim_{n \rightarrow \infty} \neg \uparrow \mathcal{E}_{\tilde{g}}[\xi_n] \leq 2$. However, $\lim_{n \rightarrow \infty} \neg \uparrow \xi_n = \{\omega^{-\frac{1}{2}}\}_{\omega \in [0,1]}$ does not belong to $L^{2,+}(\mathcal{F}_T) = \text{Dom}^+(\tilde{g})$. \square

Similar to Proposition 3.3, pasting two g -expectations at any stopping time generates another g -expectation.

Proposition 75. Let g_1, g_2 be two generators satisfying (7.2) with Lipschitz coefficients K_1 and K_2 respectively. For any $\nu \in \mathcal{S}_{0,T}$, we define the pasting of $\mathcal{E}_{g_1}, \mathcal{E}_{g_2}$ at ν to be the following continuous \mathbf{F} -adapted process

$$\mathcal{E}_{g_1, g_2}^\nu[\xi | \mathcal{F}_t] \triangleq \mathbf{1}_{\{\nu \leq t\}} \mathcal{E}_{g_2}[\xi | \mathcal{F}_t] + \mathbf{1}_{\{\nu > t\}} \mathcal{E}_{g_1}[\mathcal{E}_{g_2}[\xi | \mathcal{F}_\nu] | \mathcal{F}_t], \quad \forall t \in [0, T] \quad (7.9)$$

for any $\xi \in L^2(\mathcal{F}_T)$. Then $\mathcal{E}_{g_1, g_2}^\nu$ is exactly the g -expectation \mathcal{E}_{g^ν} with

$$g^\nu(t, \omega, z) \triangleq \mathbf{1}_{\{\nu(\omega) \leq t\}} g_2(t, \omega, z) + \mathbf{1}_{\{\nu(\omega) > t\}} g_1(t, \omega, z), \quad (t, \omega, z) \in [0, T] \times \Omega \times \mathbb{R}^d, \quad (7.10)$$

which is a generator satisfying (7.2) with the Lipschitz coefficient $K_1 \vee K_2$.

Fix $M > 0$, we denote by \mathcal{G}_M the collection of all convex generators g satisfying (7.2) with Lipschitz coefficient $K_g \leq M$. Proposition 75 shows that the family of convex g -expectations $\mathcal{E}_M \triangleq \{\mathcal{E}_g\}_{g \in \mathcal{G}_M}$ is closed under the pasting (?). To wit, \mathcal{E}_M is a stable class of g -expectations in the sense of Definition 7.10. In what follows we let \mathcal{G}' be a non-empty subset of \mathcal{G}_M such that $\mathcal{E}' \triangleq \{\mathcal{E}_g\}_{g \in \mathcal{G}'}$ is closed under pasting. Correspondingly, $\{\tilde{\mathcal{E}}_g\}_{g \in \mathcal{G}'}$ is closed under the general pasting (?), thus we see from Proposition 72 that \mathcal{E}' is a stable class in the sense of Definition 7.10. In light of Remark 73, when applying the optimal stopping theory of \mathbf{F} -expectations (as presented in Section 4 and 5) to the stable class \mathcal{E}' , we can replace $\tilde{\mathcal{E}}_g$ by \mathcal{E}_g for each $g \in \mathcal{G}'$. However, such extension does not mean that we can take away (S2) and (Y3), the lower-bound requirements on reward processes Y and \mathcal{H} respectively. In fact, as we see in the proof of Proposition 46, the deduction of the RCLL modification Z^0 of the process $\{Z(t)\}_{t \in [0, T]}$ totally relies on applying Corollary 25 to certain process $\{Z^j(t)\}_{t \in [0, T]}$. This requires that the process $\{Z^j(t)\}_{t \in [0, T]}$ is bounded from below by some $c \in \mathbb{R}$, which is in turn assured by (S2) and (Y3) (see (?)). **(compare Remark); (H3) becomes futile, which is used to assume that $\tilde{\mathcal{E}}[\xi | \mathcal{F}_\nu] \in \text{Dom}^\#(\mathcal{E})$ for any $\xi \in \text{Dom}^\#(\mathcal{E})$ and $\nu \in \mathcal{S}_{0,T}$. (fi)** Now we make the following assumptions on the reward processes:

Standing assumptions on the reward processes in this subsection.

Let Y be a continuous \mathbf{F} -adapted process with

$$\zeta'_Y \triangleq \left(\operatorname{esssup}_{t \in \mathcal{D}_T} Y_t \right)^+ \in L^2(\mathcal{F}_T) \quad (7.11)$$

and satisfying (Y3). Moreover, for any $g \in \mathcal{G}'$, we suppose that the model-dependent cumulative reward process is in the form of

$$H_t^g \triangleq \int_0^t h_s^g ds, \quad \forall t \in [0, T],$$

where $\{h_t^g, t \in [0, T]\}_{g \in \mathcal{G}'}$ is a family of predictable processes satisfying the following assumptions:

($\tilde{h}1$) There exists a $c' < 0$ such that for any $g \in \mathcal{G}'$, $h_t^g \geq c'$, $dt \times dP - a.s.$

($\tilde{h}2$) The random variable $\omega \mapsto \int_0^T h'(t, \omega) dt$ belongs to $L^2(\mathcal{F}_T)$ with $h'(t, \omega) \triangleq \left(\operatorname{esssup}_{g \in \mathcal{G}'} h_t^g(\omega) \right)^+$ (the essential supremum is taken with respect to the

product measure space $([0, T] \times \Omega, \mathcal{P}, \lambda \times P)$, where λ denotes the Lebesgue measure on $[0, T]$).

($\tilde{h}3$) For any $\nu \in \mathcal{S}_{0,T}$ and $g_1, g_2 \in \mathcal{G}'$, it holds for any $0 \leq s < t \leq T$ that

$$h_t^{g^\nu} = \mathbf{1}_{\{\nu \leq t\}} h_t^{g_2} + \mathbf{1}_{\{\nu > t\}} h_t^{g_1}, \quad dt \times dP - a.s.,$$

where g^ν is defined in (?).

Then the triple $(\mathcal{E}', \mathcal{H}' \triangleq \{H^g\}_{g \in \mathcal{G}'}, Y)$ satisfies all the conditions stated in Section 4 and 5. Thus we can carry out the optimal stopping theory developed for \mathbf{F} -expectations to $(\mathcal{E}', \mathcal{H}', Y)$ as we will see next.

Theorem 76. *The stable class \mathcal{E}' satisfies (?), the family of processes \mathcal{H}' satisfies (S1') (thus (S1) see Remark 73), (S2) and (S3), while the process Y satisfies (Y1), (4.6) (thus (Y2'), again by Remark 73) and (Y3). Moreover, the family of processes $\{Y_t^g \triangleq Y_t + H_t^g, t \in [0, T]\}_{g \in \mathcal{G}'}$ is both " \mathcal{E}' -uniformly-left-continuous" (thus satisfies (5.2), see also Remark 73) and " \mathcal{E}' -uniformly-right-continuous".*

7.2 Existence of an Optimal Prior in (4.1) for g -Expectations

For certain collections of g -expectations, we can even determine an optimal generator g^* in the following sense:

$$\mathcal{E}_g[Y_{\bar{\tau}(0)}^g] = \sup_{g \in \mathcal{G}} \mathcal{E}_g[Y_{\bar{\tau}(0)}^g] = \sup_{(g, \rho) \in \mathcal{G} \times \mathcal{S}_{0,T}} \mathcal{E}_g[Y_\rho^g],$$

where the optimal stopping time $\bar{\tau}(0)$ is defined as in Theorem 57.

Let S be a separable metric space with metric $|\cdot|_S$ such that S is a countable union of non-empty compact subsets. We denote by \mathfrak{S} the Borel σ -algebra of S and take $\mathcal{H}_{\mathbf{F}}^0([0, T]; S)$ as the space of admissible control strategies. For any $U \in \mathcal{H}_{\mathbf{F}}^0([0, T]; S)$, we define the generator

$$g_U(t, \omega, z) \triangleq g^o(t, \omega, z, U_t(\omega)), \quad (7.12)$$

where the function $g^o(t, \omega, z, u): [0, T] \times \Omega \times \mathbb{R}^d \times S \mapsto \mathbb{R}$ satisfies:

($g^o 1$) g^o is $\mathcal{P} \otimes \mathcal{B}(\mathbb{R}^d) \otimes \mathfrak{S} / \mathcal{B}(\mathbb{R})$ -measurable.

($g^o 2$) It holds $dt \times dP - a.s.$ that $g^o(t, \omega, 0, u) = 0$ for any $u \in S$.

($g^o 3$) g^o is Lipschitz in z : For some $K_o > 0$, it holds $dt \times dP - a.s.$ that

$$|g^o(t, \omega, z_1, u) - g^o(t, \omega, z_2, u)| \leq K_o |z_1 - z_2|, \quad \forall z_1, z_2 \in \mathbb{R}^d, \quad \forall u \in S.$$

($g^o 4$) g^o is convex in z : It holds $dt \times dP - a.s.$ that

$$g^o(t, \omega, \lambda z_1 \neg + \neg(1 \neg \neg \neg \lambda) z_2, u) \leq \lambda g^o(t, \omega, z_1, u) \neg + \neg(1 \neg \neg \neg \lambda) g^o(t, \omega, z_2, u), \\ \forall \lambda \in (0, 1), \quad \forall z_1, z_2 \in \mathbb{R}^d, \quad \forall u \in S.$$

Now fix a non-empty subset \mathfrak{U} of $\mathcal{H}_{\mathbf{F}}^0([0, T]; S)$ that preserves "pasting", i.e., for any $\nu \in \mathcal{S}_{0, T}$ and $U^1, U^2 \in \mathfrak{U}$,

$$U_t^\nu(\omega) \triangleq \mathbf{1}_{\{\nu(\omega) \leq t\}} U_t^2(\omega) + \mathbf{1}_{\{\nu(\omega) > t\}} U_t^1(\omega), \quad (t, \omega) \in [0, T] \times \Omega, \quad (7.13)$$

also belongs to \mathfrak{U} . Then it is easy to check that $\{\mathcal{E}_{g_U}\}_{U \in \mathfrak{U}} \subset \mathcal{E}_{K_o}$ forms a stable class of g -expectations.

Let Y still be a continuous \mathbf{F} -adapted process satisfying (?) and (Y3). For any $U \in \mathfrak{U}$, assume that the model dependent reward process has a density which is given by

$$h_t^U(\omega) \triangleq h(t, \omega, U_t(\omega)), \quad (t, \omega) \in [0, T] \times \Omega,$$

where $h(t, \omega, u): [0, T] \times \Omega \times S \mapsto \mathbb{R}$ is a $\mathcal{P} \otimes \mathfrak{S} / \mathcal{B}(\mathbb{R})$ -measurable function satisfying the following assumptions:

($\hat{h}1$) For some $c < 0$, it holds $dt \times dP - a.s.$ that $h(t, \omega, u) \geq c$ for any $u \in S$.

($\hat{h}2$) The random variable $\omega \mapsto \int_0^T \hat{h}(t, \omega) dt$ belongs to $L^2(\mathcal{F}_T)$ with $\hat{h}(t, \omega) \triangleq \left(\text{esssup}_{U \in \mathfrak{U}} h_t^U(\omega) \right)^+$ (the essential supremum is taken with respect to the product measure space $([0, T] \times \Omega, \mathcal{P}, \lambda \times P)$, where λ denotes the Lebesgue measure on $[0, T]$).

It is easy to see that $\{h_t^U, t \in [0, T]\}_{U \in \mathfrak{U}}$ is a family of predictable processes satisfying ($\hat{h}1$)-($\hat{h}3$). Hence, we can apply the optimal stopping theory developed for \mathbf{F} -expectations to the triple $(\{\mathcal{E}_{g_U}\}_{U \in \mathfrak{U}}, \{h^U\}_{U \in \mathfrak{U}}, Y)$ thanks to Theorem 76. Now let us construct a so-called *Hamiltonian* function

$$H(t, \omega, z, u) \triangleq g^o(t, \omega, z, u) + h(t, \omega, u), \quad (t, \omega, z, u) \in [0, T] \times \Omega \times \mathbb{R}^d \times S.$$

We assume that for any $(t, \omega, z) \in [0, T] \times \Omega \times \mathbb{R}^d$, there exists a $u = u^*(t, \omega, z) \in S$ such that

$$\sup_{u \in S} H(t, \omega, z, u) = H(t, \omega, z, u(t, \omega, z)). \quad (7.14)$$

(This is valid, for example, when the metric space S is compact and the mapping $u \mapsto H(t, \omega, z, u)$ is continuous.) Then it can be shown (see (, ?, Lemma 1) or (, ?, Lemma 16.34)) that the mapping $u^*: [0, T] \times \Omega \times \mathbb{R}^d \mapsto S$ can be selected to be $\mathcal{P} \otimes \mathcal{B}(\mathbb{R}^d)/\mathfrak{S}$ -measurable.

The following theorem is the main result of this subsection.

Theorem 77. *There exists a $U^* \in \mathfrak{U}$ such that $\sup_{(U, \rho) \in \mathfrak{U} \times \mathcal{S}_{0, T}} \mathcal{E}_{g_U}[Y_\rho^U] = \mathcal{E}_{g_{U^*}}[Y_{\bar{\tau}(0)}^{U^*}]$, where the stopping time $\bar{\tau}(0)$ is as in Theorem 57.*

7.3 The Cooperative Game of Karatzas and Zamfirescu [2006] Revisited

In this subsection, we apply the results of the last subsection to extend the results of [\[Kara_Zam_2006\]](#). Let us first recall their setting:

- Consider the canonical space $(\Omega, \mathcal{F}) = (C([0, T]; \mathbb{R}^d), \mathcal{B}(C([0, T]; \mathbb{R}^d)))$ endowed with Wiener measure P , under which the coordinate mapping process $B(t, \omega) = \omega(t)$, $t \in [0, T]$ becomes a standard d -dimensional Brownian motion. We still take the filtration \mathbf{F} generated by the Brownian motion B (see (??)) and let \mathcal{P} denote the predictable σ -algebra with respect to \mathbf{F} .
- It is well-known (see e.g. (, ?, Theorem 14.6)) that given a $x \in \mathbb{R}^d$, there exists a pathwise unique, strong solution $X(\cdot)$ of the stochastic equation

$$X(t) = x + \int_0^t \sigma(s, X) dB_s, \quad t \in [0, T],$$

where the diffusion term $\sigma(t, \omega)$ is a $\mathbb{R}^{d \times d}$ -valued predictable process satisfying:

($\sigma 1$) $\int_0^T |\sigma(t, \vec{0})|^2 dt < \infty$ and $\sigma(t, \omega)$ is nonsingular for any $(t, \omega) \in [0, T] \times \Omega$.

($\sigma 2$) There exists a $K > 0$ such that for any $\omega, \tilde{\tilde{A}}_\cdot \in \Omega$ and $t \in [0, T]$

$$\|\sigma^{-1}(t, \omega)\| \leq K \quad \text{and} \quad |\sigma_{ij}(t, \omega) - \sigma_{ij}(t, \tilde{\tilde{A}}_\cdot)| \leq K \|\omega - \tilde{\tilde{A}}_\cdot\|_t, \quad \forall 1 \leq i, j \leq n \quad (7.15)$$

with $\|\omega\|_t^* \triangleq \sup_{s \in [0, t]} |\omega(s)|$.

• Let $f(t, \omega, u): [0, T] \times \Omega \times S \mapsto \mathbb{R}^d$ is a $\mathcal{P} \otimes \mathfrak{G} / \mathcal{B}(\mathbb{R}^d)$ -measurable function such that

(i) For any $u \in S$, the mapping $(t, \omega) \mapsto f(t, \omega, u)$ is predictable (i.e. \mathcal{P} -measurable);

(ii) With the same K as in (7.15),

$$|f(t, \omega, u)| \leq K(1 + \|\omega\|_t), \quad \forall (t, \omega, u) \in [0, T] \times \Omega \times S. \quad (7.16)$$

For any $U \in \tilde{\mathfrak{U}} \stackrel{\Delta}{=} \mathcal{H}_F^0([0, T]; S)$, (, ?, page 166) shows that

$$\frac{d P_U \triangle}{d P} \stackrel{\Delta}{=} \exp \left\{ \int_0^T \langle \sigma^{-1}(t, X) f(t, X, U_t), d B_t \rangle - \frac{1}{2} \int_0^T \left| \sigma^{-1}(t, X) f(t, X, U_t) \right|^2 dt \right\}$$

defines a probability measure P_U .

The objective of [\[Kara_Zam_2006\]](#) is to find an optimal stopping time $\tau^* \in \mathcal{S}_{0,T}$ and an optimal control strategy $U^* \in \tilde{\mathfrak{U}}$ that maximizes the expected reward

$$E_U \left[\varphi(X(\rho)) + \int_0^\rho h(s, X, U_s) ds \right]$$

over $(\rho, U) \in \mathcal{S}_{0,T} \times \tilde{\mathfrak{U}}$. Here $\varphi: \mathbb{R}^d \mapsto \mathbb{R}$ is a bounded continuous function, and $h(t, \omega, u): [0, T] \times \Omega \times S \mapsto \mathbb{R}$ is a $\mathcal{P} \otimes \mathfrak{G} / \mathcal{B}(\mathbb{R})$ -measurable function such that $|h(t, \omega, u)| \leq K$ for any $(t, \omega, u) \in [0, T] \times \Omega \times S$ (with the same K that appears in (7.15)).

Corollary 8 of [\[Kara_Zam_2006\]](#) shows that under (7.16), the process

$$\tilde{Z}(t) \stackrel{\Delta}{=} \operatorname{esssup}_{(U, \rho) \in \tilde{\mathfrak{U}} \times \mathcal{S}_{t,T}} E_U \left[\varphi(X(\rho)) + \int_t^\rho h(s, X, U_s) ds \middle| \mathcal{F}_t \right], \quad t \in [0, T]$$

admits an RCLL modification \tilde{Z}^0 , and that the first time processes \tilde{Z}^0 and $\{\varphi(X(t))\}_{t \in [0, T]}$ meet with each other, i.e. $\bar{\tau}(0) \stackrel{\Delta}{=} \inf \{t \in [0, T] \mid \tilde{Z}_t^0 = \varphi(X(t))\}$, is an optimal stopping time. That is,

$$\sup_{(U, \rho) \in \tilde{\mathfrak{U}} \times \mathcal{S}_{0,T}} E_U \left[\varphi(X(\rho)) + \int_0^\rho h(s, X, U_s) ds \right] = \sup_{U \in \tilde{\mathfrak{U}}} E_U \left[\varphi(X(\bar{\tau}(0))) + \int_0^{\bar{\tau}(0)} h(s, X, U_s) ds \right]. \quad (7.17)$$

Moreover, if for any $(t, \omega, z) \in [0, T] \times \Omega \times \mathbb{R}^d$, there is a $u^*(t, \omega, z) \in S$ which is $\mathcal{P} \otimes \mathcal{B}(\mathbb{R}^d) / \mathfrak{G}$ -measurable such that

$$\sup_{u \in S} \tilde{H}(t, \omega, z, u) = \tilde{H}(t, \omega, z, u(t, \omega, z)) \quad (7.18)$$

with $\tilde{H}(t, \omega, z, u) \stackrel{\Delta}{=} \sigma^{-1}(t, \omega) f(t, \omega, u), z + h(t, \omega, u), (t, \omega, z, u) \in [0, T] \times \Omega \times \mathbb{R}^d \times S$, then there further exists an optimal control strategy $U^* \in \tilde{\mathfrak{U}}$ (see (7.19), Section 8)) such that

$$\sup_{(U, \rho) \in \tilde{\mathfrak{U}} \times \mathcal{S}_{0,T}} E_U \left[\varphi(X(\rho)) + \int_0^\rho h(s, X, U_s) ds \right] = E_U \left[\varphi(X(\bar{\tau}(0))) + \int_0^{\bar{\tau}(0)} h(s, X, U_s) ds \right]. \quad (7.19)$$

In the main result of this subsection, we will show that the assumption of [\[Kara_Zam_2006\]](#) that φ and h are bounded from above by constants can be relaxed and replaced by linear-growth conditions. This comes, however, at the cost of strengthening the assumption stated in (7.16).

Proposition 78. *With the same K as in (7.15), we assume that*

$$-K \leq \varphi(x) \leq K|x|, \quad \forall x \in \mathbb{R}^d \quad (7.20)$$

and that for a.e. $t \in [0, T]$

$$|f(t, \omega, u)| \leq K \quad \langle \text{hbox} | \text{a n d} \rangle \quad -K \leq h(t, \omega, u) \leq K\|\omega\|_T, \quad \forall (\omega, u) \in \Omega \times S. \quad (7.21)$$

Then the process $\{\tilde{Z}(t)\}_{t \in [0, T]}$ has a RCLL modification \tilde{Z}^0 , and the first time $\bar{\tau}(0)$ when the process \tilde{Z}^0 meets the process $\{\varphi(X(t))\}_{t \in [0, T]}$ is an optimal stopping time; i.e., it satisfies (7.17). Moreover, if there exists a measurable mapping $u^: [0, T] \times \Omega \times \mathbb{R}^d \mapsto S$ satisfying (7.18), then there exists an optimal control strategy $U^* \in \tilde{\mathfrak{U}}$ such that (7.19) holds.*

7.4 Quadratic g -Expectations

Now we consider a *quadratic* generator $\hat{g} = \hat{g}(t, \omega, z): [0, T] \times \Omega \times \mathbb{R}^d \mapsto \mathbb{R}$ that satisfies

$$\left\{ \begin{array}{l} (i) \quad \hat{g}(t, \omega, 0) = 0, \quad dt \times dP - a.s. \\ (ii) \quad \langle \text{hbox} | \text{F o r s o m e } \kappa > 0, \text{ it holds } dt \times dP - a.s. \text{ that} \rangle \\ \quad \left| \frac{\partial \hat{g}}{\partial z}(t, \omega, z) \right| \leq \kappa(1 + |z|), \quad \forall z \in \mathbb{R}^d. \\ (iii) \quad \hat{g} \langle \text{hbox} | \text{i s c o n v e x i n } z \text{ in the sense of (7.5).} \rangle \end{array} \right. \quad (7.22)$$

Note that under (ii), (i) is equivalent to the following statement: It holds $dt \times dP - a.s.$ that

$$|\hat{g}(t, \omega, z)| \leq \kappa \left(|z| + \langle \text{hbox} | \frac{1}{2} \rangle |z|^2 \right), \quad \forall z \in \mathbb{R}^d. \quad (7.23)$$

In fact, it is clear that (7.23) implies (i). Conversely, for $d t \times d P - a.s.$ $(t, \omega) \in [0, T] \times \Omega$, one can deduce that for any $z \in \mathbb{R}^d$, $|\hat{g}(t, \omega, z)| = \left| \hat{g}(t, \omega, z) - \hat{g}(t, \omega, 0) \right| = \left| \int_0^1 \frac{\partial \hat{g}}{\partial z}(t, \lambda z) z d\lambda \right| \leq \kappa \int_0^1 (1 + \lambda |z|) |z| d\lambda = \kappa(|z| + \frac{1}{2}|z|^2)$.

For any $\xi \in L^{-e}(\mathcal{F}_T)$, (, ?, Corollary 6) (where we take $f = g$, thus $\alpha(t) \equiv \frac{\kappa}{2}$ and $(\beta, \gamma) = (0, 2\kappa)$) shows that the quadratic BSDE (ξ, \hat{g}) admits a unique solution $(\Gamma^{\xi, \hat{g}}, \Theta^{\xi, \hat{g}}) \in \mathbb{C}_{\mathbf{F}}^e([0, T]) \times M_{\mathbf{F}}([0, T]; \mathbb{R}^d)$. Hence we can correspondingly define the “quadratic” g -expectation of ξ by

$$\mathcal{E}_{\hat{g}}[\xi | \mathcal{F}_t] \triangleq \Gamma_t^{\xi, \hat{g}}, \quad t \in [0, T].$$

To show that the quadratic g -expectation $\mathcal{E}_{\hat{g}}$ is an \mathbf{F} -expectation with domain $Dom(\mathcal{E}_{\hat{g}}) = L^{-e}(\mathcal{F}_T)$, we first note that $L^{-e}(\mathcal{F}_T) \in \tilde{\mathcal{H}}_T$ (Clearly, $L^{-e}(\mathcal{F}_T)$ satisfies (D1) and (D3) and $\mathbb{R} \subset L^{-e}(\mathcal{F}_T)$). For any $\xi, \eta \in L^{-e}(\mathcal{F}_T)$, $A \in \mathcal{F}_T$ and $\lambda > 0$, we have $E[e^{\lambda |1_A \xi|}] \leq E[e^{\lambda |\xi|}] < \infty$ and $E[e^{\lambda |\xi + \eta|}] \leq E[e^{\lambda |\xi|} e^{\lambda |\eta|}] \leq \frac{1}{2} E[e^{2\lambda |\xi|}] + \frac{1}{2} E[e^{2\lambda |\eta|}] < \infty$, thus (D2) also holds for $L^{-e}(\mathcal{F}_T)$. Similar to the Lipschitz g -expectation case, the uniqueness of the solution $(\Gamma^{\xi, \hat{g}}, \Theta^{\xi, \hat{g}})$ to the quadratic BSDE (ξ, \hat{g}) implies that the family of operators $\{\mathcal{E}_{\hat{g}}[\cdot | \mathcal{F}_t]: L^{-e}(\mathcal{F}_T) \mapsto L^{-e}(\mathcal{F}_t)\}_{t \in [0, T]}$ satisfies (A2)-(A4) (cf. (, ?, Lemma 36.6) and (, ?, Lemma 2.1)), while a comparison theorem for quadratic BSDEs (see e.g. (, ?, Theorem 5)) and the following proposition show that (A1) also holds for the family $\{\mathcal{E}_{\hat{g}}[\cdot | \mathcal{F}_t]\}_{t \in [0, T]}$.

Proposition 79. *Let \hat{g} be a quadratic generator satisfying (7.22). For any $\xi^1, \xi^2 \in L^{-e}(\mathcal{F}_T)$, if $\xi^1 \geq \xi^2$, a.s., then it holds a.s. that*

$$\Gamma_t^{\xi^1, \hat{g}} \geq \Gamma_t^{\xi^2, \hat{g}}, \quad \forall t \in [0, T]. \quad (7.24)$$

Moreover, if $\Gamma_{\nu}^{\xi^1, \hat{g}} = \Gamma_{\nu}^{\xi^2, \hat{g}}$, a.s. for some $\nu \in \mathcal{S}_{0, T}$, then

$$\xi^1 = \xi^2, \quad a.s. \quad (7.25)$$

Therefore, the quadratic g -expectation $\mathcal{E}_{\hat{g}}$ is an \mathbf{F} -expectation with domain $Dom(\mathcal{E}_{\hat{g}}) = L^{-e}(\mathcal{F}_T)$. Similar to the Lipschitz g -expectation case, the convexity (7.22)(iii) of the quadratic generator \hat{g} as well as Theorem 5 of [\[BH-07\]](#) determine that $\mathcal{E}_{\hat{g}}[\cdot | \mathcal{F}_t]$ is a convex operator on $L^{-e}(\mathcal{F}_T)$ for any $t \in [0, T]$. Hence, $\mathcal{E}_{\hat{g}}$ satisfies (H0) thanks to Lemma 63. To see $\mathcal{E}_{\hat{g}}$ also satisfying (H1)-(H3), we need the following stability result.

Lemma 80. *If $\xi_n \rightarrow \xi$, a.s. and $E[e^{\lambda |\xi|}] + \sup_{n \in \mathbb{N}} E[e^{\lambda |\xi_n|}] < \infty$ for any $\lambda > 0$, then*

$$\lim_{n \rightarrow \infty} E \left[\sup_{t \in [0, T]} \left| \mathcal{E}_{\hat{g}}[\xi_n | \mathcal{F}_t] - \mathcal{E}_{\hat{g}}[\xi | \mathcal{F}_t] \right| \right] = 0. \quad (7.26)$$

Proof: Taking $f_n \equiv g$ and $f = g$ in Proposition 7 of [\[BH-07\]](#) yields that

$$\lim_{n \rightarrow \infty} E \left[\exp \left\{ p \sup_{t \in [0, T]} \left| \mathcal{E}_{\hat{g}}[\xi_n | \mathcal{F}_t] - \mathcal{E}_{\hat{g}}[\xi | \mathcal{F}_t] \right| \right\} \right] = 0, \quad \forall p \geq 1.$$

Then the strict monotonicity of (a1) implies that $\xi = \eta$, a.s., proving (A1).

2) Let $0 \leq s \leq t \leq T$, for any $A \in \mathcal{F}_s \subset \mathcal{F}_t$ and $\gamma \in \Lambda_s \subset \Lambda_t$, one can deduce that

$$\mathcal{E}^o[\mathbf{1}_A \mathcal{E}^o[\mathcal{E}^o[\xi|\mathcal{F}_t]|\mathcal{F}_s] + \gamma] = \mathcal{E}^o[\mathbf{1}_A \mathcal{E}^o[\xi|\mathcal{F}_t] + \gamma] = \mathcal{E}^o[\mathbf{1}_A \xi + \gamma].$$

Since $\mathcal{E}^o[\mathcal{E}^o[\xi|\mathcal{F}_t]|\mathcal{F}_s] \in \mathcal{F}_s$, (a2) implies that $\mathcal{E}^o[\xi|\mathcal{F}_s] = \xi_s = \mathcal{E}^o[\mathcal{E}^o[\xi|\mathcal{F}_t]|\mathcal{F}_s]$, proving (A2).

3) Fix $A \in \mathcal{F}_t$, for any $\tilde{A} \in \mathcal{F}_t$ and $\gamma \in \Lambda_t$, we have

$$\mathcal{E}^o[\mathbf{1}_{\tilde{A}}(\mathbf{1}_A \mathcal{E}^o[\xi|\mathcal{F}_t]) + \gamma] = \mathcal{E}^o[\mathbf{1}_{\tilde{A} \cap A} \mathcal{E}^o[\xi|\mathcal{F}_t] + \gamma] = \mathcal{E}^o[\mathbf{1}_{\tilde{A} \cap A} \xi + \gamma] = \mathcal{E}^o[\mathbf{1}_{\tilde{A}}(\mathbf{1}_A \xi) + \gamma].$$

Since $\mathbf{1}_A \mathcal{E}^o[\xi|\mathcal{F}_t] \in \mathcal{F}_t$, (a2) implies that $\mathcal{E}^o[\mathbf{1}_A \xi|\mathcal{F}_t] = \mathbf{1}_A \mathcal{E}^o[\xi|\mathcal{F}_t]$, proving (A3).

4) For any $A \in \mathcal{F}_t$ and $\eta, \gamma \in \Lambda_t$, (D2) implies that $\mathbf{1}_A \eta + \gamma \in \Lambda_t$, thus we have

$$\mathcal{E}^o[\mathbf{1}_A(\mathcal{E}^o[\xi|\mathcal{F}_t] + \eta) + \gamma] = \mathcal{E}^o[\mathbf{1}_A \mathcal{E}^o[\xi|\mathcal{F}_t] + (\mathbf{1}_A \eta + \gamma)] = \mathcal{E}^o[\mathbf{1}_A \xi + (\mathbf{1}_A \eta + \gamma)] = \mathcal{E}^o[\mathbf{1}_A(\xi + \eta) + \gamma].$$

Then it follows from (a2) that $\mathcal{E}^o[\xi + \eta|\mathcal{F}_t] = \mathcal{E}^o[\xi|\mathcal{F}_t] + \eta$, proving (A4). \square

Proof of Proposition 8: (1) For any $A \in \mathcal{F}_t$, using (A3) twice, we obtain

$$\begin{aligned} \mathcal{E}[\mathbf{1}_A \xi \neg + \neg \mathbf{1}_{A^c} \eta | \mathcal{F}_t] & \neg \neg \neg = \neg \neg \neg \mathbf{1}_A \mathcal{E}[\mathbf{1}_A \xi \neg + \neg \mathbf{1}_{A^c} \eta | \mathcal{F}_t] \neg + \neg \\ & \mathbf{1}_{A^c} \mathcal{E}[\mathbf{1}_A \xi \neg + \neg \mathbf{1}_{A^c} \eta | \mathcal{F}_t] \neg = \neg \mathcal{E}[\mathbf{1}_A (\mathbf{1}_A \xi \neg \\ & + \neg \mathbf{1}_{A^c} \eta) | \mathcal{F}_t] \neg + \neg \mathcal{E}[\mathbf{1}_{A^c} (\mathbf{1}_A \xi \neg + \neg \\ & \mathbf{1}_{A^c} \eta) | \mathcal{F}_t] \\ & \neg \neg \neg = \neg \neg \neg \mathcal{E}[\mathbf{1}_A \xi | \mathcal{F}_t] + \mathcal{E}[\mathbf{1}_{A^c} \eta | \mathcal{F}_t] = \mathbf{1}_A \mathcal{E}[\xi | \mathcal{F}_t] + \\ & \mathbf{1}_{A^c} \mathcal{E}[\eta | \mathcal{F}_t], \quad a. s. \end{aligned}$$

(2) Applying (A3) with a null set A and $\xi = 0$, we obtain $\mathcal{E}[0|\mathcal{F}_t] = \mathcal{E}[\mathbf{1}_A 0|\mathcal{F}_t] = \mathbf{1}_A \mathcal{E}[0|\mathcal{F}_t] = 0$, a.s. If $\xi \in Dom_t(\mathcal{E})$, (A4) implies that $\mathcal{E}[\xi|\mathcal{F}_t] = \mathcal{E}[0 + \xi|\mathcal{F}_t] = \mathcal{E}[0|\mathcal{F}_t] + \xi = \xi$, a.s.

(3) If $\xi \leq \eta$, a.s., (A1) directly implies that for any $A \in \mathcal{F}_\nu$, $\mathcal{E}[\mathbf{1}_A \xi] \leq \mathcal{E}[\mathbf{1}_A \eta]$. On the other hand, suppose that $\mathcal{E}[\mathbf{1}_A \xi] \leq \mathcal{E}[\mathbf{1}_A \eta]$ for any $A \in \mathcal{F}_\nu$. We set $\tilde{A} \triangleq \{\xi > \eta\} \in \mathcal{F}_\nu$, thus $\mathbf{1}_{\tilde{A}} \xi \geq \mathbf{1}_{\tilde{A}} \eta \geq c \wedge 0$, a.s. Using (A1) we see that $\mathcal{E}[\mathbf{1}_{\tilde{A}} \xi] \geq \mathcal{E}[\mathbf{1}_{\tilde{A}} \eta]$; hence $\mathcal{E}[\mathbf{1}_{\tilde{A}} \xi] = \mathcal{E}[\mathbf{1}_{\tilde{A}} \eta]$. Then (A4) implies that

$$\mathcal{E}[\mathbf{1}_{\tilde{A}} \xi - c \wedge 0] = \mathcal{E}[\mathbf{1}_{\tilde{A}} \xi] - c \wedge 0 = \mathcal{E}[\mathbf{1}_{\tilde{A}} \eta] - c \wedge 0 = \mathcal{E}[\mathbf{1}_{\tilde{A}} \eta - c \wedge 0].$$

Applying the second part of (A1), we obtain that $\mathbf{1}_{\tilde{A}} \xi - c \wedge 0 = \mathbf{1}_{\tilde{A}} \eta - c \wedge 0$, a.s., which implies that $P(\tilde{A}) = 0$, i.e. $\xi \leq \eta$, a.s. \square

Proof of Proposition 24: We shall only consider the \mathcal{E} -supermartingale case, as the other cases can be deduced similarly. We first show that for any $s \in [0, T]$ and $\nu \in \mathcal{S}_{0,T}^F$

$$\mathcal{E}[X_\nu | \mathcal{F}_s] \leq X_{\nu \wedge s}, \quad a. s. \quad (8.1)$$

To see this, we note that since $\{\nu \leq s\} \in \mathcal{F}_s$, (A3) and Proposition 8 (2) imply that

$$\begin{aligned} \mathcal{E}[X_\nu | \mathcal{F}_s] &= \mathbf{1}_{\{\nu > s\}} \mathcal{E}[X_\nu | \mathcal{F}_s] + \mathbf{1}_{\{\nu \leq s\}} \mathcal{E}[X_\nu | \mathcal{F}_s] = \mathcal{E}[\mathbf{1}_{\{\nu > s\}} X_{\nu \vee s} | \mathcal{F}_s] + \\ &\quad \mathcal{E}[\mathbf{1}_{\{\nu \leq s\}} X_{\nu \wedge s} | \mathcal{F}_s] \\ &= \mathbf{1}_{\{\nu > s\}} \mathcal{E}[X_{\nu \vee s} | \mathcal{F}_s] + \mathbf{1}_{\{\nu \leq s\}} \mathcal{E}[X_{\nu \wedge s} | \mathcal{F}_s] = \\ &\quad \mathbf{1}_{\{\nu > s\}} \mathcal{E}[X_{\nu \vee s} | \mathcal{F}_s] + \mathbf{1}_{\{\nu \leq s\}} X_{\nu \wedge s}, \quad a. s. \end{aligned} \quad (8.2)$$

Suppose that $\nu_s \stackrel{\Delta}{=} \nu \vee s$ takes values in a finite subset $\{t_1 < \dots < t_n\}$ of $[s, T]$. Then (A4) implies that

$$\mathcal{E}[X_{\nu_s} | \mathcal{F}_{t_{n-1}}] = \mathcal{E}[\mathbf{1}_{\{\nu_s = t_n\}} X_{t_n} | \mathcal{F}_{t_{n-1}}] + \sum_{i=1}^{n-1} \mathbf{1}_{\{\nu_s = t_i\}} X_{t_i}, \quad a. s.$$

Since $\{\nu_s = t_n\} = \{\nu_s > t_{n-1}\} \in \mathcal{F}_{t_{n-1}}$, (A3) shows that

$$\mathcal{E}[\mathbf{1}_{\{\nu_s = t_n\}} X_{t_n} | \mathcal{F}_{t_{n-1}}] = \mathbf{1}_{\{\nu_s = t_n\}} \mathcal{E}[X_{t_n} | \mathcal{F}_{t_{n-1}}] \leq \mathbf{1}_{\{\nu_s = t_n\}} X_{t_{n-1}}, \quad a. s.$$

Thus it holds a.s. that $\mathcal{E}[X_{\nu_s} | \mathcal{F}_{t_{n-1}}] \leq \mathbf{1}_{\{\nu_s > t_{n-2}\}} X_{t_{n-1}} + \sum_{i=1}^{n-2} \mathbf{1}_{\{\nu_s = t_i\}} X_{t_i}$. Applying $\mathcal{E}[\cdot | \mathcal{F}_{t_{n-2}}]$ on both sides, we can further deduce from (A2)-(A4) that

$$\begin{aligned} \mathcal{E}[X_{\nu_s} | \mathcal{F}_{t_{n-2}}] &= \mathcal{E}[\mathcal{E}[X_{\nu_s} | \mathcal{F}_{t_{n-1}}] | \mathcal{F}_{t_{n-2}}] \leq \mathbf{1}_{\{\nu_s > t_{n-2}\}} \mathcal{E}[X_{t_{n-1}} | \mathcal{F}_{t_{n-2}}] + \\ &\quad \sum_{i=1}^{n-2} \mathbf{1}_{\{\nu_s = t_i\}} X_{t_i} \\ &\leq \mathbf{1}_{\{\nu_s > t_{n-2}\}} X_{t_{n-2}} + \sum_{i=1}^{n-2} \mathbf{1}_{\{\nu_s = t_i\}} X_{t_i} = \\ &\quad \mathbf{1}_{\{\nu_s > t_{n-3}\}} X_{t_{n-2}} + \sum_{i=1}^{n-3} \mathbf{1}_{\{\nu_s = t_i\}} X_{t_i}, \quad a. s. \end{aligned}$$

Inductively, it follows that $\mathcal{E}[X_{\nu_s} | \mathcal{F}_{t_1}] \leq X_{t_1}$, a.s. Applying (A2) once again, we obtain

$$\mathcal{E}[X_{\nu_s} | \mathcal{F}_s] = \mathcal{E}[\mathcal{E}[X_{\nu_s} | \mathcal{F}_{t_1}] | \mathcal{F}_s] \leq \mathcal{E}[X_{t_1} | \mathcal{F}_s] \leq X_s, \quad a. s.,$$

which together with (8.1) implies that

$$\mathcal{E}[X_\nu | \mathcal{F}_s] \leq \mathbf{1}_{\{\nu > s\}} X_s + \mathbf{1}_{\{\nu \leq s\}} X_{\nu \wedge s} = X_{\nu \wedge s}, \quad a. s., \quad \langle \text{hbox} \mid \text{proving(8.1)} \rangle.$$

Let $\sigma \in \mathcal{S}_{0,T}^F$ taking values in a finite set $\{s_1 < \dots < s_m\}$, then

$$\begin{aligned} \mathcal{E}[X_\nu | \mathcal{F}_\sigma] &= \mathcal{E}[\mathcal{E}[X_\nu | \mathcal{F}_{s_j}] | \mathcal{F}_\sigma] \leq \\ &\quad \sum_{j=1}^m \mathbf{1}_{\{\sigma = s_j\}} X_{\nu \wedge s_j} = X_{\nu \wedge \sigma}, \\ &\quad a. s. \end{aligned} \quad \langle \text{hbox} \mid \square \rangle$$

Proof of Proposition 26: Given $\xi \in Dom(\mathcal{E})$, we let $\nu \in \mathcal{S}_{0,T}^F$ take values in a finite set $\{t_1 < \dots < t_n\}$.

1) For any $\eta \in Dom(\mathcal{E})$ with $\xi \leq \eta$, a. s., (A1) implies that

$$\mathcal{E}[\xi|\mathcal{F}_\nu] = \sum_{i=1}^n \mathbf{1}_{\{\nu=t_i\}} \mathcal{E}[\xi|\mathcal{F}_{t_i}] \leq \sum_{i=1}^n \mathbf{1}_{\{\nu=t_i\}} \mathcal{E}[\eta|\mathcal{F}_{t_i}] = \mathcal{E}[\eta|\mathcal{F}_\nu], \quad a. s.$$

Moreover, if $0 \leq \xi \leq \eta$, a.s. and $\mathcal{E}[\xi|\mathcal{F}_\sigma] = \mathcal{E}[\eta|\mathcal{F}_\sigma]$, a.s. for some $\sigma \in \mathcal{S}_{0,T}^F$, we can apply Corollary 25 to obtain

$$\mathcal{E}[\xi] = \mathcal{E}[\mathcal{E}[\xi|\mathcal{F}_\sigma]] = \mathcal{E}[\mathcal{E}[\eta|\mathcal{F}_\sigma]] = \mathcal{E}[\eta].$$

The second part of (A1) then implies that $\xi = \eta$, a.s., proving (1).

2) For any $A \in \mathcal{F}_\nu$, it is clear that $A \cap \{\nu = t_i\} \in \mathcal{F}_{t_i}$ for each $i \in \{1, \dots, n\}$. Hence we can deduce from (A3) that

$$\begin{aligned} \mathcal{E}[\mathbf{1}_A \xi|\mathcal{F}_\nu] &= \sum_{i=1}^n \mathbf{1}_{\{\nu=t_i\}} \mathcal{E}[\mathbf{1}_A \xi|\mathcal{F}_{t_i}] = \sum_{i=1}^n \mathcal{E}[\mathbf{1}_{\{\nu=t_i\} \cap A} \xi|\mathcal{F}_{t_i}] = \\ &= \sum_{i=1}^n \mathbf{1}_{\{\nu=t_i\} \cap A} \mathcal{E}[\xi|\mathcal{F}_{t_i}] \\ &= \mathbf{1}_A \sum_{i=1}^n \mathbf{1}_{\{\nu=t_i\}} \mathcal{E}[\xi|\mathcal{F}_{t_i}] = \mathbf{1}_A \mathcal{E}[\xi|\mathcal{F}_\nu], \quad a. s., \quad \langle \text{hbox |} \\ &\quad \text{proving(2).} \rangle \end{aligned}$$

3) For any $\eta \in Dom_\nu(\mathcal{E})$, since $\mathbf{1}_{\{\nu=t_i\}} \eta \in Dom_{t_i}(\mathcal{E})$ for each $i \in \{1, \dots, n\}$, (A3) and (A4) imply that

$$\begin{aligned} \mathcal{E}[\xi + \eta|\mathcal{F}_\nu] &= \sum_{i=1}^n \mathbf{1}_{\{\nu=t_i\}} \mathcal{E}[\xi + \eta|\mathcal{F}_{t_i}] = \sum_{i=1}^n \mathcal{E}[\mathbf{1}_{\{\nu=t_i\}} \xi + \mathbf{1}_{\{\nu=t_i\}} \eta|\mathcal{F}_{t_i}] = \\ &= \sum_{i=1}^n (\mathcal{E}[\mathbf{1}_{\{\nu=t_i\}} \xi|\mathcal{F}_{t_i}] + \mathbf{1}_{\{\nu=t_i\}} \eta) \\ &= \sum_{i=1}^n \mathbf{1}_{\{\nu=t_i\}} \mathcal{E}[\xi|\mathcal{F}_{t_i}] + \eta = \mathcal{E}[\xi|\mathcal{F}_\nu] + \eta, \quad a. s., \quad \langle \text{hbox |} \\ &\quad \text{proving(3).} \rangle \end{aligned}$$

The proof of (4) and (5) is similar to that of Proposition 8 (1) and (2) by applying the just obtained "Zero-one Law" and "Translation Invariance". \square

Proof of Theorem 16: (H1) is an easy consequence of the lower semi-continuity (2.2). In fact, for any $\xi \in Dom^+(\mathcal{E})$ and any $\{A_n\}_{n \in \mathbb{N}} \subset \mathcal{F}_T$ with $\lim_{n \rightarrow \infty} \neg \uparrow \mathbf{1}_{A_n} = 1$ a.s., $\{\mathbf{1}_{A_n} \xi\}_{n \in \mathbb{N}}$ is an increasing sequence converging to ξ . Then applying the lower semi-continuity with $\nu = 0$ and using (A1), we obtain $\mathcal{E}[\xi] \leq \lim_{n \rightarrow \infty} \neg \uparrow \mathcal{E}[\mathbf{1}_{A_n} \xi] \leq \mathcal{E}[\xi]$; so (H1) follows.

On the other hand, to show that (H1) implies the lower semi-continuity, we first extend (H1) as follows: For any $\xi \in Dom^+(\mathcal{E})$ and any $\{A_n\}_{n \in \mathbb{N}} \subset \mathcal{F}_T$ with $\lim_{n \rightarrow \infty} \neg \uparrow \mathbf{1}_{A_n} = 1$, a.s., it holds for any $t \in [0, T]$ that

$$\lim_{n \rightarrow \infty} \neg \uparrow \mathcal{E}[\mathbf{1}_{A_n} \xi|\mathcal{F}_t] = \mathcal{E}[\xi|\mathcal{F}_t], \quad a. s. \quad (8.3)$$

In fact, by (A1), it holds a.s. that $\{\mathcal{E}[\mathbf{1}_{A_n}\xi|\mathcal{F}_t]\}_{n \in \mathbb{N}}$ is an increasing sequence bounded from above by $\mathcal{E}[\xi|\mathcal{F}_t]$. Hence, $\lim_{n \rightarrow \infty} \neg \uparrow \mathcal{E}[\mathbf{1}_{A_n}\xi|\mathcal{F}_t] \leq \mathcal{E}[\xi|\mathcal{F}_t]$, a.s. Assuming that $\lim_{n \rightarrow \infty} \neg \uparrow \mathcal{E}[\mathbf{1}_{A_n}\xi|\mathcal{F}_t] < \mathcal{E}[\xi|\mathcal{F}_t]$ with a positive probability, we can find an $\varepsilon > 0$ such that the set $A_\varepsilon = \left\{ \lim_{n \rightarrow \infty} \neg \uparrow \mathcal{E}[\mathbf{1}_{A_n}\xi|\mathcal{F}_t] \leq \mathcal{E}[\xi|\mathcal{F}_t] - \varepsilon \right\} \in \mathcal{F}_t$ still has positive probability. Hence for any $n \in \mathbb{N}$, we have

$$\mathbf{1}_{A_\varepsilon} \mathcal{E}[\mathbf{1}_{A_n}\xi|\mathcal{F}_t] \leq \mathbf{1}_{A_\varepsilon} \lim_{n \rightarrow \infty} \neg \uparrow \mathcal{E}[\mathbf{1}_{A_n}\xi|\mathcal{F}_t] \leq \mathbf{1}_{A_\varepsilon} (\mathcal{E}[\xi|\mathcal{F}_t] - \varepsilon), \quad a.s.$$

Then (A1)-(A4) imply that

$$\begin{aligned} \mathcal{E}[\mathbf{1}_{A_\varepsilon}\mathbf{1}_{A_n}\xi] + \varepsilon &= \mathcal{E}[\mathbf{1}_{A_\varepsilon}\mathbf{1}_{A_n}\xi + \varepsilon] = \mathcal{E}[\mathcal{E}[\mathbf{1}_{A_\varepsilon}\mathbf{1}_{A_n}\xi + \varepsilon|\mathcal{F}_t]] = \\ &= \mathcal{E}[\mathbf{1}_{A_\varepsilon}\mathcal{E}[\mathbf{1}_{A_n}\xi|\mathcal{F}_t] + \varepsilon] \\ &\leq \mathcal{E}[\mathbf{1}_{A_\varepsilon}\mathcal{E}[\xi|\mathcal{F}_t] + \varepsilon\mathbf{1}_{A_\varepsilon}] = \mathcal{E}[\mathcal{E}[\mathbf{1}_{A_\varepsilon}\xi + \varepsilon\mathbf{1}_{A_\varepsilon}|\mathcal{F}_t]] = \\ &= \mathcal{E}[\mathbf{1}_{A_\varepsilon}\xi + \varepsilon\mathbf{1}_{A_\varepsilon}]. \end{aligned}$$

Using (A4), (H1) and (A1), we obtain

$$\mathcal{E}[\mathbf{1}_{A_\varepsilon}\xi + \varepsilon] = \mathcal{E}[\mathbf{1}_{A_\varepsilon}\xi] + \varepsilon = \lim_{n \rightarrow \infty} \neg \uparrow \mathcal{E}[\mathbf{1}_{A_n}\mathbf{1}_{A_\varepsilon}\xi] + \varepsilon \leq \mathcal{E}[\mathbf{1}_{A_\varepsilon}\xi + \varepsilon\mathbf{1}_{A_\varepsilon}] \leq \mathcal{E}[\mathbf{1}_{A_\varepsilon}\xi + \varepsilon],$$

thus $\mathcal{E}[\mathbf{1}_{A_\varepsilon}\xi + \varepsilon] = \mathcal{E}[\mathbf{1}_{A_\varepsilon}\xi + \varepsilon\mathbf{1}_{A_\varepsilon}]$. Then the second part of (A1) implies that $\mathbf{1}_{A_\varepsilon}\xi + \varepsilon = \mathbf{1}_{A_\varepsilon}\xi + \varepsilon\mathbf{1}_{A_\varepsilon}$, a.s., which can hold only if $P(A_\varepsilon) = 0$. This results in a contradiction. Thus $\lim_{n \rightarrow \infty} \neg \uparrow \mathcal{E}[\mathbf{1}_{A_n}\xi|\mathcal{F}_t] = \mathcal{E}[\xi|\mathcal{F}_t]$, a.s., proving (8.3).

Next, we show that (2.2) holds for each deterministic stopping time $\nu = t \in [0, T]$. For any $j, n \in \mathbb{N}$, we define $A_n^j \triangleq \cap_{k=n}^\infty \{|\xi - \xi_k| < 1/j\} \in \mathcal{F}_T$. (A1) and (A4) imply that for any $k \geq n$

$$\mathcal{E}[\mathbf{1}_{A_n^j}\xi|\mathcal{F}_t] \leq \mathcal{E}[\mathbf{1}_{\{|\xi - \xi_k| < 1/j\}}\xi|\mathcal{F}_t] \leq \mathcal{E}[\xi_k + 1/j|\mathcal{F}_t] = \mathcal{E}[\xi_k|\mathcal{F}_t] + 1/j, \quad a.s.$$

Hence, except on a null set N_n^j , the above inequality holds for any $k \geq n$. As $k \rightarrow \infty$, it holds on $(N_n^j)^c$ that

$$\mathcal{E}[\mathbf{1}_{A_n^j}\xi|\mathcal{F}_t] \leq \liminf_{k \rightarrow \infty} \mathcal{E}[\xi_k|\mathcal{F}_t] + 1/j.$$

(Here it is not necessary that $\liminf_{k \rightarrow \infty} \mathcal{E}[\xi_k|\mathcal{F}_t] < \infty$, a.s.) Since $\xi_n \rightarrow \xi$, a.s. as $n \rightarrow \infty$, it is clear that $\lim_{n \rightarrow \infty} \neg \uparrow \mathbf{1}_{A_n^j} = 1$, a.s. Then (8.3) implies that $\mathcal{E}[\xi|\mathcal{F}_t] = \lim_{n \rightarrow \infty} \neg \uparrow \mathcal{E}[\mathbf{1}_{A_n^j}\xi|\mathcal{F}_t]$ holds except on a null set N_0^j . Let $N^j = \cup_{n=0}^\infty N_n^j$. It then holds on $(N^j)^c$ that

$$\mathcal{E}[\xi|\mathcal{F}_t] = \lim_{n \rightarrow \infty} \neg \uparrow \mathcal{E}[\mathbf{1}_{A_n^j}\xi|\mathcal{F}_t] \leq \liminf_{k \rightarrow \infty} \mathcal{E}[\xi_k|\mathcal{F}_t] + 1/j.$$

As $j \rightarrow \infty$, it holds except on the null set $\cup_{j=1}^\infty N^j$ that

$$\mathcal{E}[\xi|\mathcal{F}_t] \leq \liminf_{n \rightarrow \infty} \mathcal{E}[\xi_n|\mathcal{F}_t]. \quad (8.4)$$

Let $\nu \in \mathcal{S}_{0,T}^F$ taking values in a finite set $\{t_1 < \dots < t_n\}$. Then we can deduce from (8.4) that

$$\begin{aligned} \mathcal{E}[\xi|\mathcal{F}_\nu] &= \sum_{i=1}^n \mathbf{1}_{\{\nu=t_i\}} \mathcal{E}[\xi|\mathcal{F}_{t_i}] \leq \sum_{i=1}^n \mathbf{1}_{\{\nu=t_i\}} \liminf_{n \rightarrow \infty} \mathcal{E}[\xi_n|\mathcal{F}_{t_i}] = \neg \\ \liminf_{n \rightarrow \infty} \sum_{i=1}^n \mathbf{1}_{\{\nu=t_i\}} \mathcal{E}[\xi_n|\mathcal{F}_{t_i}] &= \liminf_{n \rightarrow \infty} \mathcal{E}[\xi_n|\mathcal{F}_\nu], \quad a.s., \end{aligned} \quad (8.5)$$

which completes the proof. \square

Proof of Theorem 18: We first show an extension of (H2): For any ξ , $\eta \in Dom^+(\mathcal{E})$ and any $\{A_n\}_{n \in \mathbb{N}} \subset \mathcal{F}_T$ with $\lim_{n \rightarrow \infty} \neg \mathbf{1}_{A_n} = 0$, a.s., it holds a.s. that

$$\lim_{n \rightarrow \infty} \neg \mathcal{E}[\xi + \mathbf{1}_{A_n} \eta|\mathcal{F}_t] = \mathcal{E}[\xi|\mathcal{F}_t], \quad a.s. \quad (8.6)$$

In fact, by (A1), it holds a.s. that $\{\mathcal{E}[\xi + \mathbf{1}_{A_n} \eta|\mathcal{F}_t]\}_{n \in \mathbb{N}}$ is a decreasing sequence bounded from below by $\mathcal{E}[\xi|\mathcal{F}_t]$. Hence, $\lim_{n \rightarrow \infty} \neg \mathcal{E}[\xi + \mathbf{1}_{A_n} \eta|\mathcal{F}_t] \geq \mathcal{E}[\xi|\mathcal{F}_t]$, a.s. Assume that $\lim_{n \rightarrow \infty} \neg \mathcal{E}[\xi + \mathbf{1}_{A_n} \eta|\mathcal{F}_t] > \mathcal{E}[\xi|\mathcal{F}_t]$ with a positive probability, then we can find an $\varepsilon > 0$ such that the set $A'_\varepsilon = \left\{ \lim_{n \rightarrow \infty} \neg \mathcal{E}[\xi + \mathbf{1}_{A_n} \eta|\mathcal{F}_t] \geq \mathcal{E}[\xi|\mathcal{F}_t] + \varepsilon \right\} \in \mathcal{F}_t$ still has positive probability. For any $n \in \mathbb{N}$, (A4) implies that

$$\begin{aligned} \mathbf{1}_{A'_\varepsilon} \mathcal{E}[\xi + \mathbf{1}_{A_n} \eta|\mathcal{F}_t] &\geq \mathbf{1}_{A'_\varepsilon} \lim_{n \rightarrow \infty} \neg \mathcal{E}[\xi + \mathbf{1}_{A_n} \eta|\mathcal{F}_t] \geq \mathbf{1}_{A'_\varepsilon} (\mathcal{E}[\xi|\mathcal{F}_t] + \varepsilon) = \\ \mathbf{1}_{A'_\varepsilon} \mathcal{E}[\xi + \varepsilon|\mathcal{F}_t], \quad a.s. \end{aligned}$$

Applying (A1)-(A3), we obtain

$$\begin{aligned} \mathcal{E}[\mathbf{1}_{A'_\varepsilon} \xi + \mathbf{1}_{A_n} \mathbf{1}_{A'_\varepsilon} \eta] &= \mathcal{E}[\mathcal{E}[\mathbf{1}_{A'_\varepsilon} \xi + \mathbf{1}_{A_n} \mathbf{1}_{A'_\varepsilon} \eta|\mathcal{F}_t]] = \mathcal{E}[\mathbf{1}_{A'_\varepsilon} \mathcal{E}[\xi + \\ &\quad \mathbf{1}_{A_n} \eta|\mathcal{F}_t]] \geq \mathcal{E}[\mathbf{1}_{A'_\varepsilon} \mathcal{E}[\xi + \varepsilon|\mathcal{F}_t]] \\ &= \mathcal{E}[\mathcal{E}[\mathbf{1}_{A'_\varepsilon} (\xi + \varepsilon)|\mathcal{F}_t]] = \mathcal{E}[\mathbf{1}_{A'_\varepsilon} (\xi + \varepsilon)]. \end{aligned}$$

Thanks to (H2) we further have

$$\mathcal{E}[\mathbf{1}_{A'_\varepsilon} \xi] = \lim_{n \rightarrow \infty} \neg \mathcal{E}[\mathbf{1}_{A'_\varepsilon} \xi + \mathbf{1}_{A_n} \mathbf{1}_{A'_\varepsilon} \eta] \geq \mathcal{E}[\mathbf{1}_{A'_\varepsilon} (\xi + \varepsilon)] \geq \mathcal{E}[\mathbf{1}_{A'_\varepsilon} \xi],$$

thus $\mathcal{E}[\mathbf{1}_{A'_\varepsilon} \xi] = \mathcal{E}[\mathbf{1}_{A'_\varepsilon} (\xi + \varepsilon)]$. Then the second part of (A1) implies that $P(A'_\varepsilon) = 0$, which yields a contradiction. Therefore, $\lim_{n \rightarrow \infty} \neg \mathcal{E}[\xi + \mathbf{1}_{A_n} \eta|\mathcal{F}_t] = \mathcal{E}[\xi|\mathcal{F}_t]$, a.s., proving (8.6).

Since the sequence $\{\xi_n\}_{n \in \mathbb{N}}$ is bounded above by η , it holds a.s. that $\xi = \lim_{n \rightarrow \infty} \xi_n \leq \eta$, thus (D3) implies that $\xi \in Dom(\mathcal{E})$. Then Fatou's Lemma (Theorem 16) implies that for any $\nu \in \mathcal{S}_{0,T}^F$,

$$\mathcal{E}[\xi|\mathcal{F}_\nu] \leq \liminf_{n \rightarrow \infty} \mathcal{E}[\xi_n|\mathcal{F}_\nu], \quad a.s. \quad (8.7)$$

On the other hand, we first fix $t \in [0, T]$. For any $j, n \in \mathbb{N}$, define $A_n^j \triangleq \cap_{k=n}^\infty \{|\xi - \xi_k| < 1/j\} \in \mathcal{F}_T$. Then one can deduce that for any $k \geq n$

$$\begin{aligned} \mathcal{E}[\xi_k|\mathcal{F}_t] &\leq \mathcal{E}[\mathbf{1}_{A_n^j} (\xi + 1/j) + \mathbf{1}_{(A_n^j)^c} \eta|\mathcal{F}_t] \leq \mathcal{E}[\xi + 1/j + \mathbf{1}_{(A_n^j)^c} (\eta - \xi)|\mathcal{F}_t], \\ a.s. \end{aligned}$$

Hence, except on a null set N_n^j , the above inequality holds for any $k \geq n$. As $k \rightarrow \infty$, it holds on $(N_n^j)^c$ that

$$\limsup_{k \rightarrow \infty} \mathcal{E}[\xi_k | \mathcal{F}_t] \leq \mathcal{E}[\xi + 1/j + \mathbf{1}_{(A_n^j)^c}(\eta - \xi) | \mathcal{F}_t].$$

Since $\xi \in L^0(\mathcal{F}_T)$ and $\xi_n \rightarrow \xi$, a.s. as $n \rightarrow \infty$, it is clear that $\lim_{n \rightarrow \infty} \neg \neg \uparrow \mathbf{1}_{A_n^j} = 1$, a.s. Then (8.6) and (A4) imply that except on a null set N_0^j , we have

$$\lim_{n \rightarrow \infty} \neg \downarrow \mathcal{E}[\xi + 1/j + \mathbf{1}_{(A_n^j)^c}(\eta - \xi) | \mathcal{F}_t] = \mathcal{E}[\xi + 1/j | \mathcal{F}_t] = \mathcal{E}[\xi | \mathcal{F}_t] + 1/j.$$

Let $N^j = \cup_{n=0}^{\infty} N_n^j$, thus it holds on $(N^j)^c$ that

$$\limsup_{k \rightarrow \infty} \mathcal{E}[\xi_k | \mathcal{F}_t] \leq \mathcal{E}[\xi | \mathcal{F}_t] + 1/j.$$

As $j \rightarrow \infty$, it holds except on the null set $\cup_{j=1}^{\infty} N^j$ that $\limsup_{n \rightarrow \infty} \mathcal{E}[\xi_n | \mathcal{F}_t] \leq \mathcal{E}[\xi | \mathcal{F}_t]$. Then for any $\nu \in \mathcal{S}_{0,T}^F$, using an argument similar to (8.5), we can deduce that

$$\limsup_{n \rightarrow \infty} \mathcal{E}[\xi_n | \mathcal{F}_\nu] \leq \mathcal{E}[\xi | \mathcal{F}_\nu], \quad a.s.,$$

which together with (8.7) proves the theorem. \square

Proof of Theorem 20: Let $F = \{t_1 < t_2 < \dots < t_d\}$ be any finite subset of \mathcal{D}_T . For $j = 1, \dots, d$, we define $A_j = \{\nu_j < T\} \in \mathcal{F}_{\nu_j}$, clearly, $A_j \supset A_{j+1}$. Let $d' = \lfloor \frac{d}{2} \rfloor$, one can deduce that $U_F(a, b; X) = \sum_{j=1}^{d'} \mathbf{1}_{A_{2j}}$ and that

$$\mathbf{1}_{\cup_{j=1}^{d'} (A_{2j-1} \setminus A_{2j})} (X_T - a) \geq \mathbf{1}_{\cup_{j=1}^{d'} (A_{2j-1} \setminus A_{2j})} \mathbf{1}_{\{X_T < a\}} (X_T - a) \geq \mathbf{1}_{\{X_T < a\}} (X_T - a) = -(a - X_T)^+.$$

Since $X_T \in \text{Dom}(\mathcal{E})$ and $L^\infty(\mathcal{F}_T) \subset \text{Dom}(\mathcal{E})$ (by Lemma 19), we can deduce from (D2) that

$$(b - a) U_F(a, b; X) - (a - X_T)^+ = \sum_{j=1}^{d'} \mathbf{1}_{A_{2j}} (b - a) + \mathbf{1}_{\{X_T < a\}} (X_T - a) \in \text{Dom}(\mathcal{E}).$$

Then Proposition 26 (1)-(3) and Proposition 24 imply that

$$\begin{aligned} \mathcal{E}[(b - a) U_F(a, b; X) - (a - X_T)^+ | \mathcal{F}_{\nu_{2d'}}] &\leq (b - a) \sum_{j=1}^{d'} \mathbf{1}_{A_{2j}} + \\ &\quad \mathcal{E}[\mathbf{1}_{\cup_{j=1}^{d'} (A_{2j-1} \setminus A_{2j})} (X_T - a) | \mathcal{F}_{\nu_{2d'}}] \\ &= \neg \neg \neg (b - a) \sum_{j=1}^{d'} \mathbf{1}_{A_{2j}} \neg + \mathbf{1}_{\cup_{j=1}^{d'} (A_{2j-1} \setminus A_{2j})} (\mathcal{E}[X_T | \mathcal{F}_{\nu_{2d'}}] - a) \leq (b - \\ &\quad a) \sum_{j=1}^{d'} \mathbf{1}_{A_{2j}} \neg + \mathbf{1}_{\cup_{j=1}^{d'} (A_{2j-1} \setminus A_{2j})} (X_{\nu_{2d'}} \neg - a), \quad a.s. \end{aligned} \quad (8.8)$$

Applying $\mathcal{E}[\cdot|\mathcal{F}_{\nu_{2d'-1}}]$ to the above inequality, using Proposition 26 (1)-(3) and Proposition 24 once again, we obtain

$$\begin{aligned}
& \mathcal{E}[(b-a)U_F(a,b;X) - (a-X_T)^+|\mathcal{F}_{\nu_{2d'-1}}] \\
& \leq \mathcal{E}\left[(b-a)\sum_{j=1}^{d'-1}\mathbf{1}_{A_{2j}} + (\mathbf{1}_{A_{2d'-1}} + \mathbf{1}_{\cup_{j=1}^{d'-1}(A_{2j-1}\setminus A_{2j})})(X_{\nu_{2d'}} - a)\middle|\mathcal{F}_{\nu_{2d'-1}}\right] \\
& = (b-a)\sum_{j=1}^{d'-1}\mathbf{1}_{A_{2j}} + \mathcal{E}[(\mathbf{1}_{A_{2d'-1}} + \mathbf{1}_{\cup_{j=1}^{d'-1}(A_{2j-1}\setminus A_{2j})})(X_{\nu_{2d'}} - a)|\mathcal{F}_{\nu_{2d'-1}}] \\
& = (b-a)\sum_{j=1}^{d'-1}\mathbf{1}_{A_{2j}} + (\mathbf{1}_{A_{2d'-1}} + \mathbf{1}_{\cup_{j=1}^{d'-1}(A_{2j-1}\setminus A_{2j})})(\mathcal{E}[X_{\nu_{2d'}}|\mathcal{F}_{\nu_{2d'-1}}] - a) \\
& \leq (b-a)\sum_{j=1}^{d'-1}\mathbf{1}_{A_{2j}} + (\mathbf{1}_{A_{2d'-1}} + \mathbf{1}_{\cup_{j=1}^{d'-1}(A_{2j-1}\setminus A_{2j})})(X_{\nu_{2d'-1}} - a) \\
& \leq (b-a)\sum_{j=1}^{d'-1}\mathbf{1}_{A_{2j}} + \mathbf{1}_{\cup_{j=1}^{d'-1}(A_{2j-1}\setminus A_{2j})}(X_{\nu_{2d'-1}} - a), \quad a.s.,
\end{aligned} \tag{8.9}$$

where we used the fact that $X_{\nu_{2d'}} > b$ on $A_{2d'}$ in the first inequality and the fact that $X_{\nu_{2d'-1}} < a$ on $A_{2d'-1}$ in the last inequality. Similarly, applying $\mathcal{E}[\cdot|\mathcal{F}_{\nu_{2d'-2}}]$ to the above inequality yields that

$$\begin{aligned}
& \mathcal{E}[(b-a)U_F(a,b;X) - (a-X_T)^+|\mathcal{F}_{\nu_{2d'-2}}] \leq (b-a)\sum_{j=1}^{d'-1}\mathbf{1}_{A_{2j}} + \\
& \mathbf{1}_{\cup_{j=1}^{d'-1}(A_{2j-1}\setminus A_{2j})}(X_{\nu_{2d'-2}} - a), \quad a.s.
\end{aligned}$$

Iteratively applying $\mathcal{E}[\cdot|\mathcal{F}_{\nu_{2d'-3}}]$, $\mathcal{E}[\cdot|\mathcal{F}_{\nu_{2d'-4}}]$ and so on, we eventually obtain that

$$\mathcal{E}[(b-a)U_F(a,b;X) - (a-X_T)^+] \leq 0. \tag{8.10}$$

We assume first that $X_T \geq c$, a.s. for some $c \in \mathbb{R}$. Since $(a-X_T)^+ \leq \neg|a| + |c|$, it directly follows from (A4) that

$$0 \geq \mathcal{E}[(b-a)U_F(a,b;X) - (a-X_T)^+] \geq \mathcal{E}[(b-a)U_F(a,b;X)] - (|a| + |c|). \tag{8.11}$$

Let $\{F_n\}_{n \in \mathbb{N}}$ be an increasing sequence of finite subsets of \mathcal{D}_T with $\cup_{n \in \mathbb{N}} F_n = \mathcal{D}_T$, thus $\lim_{n \rightarrow \infty} \neg \uparrow U_{F_n}(a, b; X) = U_{\mathcal{D}_T}(a, b; X)$. Fix $M \in \mathbb{N}$, we see that

$$\lim_{n \rightarrow \infty} \neg \uparrow \mathbf{1}\mathbf{1}_{\{U_{F_n}(a, b; X) > M\}} = \mathbf{1}\mathbf{1}_{\cup_n \{U_{F_n}(a, b; X) > M\}} = \mathbf{1}\mathbf{1}_{\{U_{\mathcal{D}_T}(a, b; X) > M\}}. \quad (8.12)$$

For any $n \in \mathbb{N}$, we know from (8.11) that $\mathcal{E}[(b-a) M \mathbf{1}\mathbf{1}_{\{U_{F_n}(a, b; X) > M\}}] \leq \mathcal{E}[(b-a) U_{F_n}(a, b; X)] \leq |a| + |c|$, thus Fatou's Lemma (Theorem 16) implies that

$$\begin{aligned} \mathcal{E}[(b-a) M \mathbf{1}\mathbf{1}_{\{U_{\mathcal{D}_T}(a, b; X) = \infty\}}] &\leq \mathcal{E}[(b-a) M \mathbf{1}\mathbf{1}_{\{U_{\mathcal{D}_T}(a, b; X) > M\}}] \\ &\leq \lim_{n \rightarrow \infty} \neg \uparrow \mathcal{E}[(b-a) M \mathbf{1}\mathbf{1}_{\{U_{F_n}(a, b; X) > M\}}] \leq \\ &|a| + |c|. \end{aligned} \quad (8.13)$$

On the other hand, if $\mathcal{E}[\cdot]$ is concave, then we can deduce from (8.10) that

$$0 \geq \mathcal{E}[(b-a) U_F(a, b; X) - (a - X_T)^+] \geq \frac{1}{2} \mathcal{E}[2(b-a) U_F(a, b; X)] + \frac{1}{2} \mathcal{E}[-2(a - X_T)^+].$$

Mimicking the arguments in (8.12) and (8.12), we obtain that

$$\mathcal{E}[(b-a) 2 M \mathbf{1}\mathbf{1}_{\{U_{\mathcal{D}_T}(a, b; X) = \infty\}}] \leq -\mathcal{E}[-2(a - X_T)^+].$$

where $-2(a - X_T)^+ = \mathbf{1}\mathbf{1}_{\{X_T < a\}} 2(X_T - a) \in \text{Dom}(\mathcal{E})$ thanks to (D2). Also note that (A1) and Proposition 26 (5) imply that $\mathcal{E}[-2(a - X_T)^+] \leq \mathcal{E}[0] = 0$.

Using (H0) in both cases above yields that $P(U_{\mathcal{D}_T}(a, b; X) = \infty) = 0$, i.e., $U_{\mathcal{D}_T}(a, b; X) < \infty$, a.s. Then a classical argument (see e.g. (, ?, Proposition 1.3.14)) shows that

$$P\left(\langle \text{hbox|both} \rangle \lim_{s \nearrow t, s \in \mathcal{D}_T} X_s \langle \text{hbox|and} \rangle \lim_{s \searrow t, s \in \mathcal{D}_T} X_s \langle \text{hbox|exist for any} \rangle t \in [0, T] \right) = 1.$$

This completes the proof. \square

Proof of Proposition 82: We can deduce from (2.4) that except on a null set N

$$\begin{aligned} X_t^+ &= \lim_{n \rightarrow \infty} X_{q_n^+(t)} \leq \text{esssup}_{s \in \mathcal{D}_T} X_s \langle \text{hbox|for any} \rangle t \in \\ &[0, T], \end{aligned} \quad (8.14)$$

$$\langle \text{error|compound hbox} \rangle X_\nu^+ = \lim_{n \rightarrow \infty} X_{q_n^+(\nu)} \leq \text{esssup}_{s \in \mathcal{D}_T} X_s \langle \text{hbox|for any} \rangle \nu \in \mathcal{S}_{0,T}. \quad (8.15)$$

Proof of (1): Case I. For any $\nu \in \mathcal{S}_{0,T}$, if $\text{esssup}_{s \in \mathcal{D}_T} X_s \in \text{Dom}^+(\mathcal{E})$, (D3) and (8.15) directly imply that X_ν^+ belongs to $\text{Dom}(\mathcal{E})$.

Case II. Assume that \mathcal{E} satisfies (8.6). For any $n \in \mathbb{N}$, since X is an \mathcal{E} -supermartingale and since $q_n^+(\nu) \in \mathcal{S}_{0,T}^F$, Corollary 25 and Proposition 24 imply that

$$\mathcal{E}[X_{q_n^+(\nu)}] = \mathcal{E}[\mathcal{E}[X_{q_n^+(\nu)} | \mathcal{F}_{q_{n+1}^+(\nu)}]] \leq \mathcal{E}[X_{q_{n+1}^+(\nu)}] \leq X_0.$$

Hence, $\{\mathcal{E}[X_{q_n^+}^+]\}_{n \in \mathbb{N}}$ is an increasing non-negative sequence that is bounded from above by $X_0 \in [0, \infty)$. (8.6) and (8.15) then imply that X_ν^+ belongs to $\text{Dom}(\mathcal{E})$, proving statement (1).

Proof of (2): Now suppose that $X_t^+ \in \text{Dom}^+(\mathcal{E})$ for any $t \in [0, T]$. First, we shall show that for $t \in [0, T]$ and $A \in \mathcal{F}_t$

$$\mathcal{E}[\mathbf{1}_A X_t^+] = \lim_{n \rightarrow \infty} \mathcal{E}[\mathbf{1}_A X_{q_n^+}^+]. \quad (8.16)$$

Since the distribution function $x \mapsto P\{X_t^+ \leq x\}$ jumps up at most on a countable subset S of $[0, \infty)$, we can find a sequence $\{K_j\}_{j=1}^\infty \subset [0, \infty) \setminus S$ increasing to ∞ . Fix $m, j \in \mathbb{N}$, (A1)-(A3) imply that for any $n \geq m$

$$\begin{aligned} \mathcal{E}[\mathbf{1}_A \mathbf{1}_{\{X_{q_n^+}^+ < K_j\}} (X_{q_n^+}^+ \wedge K_j)] &= \mathcal{E}[\mathbf{1}_A \mathbf{1}_{\{X_{q_n^+}^+ < K_j\}} X_{q_n^+}^+] \geq \\ &\mathcal{E}[\mathbf{1}_A \mathbf{1}_{\{X_{q_n^+}^+ < K_j\}} \mathcal{E}[X_{q_n^+}^+ | \mathcal{F}_{q_n^+}]] \\ &= \mathcal{E}[\mathcal{E}[\mathbf{1}_A \mathbf{1}_{\{X_{q_n^+}^+ < K_j\}} X_{q_n^+}^+ | \mathcal{F}_{q_n^+}]] = \\ &\mathcal{E}[\mathbf{1}_A \mathbf{1}_{\{X_{q_n^+}^+ < K_j\}} X_{q_n^+}^+]. \end{aligned} \quad (8.17)$$

Since $K_j \notin S$, $P\{X_t^+ = K_j\} = 0$, one can easily deduce from (8.14) that $\lim_{n \rightarrow \infty} \mathbf{1}_{\{X_{q_n^+}^+ < K_j\}} = \mathbf{1}_{\{X_t^+ < K_j\}}$, a.s. (In fact, for almost every $\omega \in \{X_t^+ < K_j\}$ (resp. $\{X_t^+ > K_j\}$), there exists an $N(\omega) \in \mathbb{N}$ such that $X_{q_n^+}^+ < (\text{hbox} | \text{resp.} >) K_j$ for any $n \geq N(\omega)$, which means $\lim_{n \rightarrow \infty} \mathbf{1}_{\{X_{q_n^+}^+ < K_j\}}(\omega) = 1(\text{hbox} | \text{resp.} >) 0 = \mathbf{1}_{\{X_t^+ < K_j\}}(\omega)$). Applying the Dominated Convergence Theorem (Theorem 18) twice, we obtain

$$\begin{aligned} \mathcal{E}[\mathbf{1}_A \mathbf{1}_{\{X_t^+ < K_j\}} X_t^+] &= \mathcal{E}[\mathbf{1}_A \mathbf{1}_{\{X_t^+ < K_j\}} (X_t^+ \wedge K_j)] = \\ &\lim_{n \rightarrow \infty} \mathcal{E}[\mathbf{1}_A \mathbf{1}_{\{X_{q_n^+}^+ < K_j\}} (X_{q_n^+}^+ \wedge K_j)] \\ &\geq \lim_{n \rightarrow \infty} \mathcal{E}[\mathbf{1}_A \mathbf{1}_{\{X_{q_n^+}^+ < K_j\}} X_{q_n^+}^+] = \\ &\mathcal{E}[\mathbf{1}_A \mathbf{1}_{\{X_t^+ < K_j\}} X_{q_m^+}^+]. \end{aligned}$$

Since $\lim_{j \rightarrow \infty} \neg \neg \mathbf{1}_{\{X_t^+ < K_j\}} = 1$, a.s., the Dominated Convergence Theorem again implies that

$$\begin{aligned} \mathcal{E}[\mathbf{1}_A X_t^+] &= \lim_{j \rightarrow \infty} \mathcal{E}[\mathbf{1}_A \mathbf{1}_{\{X_t^+ < K_j\}} X_t^+] \geq \lim_{j \rightarrow \infty} \mathcal{E}[\mathbf{1}_A \mathbf{1}_{\{X_t^+ < K_j\}} X_{q_m^+}^+] = \\ &\mathcal{E}[\mathbf{1}_A X_{q_m^+}^+], \end{aligned}$$

which leads to that $\mathcal{E}[\mathbf{1}_A X_t^+] \geq \limsup_{m \rightarrow \infty} \mathcal{E}[\mathbf{1}_A X_{q_m^+}^+]$. Fatou's Lemma (Theorem 16) gives the reverse inequality, thus proving (8.16). Since X is an \mathcal{E} -supermartingale, using (8.16), (A2) and (A3), we obtain

$$\begin{aligned} \mathcal{E}[\mathbf{1}_A X_t^+] &= \lim_{n \rightarrow \infty} \mathcal{E}[\mathbf{1}_A X_{q_n^+}^+] = \lim_{n \rightarrow \infty} \mathcal{E}[\mathcal{E}[\mathbf{1}_A X_{q_n^+}^+ | \mathcal{F}_t]] = \\ &\lim_{n \rightarrow \infty} \mathcal{E}[\mathbf{1}_A \mathcal{E}[X_{q_n^+}^+ | \mathcal{F}_t]] \leq \mathcal{E}[\mathbf{1}_A X_t] \end{aligned}$$

for any $A \in \mathcal{F}_t$, which further implies that $X_t^+ \leq X_t$, a.s. thanks to Proposition 8 (3).

Next, we show that X^+ is an \mathcal{E} -supermartingale: For any $0 \leq s < t \leq T$, it is clear that $q_n^+(s) \leq q_n^+(t)$ for any $n \in \mathbb{N}$. For any $A \in \mathcal{F}_s$, (A3) and Corollary 25 imply that for any $n \in \mathbb{N}$

$$\mathcal{E}[\mathbf{1}_A X_{q_n^+(s)}] \geq \mathcal{E}[\mathbf{1}_A \mathcal{E}[X_{q_n^+(t)} | \mathcal{F}_{q_n^+(s)}]] = \mathcal{E}[\mathcal{E}[\mathbf{1}_A X_{q_n^+(t)} | \mathcal{F}_{q_n^+(s)}]] = \mathcal{E}[\mathbf{1}_A X_{q_n^+(t)}].$$

As $n \rightarrow \infty$, (8.16), (A2) and (A3) imply that

$$\begin{aligned} \mathcal{E}[\mathbf{1}_A X_s^+] &= \lim_{n \rightarrow \infty} \mathcal{E}[\mathbf{1}_A X_{q_n^+(s)}] \geq \lim_{n \rightarrow \infty} \mathcal{E}[\mathbf{1}_A X_{q_n^+(t)}] = \mathcal{E}[\mathbf{1}_A X_t^+] = \\ &= \mathcal{E}[\mathcal{E}[\mathbf{1}_A X_t^+ | \mathcal{F}_s]] = \mathcal{E}[\mathbf{1}_A \mathcal{E}[X_t^+ | \mathcal{F}_s]]. \end{aligned}$$

Then Proposition 8 (3) implies that $X_s^+ \geq \mathcal{E}[X_t^+ | \mathcal{F}_s]$, a.s., thus $\{X_t^+\}_{t \in [0, T]}$ is an RCLL \mathcal{E} -supermartingale.

Proof of (3): If $t \mapsto \mathcal{E}[X_t]$ is right continuous, for any $t \in [0, T]$, (8.16) implies that

$$\mathcal{E}[X_t^+] = \lim_{n \rightarrow \infty} \mathcal{E}[X_{q_n^+(t)}] = \mathcal{E}[X_t].$$

Then the second part of (A1) imply that $X_t^+ = X_t$, a.s., which means that X^+ is an RCLL modification of X . On the other hand, if \tilde{X} is a right-continuous modification of X , we see from (2.4) that except on a null set \tilde{N}

$$\begin{aligned} X_t^+ &= \lim_{n \rightarrow \infty} X_{q_n^+(t)}, \quad \tilde{X}_t = \lim_{n \rightarrow \infty} \tilde{X}_{q_n^+(t)}, \quad \tilde{X}_t = X_t, \quad \langle \text{hbox} \mid a n d \rangle \quad \tilde{X}_{q_n^+(t)} = \\ &X_{q_n^+(t)} \langle \text{hbox} \mid f o r a n y \rangle n \in \mathbb{N}. \end{aligned}$$

Putting them together, it holds on \tilde{N}^c that

$$X_t^+ = \lim_{n \rightarrow \infty} X_{q_n^+(t)} = \lim_{n \rightarrow \infty} \tilde{X}_{q_n^+(t)} = \tilde{X}_t = X_t. \quad (8.18)$$

Since X is an \mathcal{E} -supermartingale, (A2) implies that for any $0 \leq t_1 < t_2 \leq T$, $\mathcal{E}[X_{t_1}] \geq \mathcal{E}[\mathcal{E}[X_{t_2} | \mathcal{F}_{t_1}]] = \mathcal{E}[X_{t_2}]$, which shows that the function $t \mapsto \mathcal{E}[X_t]$ is decreasing. Then (8.16) and (8.18) imply that for any $t \in [0, T]$

$$\mathcal{E}[X_t] \geq \lim_{s \downarrow t} \mathcal{E}[X_s] = \lim_{n \rightarrow \infty} \mathcal{E}[X_{q_n^+(t)}] = \mathcal{E}[X_t^+] = \mathcal{E}[X_t],$$

thus $\lim_{s \downarrow t} \mathcal{E}[X_s] = \mathcal{E}[X_t]$, i.e., the function $t \mapsto \mathcal{E}[X_t]$ is right continuous. \square

Proof of Corollary 25 : Since $\text{essinf}_{t \in [0, T]} X_t \geq c$, a.s., we can deduce from (A4) that $X^c \triangleq \{X_t - c\}_{t \in [0, T]}$ is a non-negative \mathcal{E} -supermartingale. If $\text{esssup}_{t \in \mathcal{D}_T} X_t \in D o m^\#(\mathcal{E})$ (D2) implies that $\text{esssup}_{t \in \mathcal{D}_T} X_t \in D o m^\#(\mathcal{E})$ is equivalent to $\text{esssup}_{t \in \mathcal{D}_T} X_t^c \in D o m^+(\mathcal{E})$ or if (8.6) holds, Proposition 82 (1) shows that for any $\nu \in \mathcal{S}_{0, T}$, $(X^c)_\nu^+$ belongs to $D o m^+(\mathcal{E})$. Because

$$(X^c)_t^+ = X_t^+ - c, \quad \forall t \in [0, T], \quad (8.19)$$

(D2) and the non-negativity of $(X^c)^+$ imply that

$$X_\nu^+ = (X^c)_\nu^+ + c \in Dom^\#(\mathcal{E}).$$

On the other hand, if $X_t^+ \in Dom^\#(\mathcal{E})$ for any $t \in [0, T]$, (D2) implies that the non-negative random variable $(X^c)_t^+ = X_t^+ - c$ belongs to $Dom^+(\mathcal{E})$. Hence, Proposition 82 (2) show that $(X^c)^+$ is an RCLL \mathcal{E} -supermartingale such that for any $t \in [0, T]$, $(X^c)_t^+ \leq X_t^c$, a.s. Then (8.19), (2.8) and (A4) imply that X^+ is an RCLL $\tilde{\mathcal{E}}$ -supermartingale such that for any $t \in [0, T]$, $X_t^+ \leq X_t$, a.s. Moreover, if $t \mapsto \tilde{\mathcal{E}}[X_t]$ is a right-continuous function (which is equivalent to the right continuity of $t \mapsto \mathcal{E}[X_t^c]$), then we know from Proposition 82 (2) that for any $t \in [0, T]$, $(X^c)_t^+ = X_t^c$, a.s., or equivalently, $X_t^+ = X_t$, a.s. Conversely, if X has a right-continuous modification, so does X^c , then Proposition 82 (2) once again shows that $t \mapsto \mathcal{E}[X_t^c]$ is right continuous, which is equivalent to the right continuity of $t \mapsto \tilde{\mathcal{E}}[X_t]$. This completes the proof. \square

Proof of Theorem 24: We shall only consider the $\tilde{\mathcal{E}}$ -supermartingale case, as the other cases can be deduced easily by similar arguments. Fix $t \in [0, T]$, we let $\{\nu_n^t\}_{n \in \mathbb{N}}$ be a decreasing sequence in $\mathcal{S}_{t,T}^F$ such that $\lim_{n \rightarrow \infty} \nu_n^t = \nu \vee t$. Since $\text{essinf}_{t \in \mathcal{D}_T} X_t \geq c$, a.s., it holds a.s. that $X_t \geq c$ for each $t \in \mathcal{D}_T$. The right-continuity of the process X then implies that except on a null set N , $X_t \geq c$ for any $t \in [0, T]$. Thus we see from (A4) that $X^c \stackrel{\Delta}{=} \{X_t - c\}_{t \in [0, T]}$ is a non-negative \mathcal{E} -supermartingale. For any $n \in \mathbb{N}$ and $A \in \mathcal{F}_t \subset \mathcal{F}_{\nu \vee t}$, (A2), (A3) and Proposition 24 imply that

$$\mathcal{E}[\mathbf{1}_A X_{\nu_n^t}^c] = \mathcal{E}[\mathcal{E}[\mathbf{1}_A X_{\nu_n^t}^c | \mathcal{F}_t]] = \mathcal{E}[\mathbf{1}_A \mathcal{E}[X_{\nu_n^t}^c | \mathcal{F}_t]] \leq \mathcal{E}[\mathbf{1}_A X_t^c]. \quad (8.20)$$

We also have that $\mathcal{E}[\mathbf{1}_A X_{\nu \vee t}^c] = \lim_{n \rightarrow \infty} \mathcal{E}[\mathbf{1}_A X_{\nu_n^t}^c]$. The proof is similar to that of (8.16). (We only need to replace X_t^+ by $X_{\nu \vee t}^c$ and $X_{q_n^+(t)}$ by $X_{\nu_n^t}^c$ in the proof of (8.16)). As $n \rightarrow \infty$ in (8.20), (A2) and (A3) imply that

$$\mathcal{E}[\mathbf{1}_A X_t^c] \geq \lim_{n \rightarrow \infty} \mathcal{E}[\mathbf{1}_A X_{\nu_n^t}^c] = \mathcal{E}[\mathbf{1}_A X_{\nu \vee t}^c] = \mathcal{E}[\mathcal{E}[\mathbf{1}_A X_{\nu \vee t}^c | \mathcal{F}_t]] = \mathcal{E}[\mathbf{1}_A \mathcal{E}[X_{\nu \vee t}^c | \mathcal{F}_t]].$$

Applying Proposition 8 (3), we obtain that $\mathcal{E}[X_{\nu \vee t}^c | \mathcal{F}_t] \leq X_t^c$, a.s. Then (A4) and (2.8) imply that

$$\tilde{\mathcal{E}}[X_{\nu \vee t} | \mathcal{F}_t] = \mathcal{E}[X_{\nu \vee t} | \mathcal{F}_t] = \mathcal{E}[X_{\nu \vee t}^c + c | \mathcal{F}_t] = \mathcal{E}[X_{\nu \vee t}^c | \mathcal{F}_t] + c \leq X_t^c + c = X_t, \quad a. s.$$

Since $\{\nu \leq t\} \in \mathcal{F}_t$, we can deduce from (A3) and (A4) that

$$\begin{aligned} \tilde{\mathcal{E}}[X_\nu | \mathcal{F}_t] &= \tilde{\mathcal{E}}[\mathbf{1}_{\{\nu > t\}} X_{\nu \vee t} + \mathbf{1}_{\{\nu \leq t\}} X_{\nu \wedge t} | \mathcal{F}_t] = \mathbf{1}_{\{\nu > t\}} \tilde{\mathcal{E}}[X_{\nu \vee t} | \mathcal{F}_t] + \\ &\quad \mathbf{1}_{\{\nu \leq t\}} X_{\nu \wedge t} \\ &\leq \mathbf{1}_{\{\nu > t\}} X_t + \mathbf{1}_{\{\nu \leq t\}} X_{\nu \wedge t} = X_{\nu \wedge t} \quad a. s. \end{aligned}$$

Hence, we can find a null set \tilde{N} such that except on \tilde{N}^c

$$\tilde{\mathcal{E}}[X_\nu|\mathcal{F}_t] \leq X_{\nu \wedge t}, \quad \langle \text{hbox} \quad | \quad \text{f o r a n y } t \in \mathcal{D}_T \text{ and the paths of } \tilde{\mathcal{E}}[X_\nu|\mathcal{F}_\cdot] \text{ and } X_{\nu \wedge \cdot} \text{ are all RCLL.} \rangle$$

As a result, on \tilde{N}^c

$$\begin{aligned} \tilde{\mathcal{E}}[X_\nu|\mathcal{F}_t] &\leq X_{\nu \wedge t}, \quad \forall t \in [0, T], \quad \langle \text{hbox} | \text{t h u s} \rangle \quad \tilde{\mathcal{E}}[X_\nu|\mathcal{F}_\sigma] \leq \\ X_{\nu \wedge \sigma}, \quad \forall \sigma \in \mathcal{S}_{0,T}. \quad \langle \text{hbox} | \quad \square \rangle \end{aligned}$$

Proof of Proposition 26: 1) If $\xi \leq \eta$, a.s., by (A1), it holds except on a null set N that

$$\tilde{\mathcal{E}}[\xi|\mathcal{F}_t] \leq \tilde{\mathcal{E}}[\eta|\mathcal{F}_t], \quad \langle \text{hbox} \quad | \quad \text{f o r a n y } t \in \mathcal{D}_T \text{ and that the paths of } \tilde{\mathcal{E}}[\xi|\mathcal{F}_\cdot] \text{ and } \tilde{\mathcal{E}}[\eta|\mathcal{F}_\cdot] \text{ are all RCLL,} \rangle$$

which implies that on N^c

$$\tilde{\mathcal{E}}[\xi|\mathcal{F}_t] \leq \tilde{\mathcal{E}}[\eta|\mathcal{F}_t], \quad \forall t \in [0, T], \quad \langle \text{hbox} | \text{t h u s} \rangle \quad \tilde{\mathcal{E}}[\xi|\mathcal{F}_\nu] \leq \tilde{\mathcal{E}}[\eta|\mathcal{F}_\nu].$$

Moreover, if $\tilde{\mathcal{E}}[\xi|\mathcal{F}_\sigma] = \tilde{\mathcal{E}}[\eta|\mathcal{F}_\sigma]$, a.s. for some $\sigma \in \mathcal{S}_{0,T}$, we can apply (2.8) and Corollary 25 to get

$$\mathcal{E}[\xi] = \tilde{\mathcal{E}}[\xi] = \tilde{\mathcal{E}}[\tilde{\mathcal{E}}[\xi|\mathcal{F}_\sigma]] = \tilde{\mathcal{E}}[\tilde{\mathcal{E}}[\eta|\mathcal{F}_\sigma]] = \tilde{\mathcal{E}}[\eta] = \mathcal{E}[\eta].$$

Then (A4) implies that $\mathcal{E}[\xi - c(\xi)] = \mathcal{E}[\xi] - c(\xi) = \mathcal{E}[\eta] - c(\xi) = \mathcal{E}[\eta - c(\xi)]$. Clearly, $0 \leq \xi - c(\xi) \leq \eta - c(\xi)$, a.s. The second part of (A1) then implies that $\xi - c(\xi) = \eta - c(\xi)$, a.s., i.e. $\xi = \eta$, a.s., proving (1).

2) For any $A \in \mathcal{F}_\nu$ and $\eta \in D o m_\nu^\#(\mathcal{E})$, we let $\{\nu_n\}_{n \in \mathbb{N}}$ be a decreasing sequence in $\mathcal{S}_{0,T}^F$ such that $\lim_{n \rightarrow \infty} \neg \downarrow \nu_n = \nu$, a.s. For any $n \in \mathbb{N}$, since $A \in \mathcal{F}_{\nu_n}$ and $\eta \in D o m_{\nu_n}^\#(\mathcal{E})$, Proposition 26 (2) and (3) imply that

$$\begin{aligned} \tilde{\mathcal{E}}[\mathbf{1}_A \xi|\mathcal{F}_{\nu_n}] &= \mathbf{1}_A \tilde{\mathcal{E}}[\xi|\mathcal{F}_{\nu_n}], \quad \langle \text{hbox} \quad | \quad \text{a n d} \rangle \quad \tilde{\mathcal{E}}[\xi + \eta|\mathcal{F}_{\nu_n}] = \tilde{\mathcal{E}}[\xi|\mathcal{F}_{\nu_n}] + \eta, \\ \text{a. s.} \end{aligned} \quad (8.21)$$

Then we can find a null set N' such that except on N'

$$\langle \text{error} | \text{compound hbox} \rangle$$

As $n \rightarrow \infty$, it holds on $(N')^c$ that

$$\begin{aligned} \tilde{\mathcal{E}}[\mathbf{1}_A \xi|\mathcal{F}_\nu] &= \lim_{n \rightarrow \infty} \tilde{\mathcal{E}}[\mathbf{1}_A \xi|\mathcal{F}_{\nu_n}] = \\ \lim_{n \rightarrow \infty} \mathbf{1}_A \tilde{\mathcal{E}}[\xi|\mathcal{F}_{\nu_n}] &= \mathbf{1}_A \tilde{\mathcal{E}}[\xi|\mathcal{F}_\nu], \\ \langle \text{error} | \text{compound hbox} \rangle \quad \tilde{\mathcal{E}}[\xi + \eta|\mathcal{F}_\nu] &= \lim_{n \rightarrow \infty} \tilde{\mathcal{E}}[\xi + \eta|\mathcal{F}_{\nu_n}] = \lim_{n \rightarrow \infty} \tilde{\mathcal{E}}[\xi|\mathcal{F}_{\nu_n}] + \\ \eta &= \tilde{\mathcal{E}}[\xi|\mathcal{F}_\nu] + \eta, \end{aligned}$$

proving (2) and (3). Proofs of (4) and (5) are similar to those of Proposition 8 (1) and (2). The proofs can be carried out by applying the just obtained "Zero-one Law" and "Translation Invariance". \square

8.2 Proofs of Section 7

Proof of Lemma 80: (1) Let \mathcal{E} be a positively-convex F -expectation. For any $A \in \mathcal{F}_T$ and $n \in \mathbb{N}$, (D1) and (D2) imply that $\mathbf{1}_A, n\mathbf{1}_A \in \text{Dom}(\mathcal{E})$. Then the positive-convexity of \mathcal{E} and Proposition 8 (2) show that

$$\begin{aligned} \mathcal{E}[\mathbf{1}_A] &= \mathcal{E}\left[\frac{1}{n}(n\mathbf{1}_A)\right] \leq \frac{1}{n}\mathcal{E}[n\mathbf{1}_A] + \left(1 - \frac{1}{n}\right)\mathcal{E}[0] = \frac{1}{n}\mathcal{E}[n\mathbf{1}_A] + \left(1 - \frac{1}{n}\right) \cdot 0 = \\ &= \frac{1}{n}\mathcal{E}[n\mathbf{1}_A]. \end{aligned} \quad (8.22)$$

Since $P(A) > 0$, one can deduce from the second part of (A1) that $\mathcal{E}[\mathbf{1}_A] > 0$. Letting $n \rightarrow \infty$ in (8.22) yields that

$$\lim_{n \rightarrow \infty} \mathcal{E}[n\mathbf{1}_A] \geq \lim_{n \rightarrow \infty} n\mathcal{E}[\mathbf{1}_A] = \infty,$$

thus \mathcal{E} satisfies (H0). Moreover, for any $\xi, \eta \in \text{Dom}^\#(\mathcal{E})$, $\lambda \in (0, 1)$ and $t \in [0, T]$, we can deduce from (2.8), (A4) and the positive-convexity of \mathcal{E} that

$$\begin{aligned} \tilde{\mathcal{E}}[\lambda\xi + (1-\lambda)\eta | \mathcal{F}_t] &= \mathcal{E}[\lambda\xi + (1-\lambda)\eta | \mathcal{F}_t] = \mathcal{E}[\lambda(\xi - c(\xi)) + (1-\lambda)(\eta - c(\eta)) | \mathcal{F}_t] + \lambda c(\xi) + (1-\lambda)c(\eta) \\ &\leq \lambda\mathcal{E}[\xi - c(\xi) | \mathcal{F}_t] + \lambda c(\xi) + (1-\lambda)\mathcal{E}[\eta - c(\eta) | \mathcal{F}_t] + (1-\lambda)c(\eta) \\ &= \lambda\mathcal{E}[\xi | \mathcal{F}_t] + (1-\lambda)\mathcal{E}[\eta | \mathcal{F}_t] = \lambda\tilde{\mathcal{E}}[\xi | \mathcal{F}_t] + (1-\lambda)\tilde{\mathcal{E}}[\eta | \mathcal{F}_t], \quad a.s., \end{aligned}$$

which shows that $\tilde{\mathcal{E}}$ is convex in the sense of (?). On the other hand, if $\tilde{\mathcal{E}}$ satisfies (?), since $\text{Dom}^+(\mathcal{E}) \subset \text{Dom}^\#(\mathcal{E})$, one can easily deduce from (2.8) that \mathcal{E} is positively-convex. \square

Proof of Proposition 75: We first check that $\mathcal{E}_{i,j}^\nu$ satisfies (A1)-(A4). For this purpose, let $\xi, \eta \in \Lambda^\#$ and $t \in [0, T]$.

1) If $\xi \leq \eta$, a.s., applying Proposition 26 (1) to $\tilde{\mathcal{E}}$ yields that $\tilde{\mathcal{E}}_j[\xi | \mathcal{F}_{\nu \vee t}] \leq \tilde{\mathcal{E}}_j[\eta | \mathcal{F}_{\nu \vee t}]$, a.s. Then (A1) of $\tilde{\mathcal{E}}_i$ and (75) imply that

$$\mathcal{E}_{i,j}^\nu[\xi | \mathcal{F}_t] = \tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_j[\xi | \mathcal{F}_{\nu \vee t}] | \mathcal{F}_t] \leq \tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_j[\eta | \mathcal{F}_{\nu \vee t}] | \mathcal{F}_t] = \mathcal{E}_{i,j}^\nu[\eta | \mathcal{F}_t], \quad a.s.$$

Moreover, if $0 \leq \xi \leq \eta$ a.s. and $\mathcal{E}_{i,j}^\nu[\xi] = \mathcal{E}_{i,j}^\nu[\eta]$ (i.e. $\tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_j[\xi | \mathcal{F}_\nu]] = \tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_j[\eta | \mathcal{F}_\nu]]$ by (75)), the second part of (A1) implies that $\tilde{\mathcal{E}}_j[\xi | \mathcal{F}_\nu] = \tilde{\mathcal{E}}_j[\eta | \mathcal{F}_\nu]$, a.s. Further applying the second part of Proposition 26 (1), we obtain $\xi = \eta$, a.s., proving (A1) for $\mathcal{E}_{i,j}^\nu$.

2) Next, we let $0 \leq s \leq t \leq T$ and set $\Xi_t \stackrel{\Delta}{=} \mathcal{E}_{i,j}^\nu[\xi | \mathcal{F}_t]$. Applying Proposition 26 (2) to $\tilde{\mathcal{E}}_i$ and $\tilde{\mathcal{E}}_j$, we obtain

$$\begin{aligned} \mathcal{E}_{i,j}^\nu[\Xi_t | \mathcal{F}_s] &\neg \neg \mathbf{1}_{\{\nu \leq s\}} \tilde{\mathcal{E}}_j[\Xi_t | \mathcal{F}_s] \neg \neg \mathbf{1}_{\{\nu > s\}} \tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_j[\Xi_t | \mathcal{F}_\nu] | \mathcal{F}_s] \neg \neg \\ &\tilde{\mathcal{E}}_j[\mathbf{1}_{\{\nu \leq s\}} \Xi_t | \mathcal{F}_s] \neg \neg \neg \tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_j[\mathbf{1}_{\{\nu > s\}} \Xi_t | \mathcal{F}_\nu] | \mathcal{F}_s], \quad a.s., \end{aligned}$$

where we used the fact that $\{\nu > s\} \in \mathcal{F}_{\nu \wedge s}$ thanks to (, ?, Lemma 1.2.16). Then (A3) and (A2) imply that

$$\begin{aligned} \tilde{\mathcal{E}}_j[11_{\{\nu \leq s\}} \Xi_t | \mathcal{F}_s] &= \tilde{\mathcal{E}}_j[11_{\{\nu \leq s\}} \tilde{\mathcal{E}}_j[\xi | \mathcal{F}_t] | \mathcal{F}_s] = 11_{\{\nu \leq s\}} \tilde{\mathcal{E}}_j[\tilde{\mathcal{E}}_j[\xi | \mathcal{F}_t] | \mathcal{F}_s] = \\ &= 11_{\{\nu \leq s\}} \tilde{\mathcal{E}}_j[\xi | \mathcal{F}_s], \quad a.s. \end{aligned} \quad (8.23)$$

On the other hand, we can deduce from (75) that

$$\begin{aligned} 11_{\{\nu > s\}} \Xi_t &= 11_{\{s < \nu \leq t\}} \tilde{\mathcal{E}}_j[\xi | \mathcal{F}_t] + 11_{\{\nu > t\}} \tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_j[\xi | \mathcal{F}_\nu] | \mathcal{F}_t] = \\ &= 11_{\{s < \nu \leq t\}} \tilde{\mathcal{E}}_j[\xi | \mathcal{F}_t] + 11_{\{\nu > t\}} \tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_j[\xi | \mathcal{F}_\nu] | \mathcal{F}_{\nu \wedge t}], \quad a.s. \end{aligned}$$

Since both $\{s < \nu \leq t\} = \{\nu > s\} \cap \{\nu > t\}^c$ and $\{\nu > t\}$ belong to $\mathcal{F}_{\nu \wedge t}$, Proposition 26 (3) and (2) as well as Corollary 25 imply that

$$\begin{aligned} \tilde{\mathcal{E}}_j[11_{\{\nu > s\}} \Xi_t | \mathcal{F}_\nu] &= \tilde{\mathcal{E}}_j[11_{\{s < \nu \leq t\}} \tilde{\mathcal{E}}_j[\xi | \mathcal{F}_t] | \mathcal{F}_\nu] + 11_{\{\nu > t\}} \tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_j[\xi | \mathcal{F}_\nu] | \mathcal{F}_{\nu \wedge t}] \\ &= 11_{\{s < \nu \leq t\}} \tilde{\mathcal{E}}_j[\tilde{\mathcal{E}}_j[\xi | \mathcal{F}_t] | \mathcal{F}_\nu] + 11_{\{\nu > t\}} \tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_j[\xi | \mathcal{F}_\nu] | \mathcal{F}_t] = \\ &= 11_{\{s < \nu \leq t\}} \tilde{\mathcal{E}}_j[\xi | \mathcal{F}_{\nu \wedge t}] + \tilde{\mathcal{E}}_i[11_{\{\nu > t\}} \tilde{\mathcal{E}}_j[\xi | \mathcal{F}_\nu] | \mathcal{F}_t] \\ &= \tilde{\mathcal{E}}_i[11_{\{s < \nu \leq t\}} \tilde{\mathcal{E}}_j[\xi | \mathcal{F}_{\nu \wedge t}] + 11_{\{\nu > t\}} \tilde{\mathcal{E}}_j[\xi | \mathcal{F}_\nu] | \mathcal{F}_t] = \\ &= \tilde{\mathcal{E}}_i[11_{\{s < \nu\}} \tilde{\mathcal{E}}_j[\xi | \mathcal{F}_\nu] | \mathcal{F}_t], \quad a.s. \end{aligned}$$

Taking $\tilde{\mathcal{E}}_i[\cdot | \mathcal{F}_s]$ of both sides as well as using (A2) and (A3) of $\tilde{\mathcal{E}}_i$, we obtain

$$\begin{aligned} \tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_j[11_{\{\nu > s\}} \Xi_t | \mathcal{F}_\nu] | \mathcal{F}_s] &= \neg \neg \tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_i[11_{\{s < \nu\}} \tilde{\mathcal{E}}_j[\xi | \mathcal{F}_\nu] | \mathcal{F}_t] | \mathcal{F}_s] = \neg \neg \\ &= \tilde{\mathcal{E}}_i[11_{\{s < \nu\}} \tilde{\mathcal{E}}_j[\xi | \mathcal{F}_\nu] | \mathcal{F}_s] = \neg \neg 11_{\{\nu > s\}} \tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_j[\xi | \mathcal{F}_\nu] | \mathcal{F}_s], \quad a.s., \end{aligned}$$

which together with (8.23) yields that

$$\begin{aligned} \mathcal{E}_{i,j}^\nu[\mathcal{E}_{i,j}^\nu[\xi | \mathcal{F}_t] | \mathcal{F}_s] &= 11_{\{\nu \leq s\}} \tilde{\mathcal{E}}_j[\xi | \mathcal{F}_s] + 11_{\{\nu > s\}} \tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_j[\xi | \mathcal{F}_\nu] | \mathcal{F}_s] = \mathcal{E}_{i,j}^\nu[\xi | \mathcal{F}_s], \\ &a.s., \quad \langle \text{hbox} | \text{proving (A2) for} \rangle \mathcal{E}_{i,j}^\nu. \end{aligned}$$

3) For any $A \in \mathcal{F}_t$, using (75), (A3) of $\tilde{\mathcal{E}}_i$ as well as applying Proposition 26 (2) to $\tilde{\mathcal{E}}_j$, we obtain

$$\begin{aligned} \mathcal{E}_{i,j}^\nu[11_A \xi | \mathcal{F}_t] &= \tilde{\mathcal{E}}_i[11_A \tilde{\mathcal{E}}_j[\xi | \mathcal{F}_{\nu \vee t}] | \mathcal{F}_t] = 11_A \tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_j[\xi | \mathcal{F}_{\nu \vee t}] | \mathcal{F}_t] = 11_A \mathcal{E}_{i,j}^\nu[\xi | \mathcal{F}_t], \\ &a.s., \quad \langle \text{hbox} | \text{proving (A3) for} \rangle \mathcal{E}_{i,j}^\nu. \end{aligned}$$

Similarly, we can show that (A4) holds for $\mathcal{E}_{i,j}^\nu$ as well. Therefore, $\mathcal{E}_{i,j}^\nu$ is an \mathbf{F} -expectation with domain $\Lambda^\#$. Since $\Lambda \in \tilde{\mathcal{D}}_T$, i.e. $\mathbb{R} \subset \Lambda$, it follows easily that $\mathbb{R} \subset \Lambda^\#$, which shows that $\Lambda^\# \in \tilde{\mathcal{D}}_T$.

4) Now we show that $\mathcal{E}_{i,j}^\nu$ satisfies (H1) and (H2): For any $\xi \in \Lambda^+$ and any $\{A_n\}_{n \in \mathbb{N}} \subset \mathcal{F}_T$ with $\lim_{n \rightarrow \infty} \neg \uparrow 11_{A_n} = 1$, a.s., the Dominated Convergence Theorem (Proposition 29) implies that $\lim_{n \rightarrow \infty} \neg \neg \uparrow \tilde{\mathcal{E}}_j[11_{A_n} \xi | \mathcal{F}_\nu] = \tilde{\mathcal{E}}_j[\xi | \mathcal{F}_\nu]$, a.s. Furthermore, using (75) and applying the Dominated Convergence Theorem to $\tilde{\mathcal{E}}_i$ yield that

$$\begin{aligned} \lim_{n \rightarrow \infty} \neg \neg \uparrow \mathcal{E}_{i,j}^\nu[11_{A_n} \xi] &= \lim_{n \rightarrow \infty} \neg \neg \uparrow \tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_j[11_{A_n} \xi | \mathcal{F}_\nu]] = \tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_j[\xi | \mathcal{F}_\nu]] = \mathcal{E}_{i,j}^\nu[\xi], \\ &\langle \text{hbox} | \text{proving (H1) for} \rangle \mathcal{E}_{i,j}^\nu. \end{aligned}$$

With a similar argument, we can show that $\mathcal{E}_{i,j}^\nu$ also satisfies (H2).

5) If both \mathcal{E}_i and \mathcal{E}_j are positively-convex, so are $\tilde{\mathcal{E}}_i$ and $\tilde{\mathcal{E}}_j$ thanks to (2.8). To see that $\mathcal{E}_{i,j}^\nu$ is convex in the sense of (?), we fix $\xi, \eta \in \Lambda^\#$, $\lambda \in (0, 1)$ and $t \in [0, T]$. For any $s \in [0, T]$, we have

$$\tilde{\mathcal{E}}_j[\lambda\xi + (1-\lambda)\eta|\mathcal{F}_s] \leq \lambda\tilde{\mathcal{E}}_j[\xi|\mathcal{F}_s] + (1-\lambda)\tilde{\mathcal{E}}_j[\eta|\mathcal{F}_s], \quad a.s.$$

Since $\tilde{\mathcal{E}}_j[\lambda\xi + (1-\lambda)\eta|\mathcal{F}_\cdot]$, $\tilde{\mathcal{E}}_j[\xi|\mathcal{F}_\cdot]$ and $\tilde{\mathcal{E}}_j[\eta|\mathcal{F}_\cdot]$ are all RCLL processes, it holds except on a null set N that

$$\tilde{\mathcal{E}}_j[\lambda\xi + (1-\lambda)\eta|\mathcal{F}_s] \leq \lambda\tilde{\mathcal{E}}_j[\xi|\mathcal{F}_s] + (1-\lambda)\tilde{\mathcal{E}}_j[\eta|\mathcal{F}_s], \quad \forall s \in [0, T],$$

$$\langle \text{hbox} \mid t \text{ h u s} \rangle \quad \tilde{\mathcal{E}}_j[\lambda\xi + (1-\lambda)\eta|\mathcal{F}_{\nu \vee t}] \leq \lambda\tilde{\mathcal{E}}_j[\xi|\mathcal{F}_{\nu \vee t}] + (1-\lambda)\tilde{\mathcal{E}}_j[\eta|\mathcal{F}_{\nu \vee t}].$$

Then (75) implies that

$$\begin{aligned} \mathcal{E}_{i,j}^\nu[\lambda\xi + (1-\lambda)\eta|\mathcal{F}_t] &= \tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_j[\lambda\xi + (1-\lambda)\eta|\mathcal{F}_{\nu \vee t}]|\mathcal{F}_t] \leq \tilde{\mathcal{E}}_i[\lambda\tilde{\mathcal{E}}_j[\xi|\mathcal{F}_{\nu \vee t}] + \\ &\quad (1-\lambda)\tilde{\mathcal{E}}_j[\eta|\mathcal{F}_{\nu \vee t}]|\mathcal{F}_t], \\ &\leq \lambda\tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_j[\xi|\mathcal{F}_{\nu \vee t}]|\mathcal{F}_t] + (1-\lambda)\tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_j[\eta|\mathcal{F}_{\nu \vee t}]|\mathcal{F}_t] = \lambda\mathcal{E}_{i,j}^\nu[\xi|\mathcal{F}_t] + \\ &\quad (1-\lambda)\mathcal{E}_{i,j}^\nu[\eta|\mathcal{F}_t], \quad a.s. \end{aligned} \quad \langle \text{hbox} \mid \square \rangle$$

8.3 Proofs of Section 4

Proof of Lemma 80: For any $i \in \mathcal{I}$, it is clear that $H_0^i = 0$ and that (4.2) directly follows from (h1). For any $s, t \in \mathcal{D}_T$ with $s < t$, we can deduce from (h2) that

$$H_{s,t}^i = \int_s^t h_r^i dr \geq c \int_s^t ds \geq cT, \quad a.s., \quad (8.24)$$

which implies that $\operatorname{ess\,inf}_{s,t \in \mathcal{D}_T; s < t} H_{s,t}^i \geq cT$, a.s. Thus (S2) holds with $C_H = cT$.

If no member of \mathcal{E} satisfies (8.6), then $\int_0^T |h_t^j| dt \in \operatorname{Dom}(\mathcal{E})$ for some $j \in \mathcal{I}$ is assumed. For any $s, t \in \mathcal{D}_T$ with $s < t$, we can deduce from (8.24) and (h2) that

$$C_H \leq H_{s,t}^j \leq \int_s^t |h_r^j| dr \leq \int_0^T |h_r^j| dr, \quad a.s.,$$

which implies that $C_H \leq \operatorname{ess\,sup}_{s,t \in \mathcal{D}_T; s < t} H_{s,t}^j \leq \int_0^T |h_r^j| dr$ a.s. Then Lemma 80

shows that $\operatorname{ess\,sup}_{s,t \in \mathcal{D}_T; s < t} H_{s,t}^j \in \operatorname{Dom}(\mathcal{E})$, i.e. (?). Moreover, we can derive (S3) directly from (h3). \square

Proof of Lemma 80: For any $i, j \in \mathcal{I}'$ and $\rho_1, \rho_2 \in \mathcal{U}$, we consider the event

$$A \triangleq \{\tilde{\mathcal{E}}_i[X(\rho_1) + H_{\nu, \rho_1}^i|\mathcal{F}_\nu] \leq \tilde{\mathcal{E}}_j[X(\rho_2) + H_{\nu, \rho_2}^j|\mathcal{F}_\nu]\} \in \mathcal{F}_\nu,$$

and define stopping times $\rho \stackrel{\Delta}{=} \rho_2 \mathbf{1}_A + \rho_1 \mathbf{1}_{A^c} \in \mathcal{U}$ and $\nu(A) \stackrel{\Delta}{=} \nu \mathbf{1}_A + T \mathbf{1}_{A^c} \in \mathcal{S}_{\nu, T}$. Since $\mathcal{E}' = \{\mathcal{E}_i\}_{i \in \mathcal{I}'}$ is a stable subclass of \mathcal{E} , Definition 7.13 assures the existence of $k = k(i, j, \nu(A)) \in \mathcal{I}'$ such that $\tilde{\mathcal{E}}_k = \mathcal{E}_{i, j}^{\nu(\neg A)}$. Applying Proposition 26 (5) to $\tilde{\mathcal{E}}_j$ and Proposition 26 (3) & (2) to $\tilde{\mathcal{E}}_i$, we can deduce from (75) that for any $\xi \in \text{Dom}(\mathcal{E})$

$$\begin{aligned} \tilde{\mathcal{E}}_k[\xi | \mathcal{F}_\nu] &= \mathcal{E}_{i, j}^{\nu(\neg A)}[\xi | \mathcal{F}_\nu] = \tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_j[\xi | \mathcal{F}_{\nu(A) \vee \nu}] | \mathcal{F}_\nu] = \tilde{\mathcal{E}}_i[\mathbf{1}_A \tilde{\mathcal{E}}_j[\xi | \mathcal{F}_\nu] + \\ &\quad \mathbf{1}_{A^c} \tilde{\mathcal{E}}_j[\xi | \mathcal{F}_T] | \mathcal{F}_\nu] \\ &= \tilde{\mathcal{E}}_i[\mathbf{1}_A \tilde{\mathcal{E}}_j[\xi | \mathcal{F}_\nu] + \mathbf{1}_{A^c} \xi | \mathcal{F}_\nu] = \mathbf{1}_A \tilde{\mathcal{E}}_j[\xi | \mathcal{F}_\nu] + \mathbf{1}_{A^c} \tilde{\mathcal{E}}_i[\xi | \mathcal{F}_\nu], \\ &\quad \text{a. s.} \end{aligned} \tag{8.25}$$

Moreover, (4.5) implies that

$$\begin{aligned} H_{\nu, \rho}^k \neg &= \neg H_{\nu(A) \wedge \nu, \nu(A) \wedge \rho}^i \neg + \neg H_{\nu(A) \vee \nu, \nu(A) \vee \rho}^j \neg = \neg \mathbf{1}_{A^c} H_{\nu, \rho_1}^i \neg + \neg \mathbf{1}_A H_{\nu, \rho_2}^j, \\ &\quad \text{a. s.} \end{aligned}$$

Then applying Proposition 26 (2) to $\tilde{\mathcal{E}}_i$ and $\tilde{\mathcal{E}}_j$, we see from (8.25) that

$$\begin{aligned} \tilde{\mathcal{E}}_k[X(\rho) + H_{\nu, \rho}^k | \mathcal{F}_\nu] &= \mathbf{1}_A \tilde{\mathcal{E}}_j[X(\rho) + H_{\nu, \rho}^k | \mathcal{F}_\nu] + \mathbf{1}_{A^c} \tilde{\mathcal{E}}_i[X(\rho) + H_{\nu, \rho}^k | \mathcal{F}_\nu] \\ &= \tilde{\mathcal{E}}_j[\mathbf{1}_A X(\rho_2) + \mathbf{1}_A H_{\nu, \rho_2}^j | \mathcal{F}_\nu] + \tilde{\mathcal{E}}_i[\mathbf{1}_{A^c} X(\rho_1) + \\ &\quad \mathbf{1}_{A^c} H_{\nu, \rho_1}^i | \mathcal{F}_\nu] \\ &= \mathbf{1}_A \tilde{\mathcal{E}}_j[X(\rho_2) + H_{\nu, \rho_2}^j | \mathcal{F}_\nu] + \mathbf{1}_{A^c} \tilde{\mathcal{E}}_i[X(\rho_1) + H_{\nu, \rho_1}^i | \mathcal{F}_\nu] \\ &= \tilde{\mathcal{E}}_i[X(\rho_1) + H_{\nu, \rho_1}^i | \mathcal{F}_\nu] \vee \tilde{\mathcal{E}}_j[X(\rho_2) + H_{\nu, \rho_2}^j | \mathcal{F}_\nu], \quad \text{a. s.} \end{aligned}$$

Similarly, taking $\rho' \stackrel{\Delta}{=} \rho_1 \mathbf{1}_A + \rho_2 \mathbf{1}_{A^c}$ and $k' = k(i, j, \nu(A^c))$, we obtain

$$\tilde{\mathcal{E}}_{k'}[X(\rho') + H_{\nu, \rho'}^{k'} | \mathcal{F}_\nu] = \tilde{\mathcal{E}}_i[X(\rho_1) + H_{\nu, \rho_1}^i | \mathcal{F}_\nu] \wedge \tilde{\mathcal{E}}_j[X(\rho_2) + H_{\nu, \rho_2}^j | \mathcal{F}_\nu], \quad \text{a. s.}$$

Hence, the family $\{\tilde{\mathcal{E}}_i[X(\rho) + H_{\nu, \rho}^i | \mathcal{F}_\nu]\}_{(i, \rho) \in \mathcal{I}' \times \mathcal{U}}$ is closed under pairwise maximization and pairwise minimization. Thanks to (4.9), Proposition VI-11-1), we can find two sequences $\{(i_n, \rho_n)\}_{n \in \mathbb{N}}$ and $\{(i'_n, \rho'_n)\}_{n \in \mathbb{N}}$ in $\mathcal{I}' \times \mathcal{U}$ such that (4.9) and (4.10) hold. \square

Proof of Lemma 46: We fix $\nu \in \mathcal{S}_{0, T}$. For any $(i, \rho) \in \mathcal{I} \times \mathcal{S}_{\nu, T}$, (?) and (?) and Proposition 26 (5) show that $\tilde{\mathcal{E}}_i[Y_\rho + H_{\nu, \rho}^i | \mathcal{F}_\nu] \geq \tilde{\mathcal{E}}_i[C_* | \mathcal{F}_\nu] = C_*$, a.s. Taking the essential supremum over $(i, \rho) \in \mathcal{I} \times \mathcal{S}_{\nu, T}$ gives

$$Z(\nu) = \text{esssup}_{(i, \rho) \in \mathcal{I} \times \mathcal{S}_{\nu, T}} \tilde{\mathcal{E}}_i[Y_\rho + H_{\nu, \rho}^i | \mathcal{F}_\nu] \geq C, \quad \text{a. s.}$$

Then for any $i \in \mathcal{I}$, (?) implies that $Z^i(\nu) = Z(\nu) + H_\nu^i \geq C_* + C_H = C_Y + 2C_H$, a.s.

If no member of \mathcal{E} satisfies (8.6) (thus (4.6) is assumed), then for any $(i, \rho) \in \mathcal{I} \times \mathcal{S}_{\nu, T}$, it holds a.s. that

$$\tilde{\mathcal{E}}_i[Y_\rho^i | \mathcal{F}_t] \leq \zeta_Y, \quad \forall t \in \mathcal{D}_T.$$

Since $\tilde{\mathcal{E}}_i[Y_\rho^i|\mathcal{F}_\cdot]$ is an RCLL process, it holds except on a null set $N = N(i, \rho)$ that

$$\tilde{\mathcal{E}}_i[Y_\rho^i|\mathcal{F}_t] \leq \zeta_Y, \quad \forall t \in [0, T], \quad \langle \text{th u s} \rangle \quad \tilde{\mathcal{E}}_i[Y_\rho^i|\mathcal{F}_\nu] \leq \zeta_Y.$$

Moreover, Proposition 26 (3) and (4.4) imply that

$$\zeta_Y \geq \tilde{\mathcal{E}}_i[Y_\rho^i|\mathcal{F}_\nu] = \tilde{\mathcal{E}}_i[Y_\rho + H_{\nu, \rho}^i|\mathcal{F}_\nu] + H_\nu^i \geq \tilde{\mathcal{E}}_i[Y_\rho + H_{\nu, \rho}^i|\mathcal{F}_\nu] + C_H, \quad a. s.$$

Taking essential supremum over $(i, \rho) \in \mathcal{I} \times \mathcal{S}_{\nu, T}$ yields that

$$Z(\nu) = \operatorname{esssup}_{(i, \rho) \in \mathcal{I} \times \mathcal{S}_{\nu, T}} \tilde{\mathcal{E}}_i[Y_\rho + H_{\nu, \rho}^i|\mathcal{F}_\nu] \leq \zeta_Y - C_H, \quad a. s.$$

where $\zeta_Y - C_H \in Dom(\mathcal{E})$ thanks to (4.6) and (D2). Hence, for any $i \in \mathcal{I}$, we have $Z^i(\nu) = Z(\nu) + H_\nu^i \leq \zeta_Y - C_H + H_\nu^i$, a.s. And (4.2) together with (D2) imply that $\zeta_Y - C_H + H_\nu^i \in Dom(\mathcal{E})$. \square

Proof of Lemma 80: If no member of \mathcal{E} satisfies (8.6), then we see from Lemma 46 that

$$C \leq Z(\nu) \leq \zeta_Y - C_H, \quad a. s.,$$

and that $\zeta_Y - C_H \in Dom(\mathcal{E})$. Hence $Z(\nu) \in Dom(\mathcal{E})$ thanks to Lemma 80.

On the other hand, if \mathcal{E}_j satisfies (8.6) for some $j \in \mathcal{I}$, letting $(X, \mathcal{I}', \mathcal{U}) = (Y, \mathcal{I}, \mathcal{S}_{\nu, T})$ in Lemma 80, we can find a sequence $\{(i_n, \rho_n)\}_{n \in \mathbb{N}}$ in $\mathcal{I} \times \mathcal{S}_{\nu, T}$ such that

$$Z(\nu) = \operatorname{esssup}_{(i, \rho) \in \mathcal{I} \times \mathcal{S}_{\nu, T}} \tilde{\mathcal{E}}_i[Y_\rho + H_{\nu, \rho}^i|\mathcal{F}_\nu] = \lim_{n \rightarrow \infty} \neg \neg \uparrow \tilde{\mathcal{E}}_{i_n}[Y_{\rho_n} + H_{\nu, \rho_n}^{i_n}|\mathcal{F}_\nu], \quad a. s.$$

For any $n \in \mathbb{N}$, it follows from Definition 7.13 that there exists $k_n = k(j, i_n, \nu) \in \mathcal{I}$ such that $\tilde{\mathcal{E}}_{k_n} = \mathcal{E}_{j, i_n}^\nu$. Applying Proposition 26 (3) to $\tilde{\mathcal{E}}_{k_n}$, we can deduce from (?), (75) and (4.5) that

$$\begin{aligned} \tilde{\mathcal{E}}_{k_n}[Y_{\rho_n}^{k_n}] - C_H &= \tilde{\mathcal{E}}_{k_n}[Y_{\rho_n} + H_{\rho_n}^{k_n} - C_H] = \tilde{\mathcal{E}}_{k_n}[Y_{\rho_n} + H_{\nu, \rho_n}^{k_n} + H_\nu^{k_n} - C_H] \geq \\ &\quad \tilde{\mathcal{E}}_{k_n}[Y_{\rho_n} + H_{\nu, \rho_n}^{k_n}] \\ &= \mathcal{E}_{j, i_n}^\nu[Y_{\rho_n} + H_{\nu, \rho_n}^{k_n}] = \tilde{\mathcal{E}}_j[\tilde{\mathcal{E}}_{i_n} \neg [Y_{\rho_n} + H_{\nu, \rho_n}^{k_n}|\mathcal{F}_\nu]] = \tilde{\mathcal{E}}_j[\tilde{\mathcal{E}}_{i_n}[Y_{\rho_n} + \\ &\quad H_{\nu, \rho_n}^{i_n}|\mathcal{F}_\nu]], \end{aligned}$$

which together with (Y2) shows that

$$\liminf_{n \rightarrow \infty} \mathcal{E}_j[\tilde{\mathcal{E}}_{i_n}[Y_{\rho_n} + H_{\nu, \rho_n}^{i_n}|\mathcal{F}_\nu]] \leq \sup_{(i, \rho) \in \mathcal{I} \times \mathcal{S}_{0, T}} \tilde{\mathcal{E}}_i[Y_\rho^i] - C_H < \infty.$$

For any $n \in \mathbb{N}$, (4.8), (4.4) and Proposition 26 (5) imply that

$$\tilde{\mathcal{E}}_{i_n}[Y_{\rho_n} + H_{\nu, \rho_n}^{i_n}|\mathcal{F}_\nu] \geq \tilde{\mathcal{E}}_{i_n}[C|\mathcal{F}_\nu] = C, \quad a. s.$$

Therefore, we can deduce from Remark 73 (1) that

$$Z(\nu) = \lim_{n \rightarrow \infty} \neg \neg \uparrow \tilde{\mathcal{E}}_{i_n}[Y_{\rho_n} + H_{\nu, \rho_n}^{i_n}|\mathcal{F}_\nu] \in Dom(\mathcal{E}).$$

For any $i \in \mathcal{I}$, (4.2) and (D2) imply that $Z^i(\nu) = Z(\nu) + H_\nu^i \in \text{Dom}(\mathcal{E})$. \square

Proof of Proposition 82: To see (4.15), we first note that the event $A \triangleq \{\nu = \sigma\}$ belong to $\mathcal{F}_{\nu \wedge \sigma}$ thanks to (4.1, Lemma 1.2.16). For any $i \in \mathcal{I}$ and $\rho \in \mathcal{S}_{\nu, T}$, we define $\rho(A) \triangleq \rho \mathbf{1}_A + T \mathbf{1}_{A^c}$, which clearly belongs to $\mathcal{S}_{\sigma, T}$. Proposition 26 (2) and (3) then imply that

$$\begin{aligned} \mathbf{1}_A \tilde{\mathcal{E}}_i[Y_\rho + H_{\nu, \rho}^i | \mathcal{F}_\nu] &= \mathbf{1}_A (\tilde{\mathcal{E}}_i[Y_\rho + H_\rho^i | \mathcal{F}_\nu] - H_\nu^i) = \mathbf{1}_A (\tilde{\mathcal{E}}_i[Y_\rho + H_\rho^i | \mathcal{F}_\sigma] - H_\nu^i) \\ &= \mathbf{1}_A \tilde{\mathcal{E}}_i[Y_\rho + H_{\sigma, \rho}^i | \mathcal{F}_\sigma] \\ &= \tilde{\mathcal{E}}_i[\mathbf{1}_A (Y_{\rho(A)} + H_{\sigma, \rho(A)}^i) | \mathcal{F}_\sigma] = \mathbf{1}_A \tilde{\mathcal{E}}_i[Y_{\rho(A)} + H_{\sigma, \rho(A)}^i | \mathcal{F}_\sigma] \\ &\leq \mathbf{1}_A \operatorname{esssup}_{(i, \gamma) \in \mathcal{I} \times \mathcal{S}_{\sigma, T}} \tilde{\mathcal{E}}_i[Y_\gamma + H_{\sigma, \gamma}^i | \mathcal{F}_\sigma] = \mathbf{1}_A Z(\sigma), \quad a. s. \end{aligned}$$

Taking the essential supremum of the left-hand-side over $(i, \rho) \in \mathcal{I} \times \mathcal{S}_{\nu, T}$ and applying Lemma 80 (2), we obtain

$$\mathbf{1}_A Z(\nu) = \mathbf{1}_A \operatorname{esssup}_{(i, \rho) \in \mathcal{I} \times \mathcal{S}_{\nu, T}} \tilde{\mathcal{E}}_i[Y_\rho + H_{\nu, \rho}^i | \mathcal{F}_\nu] = \operatorname{esssup}_{(i, \rho) \in \mathcal{I} \times \mathcal{S}_{\nu, T}} (\mathbf{1}_A \tilde{\mathcal{E}}_i[Y_\rho + H_{\nu, \rho}^i | \mathcal{F}_\nu]) \leq \mathbf{1}_A Z(\sigma), \quad a. s.$$

Reversing the roles of ν and σ , we obtain (4.15).

As to (4.15), since $\mathcal{S}_{\gamma, T} \subset \mathcal{S}_{\nu, T}$, it is clear that

$$\operatorname{esssup}_{(i, \rho) \in \mathcal{I} \times \mathcal{S}_{\gamma, T}} \tilde{\mathcal{E}}_i[Y_\rho + H_{\nu, \rho}^i | \mathcal{F}_\nu] \leq \operatorname{esssup}_{(i, \rho) \in \mathcal{I} \times \mathcal{S}_{\nu, T}} \tilde{\mathcal{E}}_i[Y_\rho + H_{\nu, \rho}^i | \mathcal{F}_\nu] = Z(\nu), \quad a. s.$$

Letting $(X, \nu, \mathcal{I}', \mathcal{U}) = (Y, \gamma, \mathcal{I}, \mathcal{S}_{\gamma, T})$ in Lemma 80, we can find a sequence $\{(i_n, \rho_n)\}_{n \in \mathbb{N}}$ in $\mathcal{I} \times \mathcal{S}_{\gamma, T}$ such that

$$Z(\gamma) = \operatorname{esssup}_{(i, \rho) \in \mathcal{I} \times \mathcal{S}_{\gamma, T}} \tilde{\mathcal{E}}_i[Y_\rho + H_{\gamma, \rho}^i | \mathcal{F}_\gamma] = \lim_{n \rightarrow \infty} \neg \neg \uparrow \tilde{\mathcal{E}}_{i_n}[Y_{\rho_n} + H_{\gamma, \rho_n}^{i_n} | \mathcal{F}_\gamma], \quad a. s.$$

Now fix $j \in \mathcal{I}$. For any $n \in \mathbb{N}$, it follows from Definition 7.13 that there exists a $k_n = k(j, i_n, \gamma) \in \mathcal{I}$ such that $\tilde{\mathcal{E}}_{k_n} = \mathcal{E}_{j, i_n}^\gamma$. Applying Proposition 26 (3) to $\tilde{\mathcal{E}}_{i_n}$, we can deduce from (75), (4.5) that

$$\begin{aligned} \operatorname{esssup}_{(i, \rho) \in \mathcal{I} \times \mathcal{S}_{\gamma, T}} \tilde{\mathcal{E}}_i[Y_\rho + H_{\nu, \rho}^i | \mathcal{F}_\nu] &\geq \tilde{\mathcal{E}}_{k_n}[Y_{\rho_n} + H_{\nu, \rho_n}^{k_n} | \mathcal{F}_\nu] = \mathcal{E}_{j, i_n}^\gamma[Y_{\rho_n} + H_{\nu, \rho_n}^{k_n} | \mathcal{F}_\nu] = \\ &\quad \tilde{\mathcal{E}}_j[\tilde{\mathcal{E}}_{i_n}[Y_{\rho_n} + H_{\nu, \rho_n}^{k_n} | \mathcal{F}_\gamma] | \mathcal{F}_\nu] \\ &= \tilde{\mathcal{E}}_j[\tilde{\mathcal{E}}_{i_n}[Y_{\rho_n} + H_{\gamma, \rho_n}^{k_n} | \mathcal{F}_\gamma] + H_{\nu, \gamma}^{k_n} | \mathcal{F}_\nu] = \tilde{\mathcal{E}}_j[\tilde{\mathcal{E}}_{i_n}[Y_{\rho_n} + H_{\gamma, \rho_n}^{i_n} | \mathcal{F}_\gamma] + H_{\nu, \gamma}^j | \mathcal{F}_\nu], \\ &\quad a. s. \end{aligned} \tag{8.26}$$

For any $n \in \mathbb{N}$, Proposition 26 (5), (4.8) and (4.4) show that

$$C_Y + 2 C_H = \tilde{\mathcal{E}}_{i_n}[C | \mathcal{F}_\gamma] + C_H \leq \tilde{\mathcal{E}}_{i_n}[Y_{\rho_n} + H_{\gamma, \rho_n}^{i_n} | \mathcal{F}_\gamma] + H_{\nu, \gamma}^j \leq Z(\gamma) + H_{\nu, \gamma}^j, \quad a. s.,$$

where $Z(\gamma) + H_{\nu,\gamma}^j \in Dom(\mathcal{E}^e)$ thanks to Lemma 80, (4.2) and (D2). Then the Dominated Convergence Theorem (Proposition 29) and (8.26) imply that

$$\begin{aligned} \tilde{\mathcal{E}}_j[Z(\gamma) + H_{\nu,\gamma}^j | \mathcal{F}_\nu] &= \lim_{n \rightarrow \infty} \tilde{\mathcal{E}}_j[\tilde{\mathcal{E}}_{i_n} \neg[Y_{\rho_n} \neg + \neg H_{\gamma,\rho_n}^{i_n} | \mathcal{F}_\gamma] \neg + \neg H_{\nu,\gamma}^j | \mathcal{F}_\nu] \leq \\ &\text{esssup}_{(i,\rho) \in \mathcal{I} \times \mathcal{S}_{\gamma,T}} \neg \neg \tilde{\mathcal{E}}_i[Y_\rho \neg + \neg H_{\nu,\rho}^i | \mathcal{F}_\nu], \quad a.s. \end{aligned}$$

Taking the essential supremum of the left-hand-side over $j \in \mathcal{I}$, we obtain

$$\text{esssup}_{j \in \mathcal{I}} \tilde{\mathcal{E}}_j[Z(\gamma) + \neg H_{\nu,\gamma}^j | \mathcal{F}_\nu] \leq \text{esssup}_{(i,\rho) \in \mathcal{I} \times \mathcal{S}_{\gamma,T}} \neg \neg \tilde{\mathcal{E}}_i[Y_\rho + H_{\nu,\rho}^i | \mathcal{F}_\nu], \quad a.s. \quad (8.27)$$

On the other hand, for any $i \in \mathcal{I}$ and $\rho \in \mathcal{S}_{\gamma,T}$, applying Corollary 25 and Proposition 26 (3), we obtain

$$\begin{aligned} \tilde{\mathcal{E}}_i[Y_\rho + H_{\nu,\rho}^i | \mathcal{F}_\nu] &= \tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_i[Y_\rho + H_{\nu,\rho}^i | \mathcal{F}_\gamma] | \mathcal{F}_\nu] = \tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_i[Y_\rho + H_{\gamma,\rho}^i | \mathcal{F}_\gamma] + H_{\nu,\gamma}^i | \mathcal{F}_\nu] \\ &\leq \tilde{\mathcal{E}}_i[Z(\gamma) + H_{\nu,\gamma}^i | \mathcal{F}_\nu] \leq \text{esssup}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[Z(\gamma) + \neg H_{\nu,\gamma}^i | \mathcal{F}_\nu], \quad a.s. \end{aligned}$$

Taking the essential supremum of the left-hand-side over $(i, \rho) \in \mathcal{I} \times \mathcal{S}_{\gamma,T}$ yields that

$$\text{esssup}_{(i,\rho) \in \mathcal{I} \times \mathcal{S}_{\gamma,T}} \tilde{\mathcal{E}}_i[Y_\rho + H_{\nu,\rho}^i | \mathcal{F}_\nu] \leq \text{esssup}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[Z(\gamma) + \neg H_{\nu,\gamma}^i | \mathcal{F}_\nu], \quad a.s.,$$

which together with (8.27) proves (4.15). \square

Proof of Proposition 46: For any $i \in \mathcal{I}$ and $\nu, \gamma \in \mathcal{S}_{0,T}$ with $\nu \leq \gamma$, a.s., Proposition 26 (3), (4.15) imply that

$$\begin{aligned} \tilde{\mathcal{E}}_i[Z^i(\rho) | \mathcal{F}_\nu] &= \tilde{\mathcal{E}}_i[Z(\rho) + \neg H_{\nu,\rho}^i | \mathcal{F}_\nu] + \neg H_{\nu,\rho}^i \leq \text{esssup}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[Z(\rho) + \neg H_{\nu,\rho}^i | \mathcal{F}_\nu] + \neg \\ &H_{\nu,\rho}^i \leq Z(\nu) + \neg H_{\nu,\rho}^i = Z^i(\nu), \quad a.s., \end{aligned}$$

which implies that $\{Z^i(t)\}_{t \in [0,T]}$ is an $\tilde{\mathcal{E}}_i$ -supermartingale. Proposition 82, Theorem 20 and (?) then show that

$\left\{ Z_t^{i,+} \triangleq \liminf_{n \rightarrow \infty} Z^i(q_n^+(t)) \right\}_{t \in [0,T]}$ defines an RCLL process. Moreover, (46) implies that

$$\text{essinf}_{t \in [0,T]} Z^i(t) \geq C_Y + 2C_H, \quad a.s. \quad (8.28)$$

If \mathcal{E}_j satisfies (8.6) for some $j \in \mathcal{I}$, Corollary 25 and (8.28) imply that

$$Z_\nu^{j,+} \in Dom^\#(\mathcal{E}_j) = Dom(\mathcal{E}^e), \quad \forall \nu \in \mathcal{S}_{0,T}, \quad (8.29)$$

and that $Z^{j,+}$ is an RCLL $\tilde{\mathcal{E}}_j$ -supermartingale such that for any $t \in [0, T]$, $Z_t^{j,+} \leq Z^j(t)$, a.s. \square (8.30)

Otherwise, if no member of \mathcal{E} satisfies (8.6), we suppose that (?) holds for some $j \in \mathcal{I}$. Then Lemma 46 and (?) imply that for any $t \in \mathcal{D}_T$,

$$C_Y + 2C_H \leq Z^j(t) = Z(t) + H_t^j \leq \zeta_Y - C_H + \zeta^j, \quad a.s.$$

Taking essential supremum of $Z^j(t)$ over $t \in \mathcal{D}_T$ yields that

$$C_Y + 2C_H \leq \operatorname{esssup}_{t \in \mathcal{D}_T} Z^j(t) \leq \zeta_Y - C_H + \zeta^j, \quad a.s.,$$

where $\zeta_Y - C_H + \zeta^j \in \operatorname{Dom}(\mathcal{E})$ thanks to (4.6), (?) and (D2). Hence Lemma 80 implies that $\operatorname{esssup}_{t \in \mathcal{D}_T} Z^j(t) \in \operatorname{Dom}(\mathcal{E}) = \operatorname{Dom}^\#(\mathcal{E}_j)$. Applying Corollary 25 and (8.28) again yields (8.29) and (8.30).

To see that $Z^{j,+}$ is a modification of $\{Z^j(t)\}_{t \in [0, T]}$, it suffices to show that for any $t \in [0, T]$, $Z_t^{j,+} \geq Z^j(t)$, a.s. Fix $t \in [0, T]$. For any $(i, \nu) \in \mathcal{I} \times \mathcal{S}_{t, T}$, Definition 7.13 assures that there exists a $k = k(j, i, t) \in \mathcal{I}$ such that $\tilde{\mathcal{E}}_k = \mathcal{E}_{j, i}^t$. (S1) and (4.5) imply that

$$H_t^k = H_{0, t}^k = H_{0, t}^j = H_t^j, \quad \textcolor{blue}{\langle \text{hbox} | a n d \rangle} \quad H_{t, \nu}^k = H_{t, \nu}^i, \quad a.s. \quad (8.31)$$

For any $n \in \mathbb{N}$, we set $t_n \triangleq q_n^+(t)$ and define $\nu_n \triangleq (\nu + 2^{-n}) \wedge T \in \mathcal{S}_{t, T}$. Let $m \geq n$, it is clear that $t_m \leq t_n \leq \nu_n$, a.s. Then Proposition 26 (3) implies that

$$\tilde{\mathcal{E}}_k[Y_{\nu_n}^k | \mathcal{F}_{t_m}] = \tilde{\mathcal{E}}_k[Y_{\nu_n} + H_{t_m, \nu_n}^k | \mathcal{F}_{t_m}] + H_{t_m}^k \leq Z(t_m) + H_{t_m}^k = Z^j(t_m) + H_{t_m}^k - H_{t_m}^j, \quad a.s.$$

As $m \rightarrow \infty$, (8.31) as well as the right-continuity of the processes $\tilde{\mathcal{E}}_k[Y_{\nu_n}^k | \mathcal{F}_\cdot]$, H^k and H^j imply that

$$\tilde{\mathcal{E}}_k[Y_{\nu_n}^k | \mathcal{F}_t] = \lim_{m \rightarrow \infty} \tilde{\mathcal{E}}_k[Y_{\nu_n}^k | \mathcal{F}_{t_m}] \leq \liminf_{m \rightarrow \infty} Z^j(t_m) + H_t^k - H_t^j = \liminf_{m \rightarrow \infty} Z^j(t_m) = Z_t^{j,+}, \quad a.s.$$

Since $\lim_{n \rightarrow \infty} \neg \neg \downarrow \nu_n = \nu$, a.s., the right continuity of the process Y^k implies that $Y_{\nu_n}^k$ converges a.s. to Y_ν^k , which belongs to $\operatorname{Dom}(\mathcal{E})$ due to assumption (Y1) and (4.2). Then (4.8) and Fatou's Lemma (Theorem 16) imply that

$$\tilde{\mathcal{E}}_k[Y_\nu^k | \mathcal{F}_t] \leq \liminf_{n \rightarrow \infty} \tilde{\mathcal{E}}_k[Y_{\nu_n}^k | \mathcal{F}_t] \leq Z_t^{j,+}, \quad a.s.$$

Applying Proposition 26 (5) and (3) to $\tilde{\mathcal{E}}_j$ and $\tilde{\mathcal{E}}_i$ respectively, we can deduce from (75) and (8.31) that

$$Z_t^{j,+} \geq \tilde{\mathcal{E}}_k[Y_\nu^k | \mathcal{F}_t] = \mathcal{E}_{j, i}^t[Y_\nu^k | \mathcal{F}_t] = \tilde{\mathcal{E}}_j[\tilde{\mathcal{E}}_i[Y_\nu^k | \mathcal{F}_t] | \mathcal{F}_t] = \tilde{\mathcal{E}}_i[Y_\nu^k | \mathcal{F}_t] = \tilde{\mathcal{E}}_i[Y_\nu + H_{t, \nu}^i | \mathcal{F}_t] + H_t^j, \quad a.s. \quad (8.32)$$

Letting (i, ν) run throughout $\mathcal{I} \times \mathcal{S}_{t, T}$ yields that

$$Z_t^{j,+} \geq \operatorname{esssup}_{(i, \nu) \in \mathcal{I} \times \mathcal{S}_{t, T}} \tilde{\mathcal{E}}_i[Y_\nu + H_{t, \nu}^i | \mathcal{F}_t] + H_t^j = Z(t) + H_t^j = Z^j(t), \quad a.s.,$$

which implies that $Z^{j,+}$ is an RCLL modification of $\{Z^j(t)\}_{t \in [0, T]}$. Correspondingly, $Z^0 \triangleq \{Z_t^{j,+} - H_t^j\}_{t \in [0, T]}$ is an RCLL modification of $\{Z(t)\}_{t \in [0, T]}$. Moreover, for any $i \in \mathcal{I}$, $Z^{i,0} \triangleq \{Z_t^0 + H_t^i\}_{t \in [0, T]}$ defines an RCLL modification of $\{Z^i(t)\}_{t \in [0, T]}$, thus it is an $\tilde{\mathcal{E}}_i$ -supermartingale. \square

Proof of Proposition 4.18: For any $t \in [0, T]$, we know from (4.11) and Proposition 46 that $Y_t \leq Z(t) = Z_t^0$, a.s. Since the processes Y and Z^0 are both right continuous, it follows from Remark 73 (2) that Z^0 dominates Y .

If $\nu \in \mathcal{S}_{0,T}^F$ takes values in a finite set $\{t_1 < \dots < t_n\}$, for any $\alpha \in \{1 \dots n\}$, we can deduce from (4.15) that

$$11_{\{\nu=t_\alpha\}} Z(\nu) = 11_{\{\nu=t_\alpha\}} Z(t_\alpha) = 11_{\{\nu=t_\alpha\}} Z_{t_\alpha}^0 = 11_{\{\nu=t_\alpha\}} Z_\nu^0, \quad a. s.$$

Summing the above expression over α , we obtain

$$Z_\nu^0 = Z(\nu), \quad a. s. \quad (8.33)$$

For general stopping time $\nu \in \mathcal{S}_{0,T}$, we let $\{\nu_n\}_{n \in \mathbb{N}}$ be a decreasing sequence in $\mathcal{S}_{0,T}^F$ such that $\lim_{n \rightarrow \infty} \nu_n = \nu$, a.s. Thus for any $i \in \mathcal{I}$, the right-continuity of the process $Z^{i,0}$ shows that

$$Z_\nu^{i,0} = \lim_{n \rightarrow \infty} Z_{\nu_n}^{i,0}, \quad a. s. \quad (8.34)$$

For any $n \in \mathbb{N}$, (8.33) and (46) imply that

$$Z_{\nu_n}^{i,0} = Z^i(\nu_n) \geq C_Y + 2C_H, \quad a. s. \quad (8.35)$$

If \mathcal{E}_j satisfies (8.6) for some $j \in \mathcal{I}$, we can deduce from (4.16) and (Y2) that

$$\tilde{\mathcal{E}}_j[Z_{\nu_n}^{j,0}] = \tilde{\mathcal{E}}_j[Z^j(\nu_n)] \leq Z^j(0) = Z(0) = \sup_{(i,\rho) \in \mathcal{I} \times \mathcal{S}_{0,T}} \tilde{\mathcal{E}}_i[Y_\rho + H_\rho^i] < \infty,$$

thus $\liminf_{n \rightarrow \infty} \tilde{\mathcal{E}}_j[Z_{\nu_n}^{j,0}] < \infty$. Then Remark 73 (1) implies that $Z_\nu^{j,0} \in Dom(\mathcal{E})$.

On the other hand, if no member of \mathcal{E} satisfies (8.6), we suppose that (?) holds for some $j \in \mathcal{I}$. In light of Proposition 46 and Lemma 46, it holds a.s. that

$$C_Y + 2C_H \leq Z_t^{j,0} = Z_t^0 + H_t^j = Z(t) + H_t^j \leq \zeta_Y - C_H + \zeta^j, \quad \forall t \in \mathcal{D}_T,$$

where $\zeta_Y - C_H + \zeta^j \in Dom(\mathcal{E})$ thanks to (4.6), (?) and (D2). Since $Z^{j,0}$ is an RCLL process, it holds except on a null set N that

$$C_Y + 2C_H \leq Z_t^{j,0} \leq \zeta_Y - C_H + \zeta^j, \quad \forall t \in [0, T], \quad \langle \text{th us} \rangle \quad C_Y + 2C_H \leq Z_\nu^{j,0} \leq \zeta_Y - C_H + \zeta^j. \quad (8.36)$$

Lemma 80 then implies that $Z_\nu^{j,0} \in Dom(\mathcal{E})$. We have seen in both cases that $Z_\nu^{j,0} \in Dom(\mathcal{E})$ for some $j \in \mathcal{I}$.

Since $Z^{j,0}$ is an RCLL $\tilde{\mathcal{E}}_j$ -supermartingale by Proposition 46, (8.35) and the Optional Sampling Theorem (Theorem 24) imply that $\tilde{\mathcal{E}}_j[Z_{\nu_n}^{j,0} | \mathcal{F}_{\nu_{n+1}}] \leq Z_{\nu_{n+1}}^{j,0}$, a.s. for any $n \in \mathbb{N}$. Applying Corollary 25 and Theorem 24 once again, we obtain

$$\tilde{\mathcal{E}}_j[Z_{\nu_n}^{j,0} | \mathcal{F}_\nu] = \tilde{\mathcal{E}}_j[\tilde{\mathcal{E}}_j[Z_{\nu_n}^{j,0} | \mathcal{F}_{\nu_{n+1}}] | \mathcal{F}_\nu] \leq \tilde{\mathcal{E}}_j[Z_{\nu_{n+1}}^{j,0} | \mathcal{F}_\nu] \leq Z_\nu^{j,0}, \quad a. s., \quad (8.37)$$

which implies that $\lim_{n \rightarrow \infty} \neg \uparrow \tilde{\mathcal{E}}_j[Z_{\nu_n}^{j,0} | \mathcal{F}_\nu] \leq Z_\nu^{j,0}$, a.s. On the other hand, using (8.34) and (8.35), we can deduce from Proposition 26 (5) and Fatou's Lemma (Theorem 16) that

$$Z_\nu^{j,0} = \tilde{\mathcal{E}}_j[Z_\nu^{j,0} | \mathcal{F}_\nu] \leq \lim_{n \rightarrow \infty} \neg \uparrow \tilde{\mathcal{E}}_j[Z_{\nu_n}^{j,0} | \mathcal{F}_\nu] \leq Z_\nu^{j,0}, \quad a.s.$$

Then (8.33) and (4.16) imply that

$$Z_\nu^{j,0} = \lim_{n \rightarrow \infty} \neg \uparrow \tilde{\mathcal{E}}_j[Z_{\nu_n}^{j,0} | \mathcal{F}_\nu] = \lim_{n \rightarrow \infty} \neg \uparrow \tilde{\mathcal{E}}_j[Z^j(\nu_n) | \mathcal{F}_\nu] \leq Z^j(\nu), \quad a.s., \quad \langle \text{hbox} | \text{th us} \rangle \quad (8.38)$$

On the other hand, for any $(i, \rho) \in \mathcal{I} \times \mathcal{S}_{\nu, T}$ and $n \in \mathbb{N}$, we define $\rho_n \stackrel{\Delta}{=} \rho \vee \nu_n \in \mathcal{S}_{\nu_n, T}$. Proposition 26 (3) implies that

$$\tilde{\mathcal{E}}_i[Y_{\rho_n}^i | \mathcal{F}_{\nu_n}] = \tilde{\mathcal{E}}_i[Y_{\rho_n} + H_{\nu_n, \rho_n}^i | \mathcal{F}_{\nu_n}] + H_{\nu_n}^i \leq Z(\nu_n) + H_{\nu_n}^i = Z^i(\nu_n), \quad a.s.$$

Taking $\tilde{\mathcal{E}}_i[\cdot | \mathcal{F}_\nu]$ on both sides, we see from Corollary 25 that

$$\tilde{\mathcal{E}}_i[Y_{\rho_n}^i | \mathcal{F}_\nu] = \tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_i[Y_{\rho_n}^i | \mathcal{F}_{\nu_n}] | \mathcal{F}_\nu] \leq \tilde{\mathcal{E}}_i[Z^i(\nu_n) | \mathcal{F}_\nu], \quad a.s.$$

It is easy to see that $\lim_{n \rightarrow \infty} \neg \downarrow \rho_n = \rho$, a.s. Using the right continuity of processes Y and H^i , we can deduce from (4.8), Fatou's Lemma (Proposition 28) and (8.38) that

$$\tilde{\mathcal{E}}_i[Y_\rho^i | \mathcal{F}_\nu] \leq \liminf_{n \rightarrow \infty} \tilde{\mathcal{E}}_i[Y_{\rho_n}^i | \mathcal{F}_\nu] \leq \lim_{n \rightarrow \infty} \neg \uparrow \tilde{\mathcal{E}}_i[Z^i(\nu_n) | \mathcal{F}_\nu] = Z_\nu^{i,0}, \quad a.s.$$

Then Proposition 26 (3) again implies that

$$\tilde{\mathcal{E}}_i[Y_\rho + H_{\nu, \rho}^i | \mathcal{F}_\nu] = \tilde{\mathcal{E}}_i[Y_\rho^i | \mathcal{F}_\nu] - H_\nu^i \leq Z_\nu^{i,0} - H_\nu^i = Z_\nu^0, \quad a.s.$$

Taking the essential supremum over $(i, \rho) \in \mathcal{I} \times \mathcal{S}_{\nu, T}$ yields that $Z(\nu) \leq Z_\nu^0$, a.s., which in conjunction with (8.38) shows that $Z_\nu^0 = Z(\nu)$, a.s., thus $Z_\nu^0 \in \text{Dom}(\mathcal{E})$ by Lemma 80. Moreover, for any $i \in \mathcal{I}$, we have

$$Z_\nu^{i,0} = Z_\nu^0 + H_\nu^i = Z(\nu) + H_\nu^i = Z^i(\nu), \quad a.s.,$$

thus $Z_\nu^{i,0} \in \text{Dom}(\mathcal{E})$ thanks to Lemma 80 once again. (4.17) is proved.

Now let X be another RCLL \mathbf{F} -adapted process dominating Y such that $X^i \stackrel{\Delta}{=} \{X_t + H_t^i\}_{t \in [0, T]}$ is an $\tilde{\mathcal{E}}_i$ -supermartingale for any $i \in \mathcal{I}$. We fix $t \in [0, T]$. For any $i \in \mathcal{I}$ and $\nu \in \mathcal{S}_{t, T}$, we let $\{\nu_n\}_{n \in \mathbb{N}}$ be a decreasing sequence in $\mathcal{S}_{t, T}^F$ such that $\lim_{n \rightarrow \infty} \neg \downarrow \nu_n = \nu$, a.s. For any $n \in \mathbb{N}$, since X^i dominates Y^i , Remark 73 (1) shows that $X_{\nu_n}^i \geq Y_{\nu_n}^i$, a.s. Moreover, (4.8) implies that

$$\text{essinf}_{t \in \mathcal{D}_T} X_t^i \geq \text{essinf}_{t \in \mathcal{D}_T} Y_t^i \geq C, \quad a.s.$$

(fi) Then (A4), Proposition 82 and the Optional Sampling Theorem (Theorem 24) imply that

$$\tilde{\mathcal{E}}_i[Y_{\nu_n} + H_{t, \nu_n}^i | \mathcal{F}_t] = \tilde{\mathcal{E}}_i[Y_{\nu_n}^i | \mathcal{F}_t] - H_t^i \leq \tilde{\mathcal{E}}_i[X_{\nu_n}^i | \mathcal{F}_t] - H_t^i \leq X_t^i - H_t^i = X_t, \quad a.s.$$

The right-continuity of the processes Y and H^i shows that $Y_\nu + H_{t,\nu}^i = \lim_{n \rightarrow \infty} (Y_{\nu_n} + H_{t,\nu_n}^i)$, a.s., thus it follows from (??), (??) and Fatou's Lemma (Proposition 28) that

$$\tilde{\mathcal{E}}_i[Y_\nu + H_{t,\nu}^i | \mathcal{F}_t] \leq \liminf_{n \rightarrow \infty} \tilde{\mathcal{E}}_i[Y_{\nu_n} + H_{t,\nu_n}^i | \mathcal{F}_t] \leq X_t, \quad a.s.$$

Taking the essential supremum of the left-hand-side over $(i, \nu) \in \mathcal{I} \times \mathcal{S}_{t,T}$, we can deduce from Proposition 46 that

$$Z_t^0 = Z(t) = \operatorname{esssup}_{(i,\nu) \in \mathcal{I} \times \mathcal{S}_{t,T}} \tilde{\mathcal{E}}_i[Y_\nu + H_{t,\nu}^i | \mathcal{F}_t] \leq X_t, \quad a.s.$$

Since both Z^0 and X are RCLL processes, Remark 73 (2) once again shows that X dominates Z^0 . \square

Proof of Lemma 80: For any $i \in \mathcal{I}$, (4.18), (4.4) as well as Proposition 26 (5) imply that

$$\tilde{\mathcal{E}}_i[Z_{\tau_\delta(\nu)}^0 + H_{\nu,\tau_\delta(\nu)}^i | \mathcal{F}_\nu] \geq \tilde{\mathcal{E}}_i[C + C_H | \mathcal{F}_\nu] = C_Y + 2C_H, \quad a.s.$$

Taking the essential supremum of the left-hand-side over $i \in \mathcal{I}$, we can deduce from (4.19) that

$$C_Y + 2C_H \leq \operatorname{esssup}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[Z_{\tau_\delta(\nu)}^0 + H_{\nu,\tau_\delta(\nu)}^i | \mathcal{F}_\nu] = J_\delta(\nu) \leq Z(\nu), \quad a.s. \quad (8.39)$$

Then Lemma 80 imply that $J_\delta(\nu) \in Dom(\mathcal{E})$. Let σ be another stopping time in $\mathcal{S}_{0,T}$. In light of (4.17) and (4.15), we see that

$$\begin{aligned} \mathbf{1}_{\{\tau_\delta(\nu)=\tau_\delta(\sigma)\}} Z_{\tau_\delta(\nu)}^0 &= \mathbf{1}_{\{\tau_\delta(\nu)=\tau_\delta(\sigma)\}} Z(\tau_\delta(\nu)) = \mathbf{1}_{\{\tau_\delta(\nu)=\tau_\delta(\sigma)\}} Z(\tau_\delta(\sigma)) = \\ &= \mathbf{1}_{\{\tau_\delta(\nu)=\tau_\delta(\sigma)\}} Z_{\tau_\delta(\sigma)}^0, \quad a.s. \end{aligned} \quad (8.40)$$

It is clear that $\{\nu = \sigma\} \subset \{\tau_\delta(\nu) = \tau_\delta(\sigma)\}$. Thus multiplying $\mathbf{1}_{\{\nu=\sigma\}}$ to both sides of (8.40) gives that

$$\mathbf{1}_{\{\nu=\sigma\}} Z_{\tau_\delta(\nu)}^0 = \mathbf{1}_{\{\nu=\sigma\}} Z_{\tau_\delta(\sigma)}^0, \quad a.s.$$

For any $i \in \mathcal{I}$, applying Proposition 26 (2) and recalling how $\tilde{\mathcal{E}}_i[\cdot | \mathcal{F}_\nu]$ and $\tilde{\mathcal{E}}_i[\cdot | \mathcal{F}_\sigma]$ are defined in (2.7), we obtain

$$\begin{aligned} \mathbf{1}_{\{\nu=\sigma\}} \tilde{\mathcal{E}}_i[Z_{\tau_\delta(\nu)}^0 + H_{\nu,\tau_\delta(\nu)}^i | \mathcal{F}_\nu] &= \mathbf{1}_{\{\nu=\sigma\}} \tilde{\mathcal{E}}_i[Z_{\tau_\delta(\nu)}^0 + H_{\nu,\tau_\delta(\nu)}^i | \mathcal{F}_\sigma] = \\ &= \tilde{\mathcal{E}}_i[\mathbf{1}_{\{\nu=\sigma\}} Z_{\tau_\delta(\sigma)}^0 + \mathbf{1}_{\{\nu=\sigma\}} H_{\sigma,\tau_\delta(\sigma)}^i | \mathcal{F}_\sigma] \\ &= \mathbf{1}_{\{\nu=\sigma\}} \tilde{\mathcal{E}}_i[Z_{\tau_\delta(\sigma)}^0 + H_{\sigma,\tau_\delta(\sigma)}^i | \mathcal{F}_\sigma], \quad a.s., \end{aligned}$$

where we use the fact that $\{\nu = \sigma\} \in \mathcal{F}_{\nu \wedge \sigma}$ thanks to (??, Lemma 1.2.16). Taking the essential supremum of both sides over $i \in \mathcal{I}$, Lemma 80 (2) implies that

$$\begin{aligned} \mathbf{1}_{\{\nu=\sigma\}} J_\delta(\nu) &= \operatorname{esssup}_{i \in \mathcal{I}} \mathbf{1}_{\{\nu=\sigma\}} \tilde{\mathcal{E}}_i[Z_{\tau_\delta(\nu)}^0 + H_{\nu,\tau_\delta(\nu)}^i | \mathcal{F}_\nu] = \\ &= \operatorname{esssup}_{i \in \mathcal{I}} \mathbf{1}_{\{\nu=\sigma\}} \tilde{\mathcal{E}}_i[Z_{\tau_\delta(\sigma)}^0 + H_{\sigma,\tau_\delta(\sigma)}^i | \mathcal{F}_\sigma] = \mathbf{1}_{\{\nu=\sigma\}} J_\delta(\sigma), \quad a.s., \end{aligned}$$

which proves the lemma. \square

Proof of Proposition 82:

Proof of 1. We fix $i \in \mathcal{I}$ and $\nu, \rho \in \mathcal{S}_{0,T}$ with $\nu \leq \rho$, a.s. Taking $(\nu, \mathcal{I}', \mathcal{U}) = (\rho, \mathcal{I}, \{\tau_\delta(\rho)\})$ and $X(\tau_\delta(\rho)) = Z_{\tau_\delta(\rho)}^0$ in Lemma 80, we can find a sequence $\{j_n\}_{n=1}^\infty$ in \mathcal{I} such that

$$J_\delta(\rho) = \operatorname{esssup}_{j \in \mathcal{I}} \tilde{\mathcal{E}}_j[Z_{\tau_\delta(\rho)}^0 + H_{\rho, \tau_\delta(\rho)}^j | \mathcal{F}_\rho] = \lim_{n \rightarrow \infty} \neg \neg \uparrow \tilde{\mathcal{E}}_{j_n}[Z_{\tau_\delta(\rho)}^0 + H_{\rho, \tau_\delta(\rho)}^{j_n} | \mathcal{F}_\rho],$$

a. s.

For any $n \in \mathbb{N}$, it follows from Definition 7.13 that there exists a $k_n = k(i, j_n, \rho) \in \mathcal{I}$ such that $\tilde{\mathcal{E}}_{k_n} = \mathcal{E}_{i, j_n}^\rho$.

$$H_{\tau_\delta(\rho)}^{k_n} - H_\rho^{k_n} = H_{\tau_\delta(\rho)}^{j_n} - H_\rho^{j_n} \quad (\text{hbox | a n d}) \quad H_\rho^{k_n} - H_\nu^{k_n} = H_\rho^i - H_\nu^i,$$

a. s. (8.41)

(fi) Applying Proposition 26 (3) to $\tilde{\mathcal{E}}_{j_n}$, we can deduce from (75) and (4.5) that

$$\begin{aligned} \tilde{\mathcal{E}}_{k_n}[Z_{\tau_\delta(\rho)}^0 + H_{\nu, \tau_\delta(\rho)}^{k_n} | \mathcal{F}_\nu] &= \mathcal{E}_{i, j_n}^\rho[Z_{\tau_\delta(\rho)}^0 + H_{\nu, \tau_\delta(\rho)}^{k_n} | \mathcal{F}_\nu] = \tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_{j_n}[Z_{\tau_\delta(\rho)}^0 + \\ &\quad H_{\nu, \tau_\delta(\rho)}^{k_n} | \mathcal{F}_\rho] | \mathcal{F}_\nu] \\ &= \tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_{j_n}[Z_{\tau_\delta(\rho)}^0 + H_{\rho, \tau_\delta(\rho)}^{j_n} | \mathcal{F}_\rho] + H_{\nu, \rho}^i | \mathcal{F}_\nu], \end{aligned}$$

a. s. (8.42)

Since $\nu \leq \rho$, a.s., we see that $\tau_\delta(\nu) \leq \tau_\delta(\rho)$, a.s. Due to (4.17) and (4.15), we have that

$$\begin{aligned} \tilde{\mathcal{E}}_{k_n}[Z_{\tau_\delta(\rho)}^0 + H_{\tau_\delta(\nu), \tau_\delta(\rho)}^{k_n} | \mathcal{F}_{\tau_\delta(\nu)}] &\leq \operatorname{esssup}_{j \in \mathcal{I}} \tilde{\mathcal{E}}_j[Z(\tau_\delta(\rho)) + H_{\tau_\delta(\nu), \tau_\delta(\rho)}^j | \mathcal{F}_{\tau_\delta(\nu)}] \leq \\ Z(\tau_\delta(\nu)) &= Z_{\tau_\delta(\nu)}^0, \quad \text{a. s.} \end{aligned}$$

Then using Corollary 25 and applying Proposition 26 (3) to $\tilde{\mathcal{E}}_{k_n}$, we obtain

$$\begin{aligned} \tilde{\mathcal{E}}_{k_n}[Z_{\tau_\delta(\rho)}^0 + H_{\nu, \tau_\delta(\rho)}^{k_n} | \mathcal{F}_\nu] &= \tilde{\mathcal{E}}_{k_n}[\tilde{\mathcal{E}}_{k_n}[Z_{\tau_\delta(\rho)}^0 + H_{\nu, \tau_\delta(\rho)}^{k_n} | \mathcal{F}_{\tau_\delta(\nu)}] | \mathcal{F}_\nu] \\ &= \tilde{\mathcal{E}}_{k_n}[\tilde{\mathcal{E}}_{k_n}[Z_{\tau_\delta(\rho)}^0 + H_{\tau_\delta(\nu), \tau_\delta(\rho)}^{k_n} | \mathcal{F}_{\tau_\delta(\nu)}] + H_{\nu, \tau_\delta(\nu)}^{k_n} | \mathcal{F}_\nu] \\ &\leq \tilde{\mathcal{E}}_{k_n}[Z_{\tau_\delta(\nu)}^0 + H_{\nu, \tau_\delta(\nu)}^{k_n} | \mathcal{F}_\nu] \leq \operatorname{esssup}_{j \in \mathcal{I}} \tilde{\mathcal{E}}_j[Z_{\tau_\delta(\nu)}^0 + H_{\nu, \tau_\delta(\nu)}^j | \mathcal{F}_\nu] = J_\delta(\nu), \quad \text{a. s.,} \end{aligned}$$

which together with (8.41) shows that

$$\tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_{j_n}[Z_{\tau_\delta(\rho)}^0 + H_{\rho, \tau_\delta(\rho)}^{j_n} | \mathcal{F}_\rho] + H_{\nu, \rho}^i | \mathcal{F}_\nu] \leq J_\delta(\nu), \quad \text{a. s.}$$

For any $n \in \mathbb{N}$, we see from (4.18), (4.4) and Proposition 26 (5) that

$$\tilde{\mathcal{E}}_{j_n}[Z_{\tau_\delta(\rho)}^0 + H_{\rho, \tau_\delta(\rho)}^{j_n} | \mathcal{F}_\rho] + H_{\nu, \rho}^i \geq \tilde{\mathcal{E}}_{j_n}[C_Y + 2C_H | \mathcal{F}_\rho] + C_H = C_Y + 3C_H, \quad \text{a. s.}$$

Then Fatou's Lemma (Proposition 28) implies that

$$\tilde{\mathcal{E}}_i[J_\delta(\rho) + H_{\nu, \rho}^i | \mathcal{F}_\nu] \leq \lim_{n \rightarrow \infty} \neg \neg \uparrow \tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_{j_n}[Z_{\tau_\delta(\rho)}^0 + H_{\rho, \tau_\delta(\rho)}^{j_n} | \mathcal{F}_\rho] + H_{\nu, \rho}^i | \mathcal{F}_\nu] \leq J_\delta(\nu),$$

a. s.

For any $\sigma \in \mathcal{S}_{0,T}$, Lemma 80, (4.2) and (D2) show that $J_\delta^i(\sigma) \stackrel{\Delta}{=} J_\delta(\sigma) + H_\sigma^i \in \text{Dom}(\mathcal{E})$. A simple application of Proposition 26 (3) yields that

$$\tilde{\mathcal{E}}_i[J_\delta^i(\rho)|\mathcal{F}_\nu] = \tilde{\mathcal{E}}_i[J_\delta(\rho) + H_{\nu,\rho}^i|\mathcal{F}_\nu] + H_\nu^i \leq J_\delta(\nu) + H_\nu^i = J_\delta^i(\nu), \quad a.s. \quad (8.43)$$

In particular, when $0 \leq s < t \leq T$, we have $\tilde{\mathcal{E}}_i[J_\delta^i(t)|\mathcal{F}_s] \leq J_\delta^i(s)$, a.s., which show that $\{J_\delta^i(t)\}_{t \in [0,T]}$ is an $\tilde{\mathcal{E}}_i$ -supermartingale.

Proof of 2. For any $i \in \mathcal{I}$ and $\nu \in \mathcal{S}_{0,T}$, (8.39) and (?) imply that

$$J_\delta^i(\nu) = J_\delta(\nu) + H_\nu^i \geq C_Y + 3C_H, \quad a.s. \quad (8.44)$$

In particular, $J_\delta^i(T) \geq C_Y + 3C_H$, a.s. Proposition 82 and Theorem 20 then show that $\left\{J_t^{\delta,i,+} \stackrel{\Delta}{=} \liminf_{n \rightarrow \infty} J_\delta^i(q_n^+(t))\right\}_{t \in [0,T]}$ defines an RCLL process. For any $t \in [0, T]$ and $\tilde{i} \in \mathcal{I}$, (4.18), (4.4) as well as Proposition 26 (5) imply that

$$\tilde{\mathcal{E}}_{\tilde{i}}[Z_{\tau_\delta(t)}^0 + H_{t,\tau_\delta(t)}^{\tilde{i}}|\mathcal{F}_t] \geq \tilde{\mathcal{E}}_{\tilde{i}}[C_Y + 2C_H|\mathcal{F}_t] = C_Y + 2C_H, \quad a.s.$$

Taking essential supremum over $\tilde{i} \in \mathcal{I}$, we can deduce from (4.19) that

$$C_Y + 2C_H \leq \text{esssup}_{\tilde{i} \in \mathcal{I}} \tilde{\mathcal{E}}_{\tilde{i}}[Z_{\tau_\delta(t)}^0 + H_{t,\tau_\delta(t)}^{\tilde{i}}|\mathcal{F}_t] = J_\delta(t) \leq Z(t), \quad a.s.,$$

Then (4.4) shows that (fi) Then (8.44) implies that

$$\text{essinf}_{t \in [0,T]} J_\delta^i(t) \geq C_Y + 3C_H, \quad a.s. \quad (8.45)$$

If \mathcal{E}_j satisfies (8.6) for some $j \in \mathcal{I}$, Corollary 25 and (8.45) imply that

$$J_\nu^{\delta,j,+} \in \text{Dom}^\#(\mathcal{E}_j) = \text{Dom}(\mathcal{E}), \quad \forall \nu \in \mathcal{S}_{0,T}, \quad (8.46)$$

$$\langle \text{hbox} | \text{and that } J^{\delta,j,+} \text{ is an RCLL } \tilde{\mathcal{E}}_j \text{ - supermartingale such that for any } t \in [0, T], J_t^{\delta,j,+} \leq J_\delta^j(t), \text{ a.s.} \rangle \quad (8.47)$$

Otherwise, if no member of \mathcal{E} satisfies (8.6), we suppose that (?) holds for some $j \in \mathcal{I}$. Then (8.44), (4.19) and (?) imply that for any $t \in \mathcal{D}_T$,

$$C_Y + 3C_H \leq J_\delta^j(t) = J_\delta(t) + H_t^j \leq Z(t) + H_t^j \leq \zeta_Y - C_H + \zeta^j, \quad a.s.$$

Taking essential supremum of $J_\delta^j(t)$ over $t \in \mathcal{D}_T$ yields that

$$C_Y + 3C_H \leq \text{esssup}_{t \in \mathcal{D}_T} J_\delta^j(t) \leq \zeta_Y - C_H + \zeta^j, \quad a.s.,$$

where $\zeta_Y - C_H + \zeta^j \in \text{Dom}(\mathcal{E})$ thanks to (4.6), (?) and (D2). Hence Lemma 80 implies that $\text{esssup}_{t \in \mathcal{D}_T} J_\delta^j(t) \in \text{Dom}(\mathcal{E}) = \text{Dom}^\#(\mathcal{E}_j)$. Applying Corollary 25 and (8.45) again yields (8.46) and (8.47).

To see that $J^{\delta,j,+}$ is a modification of $\{J_\delta^j(t)\}_{t \in [0,T]}$, it suffices to show that for any $t \in [0, T]$, $J_t^{\delta,j,+} \geq J_\delta^j(t)$, a.s. Fix $t \in [0, T]$. For any $i \in \mathcal{I}$, Definition 7.13 assures that there exists a $k = k(j, i, t) \in \mathcal{I}$ such that $\tilde{\mathcal{E}}_k = \mathcal{E}_{j,i}^t$. Moreover, (S1) and (4.5) imply that

$$H_t^k = H_{0,t}^k = H_{0,t}^j = H_t^j, \quad \langle \text{hbox} | \text{and} \rangle \quad H_{t,\tau_\delta(t)}^k = H_{t,\tau_\delta(t)}^i, \quad a.s. \quad (8.48)$$

For any $n \in \mathbb{N}$, we set $t_n \stackrel{\Delta}{=} q_n^+(t)$. Let $m \geq n$, it is clear that $t_m \leq t_n$, a.s. Then (4.17), Corollary 25, Proposition 26 (3) as well as (4.15) imply that

$$\begin{aligned}
\tilde{\mathcal{E}}_k[Z_{\tau_\delta(t_n)}^{k,0}|\mathcal{F}_{t_m}] &= \tilde{\mathcal{E}}_k[Z^k(\tau_\delta(t_n))|\mathcal{F}_{t_m}] = \tilde{\mathcal{E}}_k[\tilde{\mathcal{E}}_k[Z^k(\tau_\delta(t_n))|\mathcal{F}_{\tau_\delta(t_m)}]|\mathcal{F}_{t_m}] \\
&= \tilde{\mathcal{E}}_k[\tilde{\mathcal{E}}_k[Z(\tau_\delta(t_n)) + H_{\tau_\delta(t_m), \tau_\delta(t_n)}^k|\mathcal{F}_{\tau_\delta(t_m)}] + H_{t_m, \tau_\delta(t_m)}^k|\mathcal{F}_{t_m}] + H_{t_m}^k \\
&\leq \tilde{\mathcal{E}}_k\left[\text{esssup}_{l \in \mathcal{I}} \tilde{\mathcal{E}}_l[Z(\tau_\delta(t_n)) + H_{\tau_\delta(t_m), \tau_\delta(t_n)}^l|\mathcal{F}_{\tau_\delta(t_m)}] + H_{t_m, \tau_\delta(t_m)}^k\right] + H_{t_m}^k \\
&\leq \tilde{\mathcal{E}}_k[Z(\tau_\delta(t_m)) + H_{t_m, \tau_\delta(t_m)}^k|\mathcal{F}_{t_m}] + H_{t_m}^k \leq \text{esssup}_{l \in \mathcal{I}} \tilde{\mathcal{E}}_l[Z_{\tau_\delta(t_m)}^{k,0} + H_{t_m, \tau_\delta(t_m)}^l|\mathcal{F}_{t_m}] + H_{t_m}^k \\
&= J_\delta(t_m) + H_{t_m}^k = J_\delta^j(t_m) + H_{t_m}^k - H_{t_m}^j, \quad a.s. \tag{8.49}
\end{aligned}$$

As $m \rightarrow \infty$, (8.48) as well as the right-continuity of the processes $\tilde{\mathcal{E}}_k[Z_{\tau_\delta(t_n)}^{k,0}|\mathcal{F}_\cdot]$, H^k and H^j imply that

$$\begin{aligned}
\tilde{\mathcal{E}}_k[Z_{\tau_\delta(t_n)}^{k,0}|\mathcal{F}_t] &= \lim_{m \rightarrow \infty} \tilde{\mathcal{E}}_k[Z_{\tau_\delta(t_n)}^{k,0}|\mathcal{F}_{t_m}] \leq \liminf_{m \rightarrow \infty} J_\delta^j(t_m) + H_t^k - H_t^j = \\
\liminf_{m \rightarrow \infty} J_\delta^j(t_m) &= J_t^{\delta, j, +}, \quad a.s.
\end{aligned}$$

Since $\lim_{n \rightarrow \infty} \neg \neg \tau_\delta(t_n) = \tau_\delta(t)$ a.s., the right-continuity of the process $Z^{k,0}$ implies that $Z_{\tau_\delta(t_n)}^{k,0}$ converges a.s. to $Z_{\tau_\delta(t)}^{k,0}$, which belongs to $D \circ m(\mathcal{E})$ thanks to Proposition 4.18. Then (4.18) and Fatou's Lemma (Theorem 16) imply that

$$\tilde{\mathcal{E}}_k[Z_{\tau_\delta(t)}^{k,0}|\mathcal{F}_t] \leq \liminf_{n \rightarrow \infty} \tilde{\mathcal{E}}_k[Z_{\tau_\delta(t_n)}^{k,0}|\mathcal{F}_t] \leq J_t^{\delta, j, +}, \quad a.s.$$

Similar to (8.32), we can deduce from (75) and (8.48) that

$$J_t^{\delta, j, +} \geq \tilde{\mathcal{E}}_i[Z_{\tau_\delta(t)}^0 + H_{t, \tau_\delta(t)}^i|\mathcal{F}_t] + H_t^j, \quad a.s.$$

Letting i run throughout \mathcal{I} yields that

$$J_t^{\delta, j, +} \geq \text{esssup}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[Z_{\tau_\delta(t)}^0 + H_{t, \tau_\delta(t)}^i|\mathcal{F}_t] + H_t^j = J_\delta(t) + H_t^j = J_\delta^j(t), \quad a.s.,$$

which implies that $J^{\delta, j, +}$ is an RCLL modification of $\{J_\delta^j(t)\}_{t \in [0, T]}$. Correspondingly, $J^{\delta, 0} \stackrel{\Delta}{=} \{J_t^{\delta, j, +} - H_t^j\}_{t \in [0, T]}$ is an RCLL modification of $\{J_\delta(t)\}_{t \in [0, T]}$. Moreover, for any $i \in \mathcal{I}$, $J^{\delta, i, 0} \stackrel{\Delta}{=} \{J_t^{\delta, 0} + H_t^i\}_{t \in [0, T]}$ defines an RCLL modification of $\{J_\delta^i(t)\}_{t \in [0, T]}$, thus it is an $\tilde{\mathcal{E}}_i$ -supermartingale.

Proof of 3. Now let us show (3). Similar to (8.33), we can deduce from Lemma 80 that for any $\nu \in \mathcal{S}_{0, T}^F$

$$J_\nu^{\delta, 0} = J_\delta(\nu), \quad a.s. \tag{8.50}$$

For a general stopping time $\nu \in \mathcal{S}_{0,T}$, we let $\{\nu_n\}_{n \in \mathbb{N}}$ be a decreasing sequence in $\mathcal{S}_{0,T}^F$ such that $\lim_{n \rightarrow \infty} \neg \downarrow \nu_n = \nu$, a.s. Thus for any $i \in \mathcal{I}$, the right-continuity of the process $J^{\delta,i,0}$ shows that

$$J_\nu^{\delta,i,0} = \lim_{n \rightarrow \infty} J_{\nu_n}^{\delta,i,0}, \quad a.s. \quad (8.51)$$

In light of (8.50) and (8.44), it holds a.s. that

$$J_t^{\delta,i,0} = J_\delta^i(t) \geq C_Y + 3C_H, \quad \forall t \in \mathcal{D}_T.$$

Since $J^{\delta,i,0}$ is an RCLL process, it holds except on a null set N that

$$J_t^{\delta,i,0} \geq C_Y + 3C_H, \quad \forall t \in [0, T], \quad \langle \text{hbox} | t h u s \rangle \quad J_\sigma^{\delta,i,0} \geq C_Y + 3C_H, \quad \forall \sigma \in \mathcal{S}_{0,T}. \quad (8.52)$$

If \mathcal{E}_j satisfies (8.6) for some $j \in \mathcal{I}$, we can deduce from (8.43), (4.19) and (Y2) that

$$\tilde{\mathcal{E}}_j[J_{\nu_n}^{\delta,j,0}] = \tilde{\mathcal{E}}_j[J_\delta^j(\nu_n)] \leq J_\delta^j(0) = J_\delta(0) \leq Z(0) = \sup_{(i,\rho) \in \mathcal{I} \times \mathcal{S}_{0,T}} \tilde{\mathcal{E}}_i[Y_\rho + H_\rho^i] < \infty,$$

thus $\liminf_{n \rightarrow \infty} \tilde{\mathcal{E}}_j[J_{\nu_n}^{\delta,j,0}] < \infty$. Then Remark 73 (1) implies that $J_\nu^{\delta,j,0} \in Dom(\mathcal{E})$.

On the other hand, if no member of \mathcal{E} satisfies (8.6), we suppose that (?) holds for some $j \in \mathcal{I}$. In light of (8.50), (4.19) and (?), it holds a.s. that

$$J_t^{\delta,j,0} = J_\delta^j(t) = J_\delta(t) + H_t^j \leq Z(t) + H_t^j \leq \zeta_Y - C_H + \zeta^j, \quad \forall t \in \mathcal{D}_T,$$

where $\zeta_Y - C_H + \zeta^j \in Dom(\mathcal{E})$ thanks to (4.6), (?) and (D2). Since $J^{\delta,j,0}$ is an RCLL process, it holds except on a null set N' that

$$J_t^{\delta,j,0} \leq \zeta_Y - C_H + \zeta^j, \quad \forall t \in [0, T], \quad \langle \text{hbox} | t h u s \rangle \quad J_\nu^{\delta,j,0} \leq \zeta_Y - C_H + \zeta^j. \quad (8.53)$$

Then (8.52) and Lemma 80 imply that $J_\nu^{\delta,j,0} \in Dom(\mathcal{E})$. We have seen in both cases that $J_\nu^{\delta,j,0} \in Dom(\mathcal{E})$ for some $j \in \mathcal{I}$.

Similar to the arguments used in (8.37) through (8.38) (with (8.50)-(8.52) replacing (8.33)-(8.35) respectively, and with (8.43) replacing (4.16)), we can deduce that

$$J_\nu^{\delta,j,0} = \lim_{n \rightarrow \infty} \neg \uparrow \tilde{\mathcal{E}}_j[J_{\nu_n}^{\delta,j,0} | \mathcal{F}_\nu] = \lim_{n \rightarrow \infty} \neg \uparrow \tilde{\mathcal{E}}_j[J_\delta^j(\nu_n) | \mathcal{F}_\nu] \leq J_\delta^j(\nu), \quad \langle \text{hbox} | t h u s \rangle \quad J_\nu^{\delta,0} \leq J_\delta(\nu), \quad a.s. \quad (8.54)$$

The right-continuity of the process $J^{\delta,0}$, (8.50) and (8.39) show that

$$J_\nu^{\delta,0} = \lim_{n \rightarrow \infty} J_{\nu_n}^{\delta,0} = \lim_{n \rightarrow \infty} J_\delta(\nu_n) \geq C_Y + 2C_H, \quad a.s.$$

Lemma 80 and Lemma 80 thus imply that $J_\nu^{\delta,0} \in Dom(\mathcal{E})$. For any $i \in \mathcal{I}$, (4.2) and (D2) show that $J_\nu^{\delta,i,0} = J_\nu^{\delta,0} + H_\nu^i \in Dom(\mathcal{E})$.

On the other hand, for any $i \in \mathcal{I}$ and $n \in \mathbb{N}$, it is clear that $\nu \leq \nu_n \leq \tau_\delta(\nu_n)$, a.s. Then Corollary 25, (8.54) and (8.43) imply that

$$\begin{aligned} \tilde{\mathcal{E}}_i[J_{\tau_\delta(\nu_n)}^{\delta,i,0}|\mathcal{F}_\nu] &= \tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_i[J_{\tau_\delta(\nu_n)}^{\delta,i,0}|\mathcal{F}_{\nu_n}]|\mathcal{F}_\nu] \leq \tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_i[J_\delta^i(\tau_\delta(\nu_n))|\mathcal{F}_{\nu_n}]|\mathcal{F}_\nu] \leq \\ &\tilde{\mathcal{E}}_i[J_\delta^i(\nu_n)|\mathcal{F}_\nu], \quad a.s. \end{aligned}$$

It is easy to see that $\lim_{n \rightarrow \infty} \neg \neg \downarrow \tau_\delta(\nu_n) = \tau_\delta(\nu)$, a.s. Using the right continuity of the process $J^{\delta,i,0}$, we can deduce from (8.52), Fatou's Lemma (Proposition 28) and (8.54) that

$$\tilde{\mathcal{E}}_i[J_{\tau_\delta(\nu)}^{\delta,i,0}|\mathcal{F}_\nu] \leq \liminf_{n \rightarrow \infty} \tilde{\mathcal{E}}_i[J_{\tau_\delta(\nu_n)}^{\delta,i,0}|\mathcal{F}_\nu] \leq \lim_{n \rightarrow \infty} \neg \neg \uparrow \tilde{\mathcal{E}}_i[J_\delta^i(\nu_n)|\mathcal{F}_\nu] = J_\nu^{\delta,i,0}, \quad a.s.$$

Proposition 26 (3) further implies that

$$\tilde{\mathcal{E}}_i[J_{\tau_\delta(\nu)}^{\delta,i,0} + H_{\nu, \tau_\delta(\nu)}^i|\mathcal{F}_\nu] = \tilde{\mathcal{E}}_i[J_{\tau_\delta(\nu)}^{\delta,i,0}|\mathcal{F}_\nu] - H_\nu^i \leq J_\nu^{\delta,i,0} - H_\nu^i = J_\nu^{\delta,0}, \quad a.s.$$

Taking the essential supremum over $i \in \mathcal{I}$ gives

$$J_\delta(\nu) = \operatorname{esssup}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[J_{\tau_\delta(\nu)}^{\delta,0} + H_{\nu, \tau_\delta(\nu)}^i|\mathcal{F}_\nu] \leq J_\nu^{\delta,0}, \quad a.s.,$$

which together with (8.54) shows that $J_\nu^{\delta,0} = J_\delta(\nu)$, a.s. \square

Proof of Theorem 57: We first show that for any $\delta \in (0, 1)$ and $\nu \in \mathcal{S}_{0,T}$

$$J_\delta(\nu) = Z_\nu^0 = Z(\nu), \quad a.s. \quad (8.55)$$

Fix $i \in \mathcal{I}$. Lemma 80 indicates that $\tilde{\mathcal{E}}_i$ is a convex \mathbf{F} -expectation on $Dom(\mathcal{E})$. Since $Z^{i,0}$ and $J^{\delta,i,0}$ are both $\tilde{\mathcal{E}}_i$ -supermartingales, we can deduce that for any $0 \leq s < t \leq T$,

$$\begin{aligned} \tilde{\mathcal{E}}_i[\delta Z_t^0 + (1 - \delta) J_t^{\delta,i,0} + H_t^i|\mathcal{F}_s] &= \tilde{\mathcal{E}}_i[\delta Z_t^{i,0} + (1 - \delta) J_t^{\delta,i,0}|\mathcal{F}_s] \leq \delta \tilde{\mathcal{E}}_i[Z_t^{i,0}|\mathcal{F}_s] + (1 - \\ &\delta) \tilde{\mathcal{E}}_i[J_t^{\delta,i,0}|\mathcal{F}_s] \\ &\leq \delta Z_s^{i,0} + (1 - \delta) J_s^{\delta,i,0} = \delta Z_s^0 + (1 - \delta) J_s^{\delta,0} + H_s^i, \quad a.s., \end{aligned}$$

which shows that $\{\delta Z_t^0 + (1 - \delta) J_t^{\delta,i,0} + H_t^i\}_{t \in [0, T]}$ is an RCLL $\tilde{\mathcal{E}}_i$ -supermartingale.

Now we fix $t \in [0, T]$ and define $A \triangleq \{\tau_\delta(t) = t\} \in \mathcal{F}_t$. Using Proposition 82 (3), Lemma 80 (2) as well as applying Proposition 26 (2) and (5) to each $\tilde{\mathcal{E}}_i$, we obtain

$$\begin{aligned} \mathbf{1}_A J_t^{\delta,0} &= \mathbf{1}_A J_\delta(t) = \mathbf{1}_A \operatorname{esssup}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[Z_{\tau_\delta(t)}^0 + H_{t, \tau_\delta(t)}^i|\mathcal{F}_t] = \\ &\operatorname{esssup}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[\mathbf{1}_A (Z_{\tau_\delta(t)}^0 + H_{t, \tau_\delta(t)}^i)|\mathcal{F}_t] \\ &= \operatorname{esssup}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[\mathbf{1}_A Z_t^0|\mathcal{F}_t] = \mathbf{1}_A Z_t^0, \quad a.s. \end{aligned} \quad (8.56)$$

Then (4.17) and (4.11) imply that

$$\mathbf{1}_A (\delta Z_t^0 + (1 - \delta) J_t^{\delta,i,0}) = \mathbf{1}_A Z_t^0 = \mathbf{1}_A Z(t) \geq \mathbf{1}_A Y_t, \quad a.s. \quad (8.57)$$

Moreover, we see from the definition of $\tau_\delta(t)$ that for any $\omega \in A^c$

$$Y_s(\omega) < \delta Z_s^0(\omega) + (1 - \delta) (C_Y + 2 C_H), \quad \forall s \in [t, \tau_\delta(t)(\omega)) \quad (8.58)$$

Since both Z^0 and Y are right-continuous processes, (8.58) and (8.39) imply that

$$Y_t \leq \delta Z_t^0 + (1 - \delta)(C_Y + 2C_H) \leq \delta Z_t^0 + (1 - \delta)J_t^{\delta,0} \quad a.s. \langle \text{hbox} | \text{on } \mathcal{A}^c \rangle,$$

which in conjunction with (8.57) and Remark 73 (2) shows that the RCLL process $\delta Z^0 + (1 - \delta)J^{\delta,0}$ dominates Y , thus dominates Z^0 thanks to Proposition 4.18. It follows that $J^{\delta,0}$ also dominates Z^0 . Then for any $\nu \in \mathcal{S}_{0,T}$, Proposition 82 (3), Remark 73 (1) and (4.17) imply that $J_\delta(\nu) = J_\nu^{\delta,0} \geq Z_\nu^0 = Z(\nu)$, a.s., The reverse inequality comes from (4.19). This proves (8.55).

Next, we fix $\nu \in \mathcal{S}_{0,T}$ and set $\delta^n = \frac{n-1}{n}$, $n \in \mathbb{N}$. It is clear that the sequence $\{\tau_{\delta^n}(\nu)\}_{n \in \mathbb{N}}$ increasing a.s. to $\bar{\tau}(\nu)$. Since the family of processes $\{Y_\nu^i\}_{i \in \mathcal{I}}$ is " \mathcal{E} -uniformly-left-continuous", we can find a subsequence $\{\delta^{n-k}\}_{k \in \mathbb{N}}$ of $\{\delta^n\}_{n \in \mathbb{N}}$ such that

$$\lim_{k \rightarrow \infty} \operatorname{esssup}_{i \in \mathcal{I}} \left| \tilde{\mathcal{E}}_i \left[\langle \text{hbox} | \frac{n-k}{n-k-1} \rangle Y_{\tau_{\delta^{n-k}}(\nu)} + H_{\tau_{\delta^{n-k}}(\nu)}^i \middle| \mathcal{F}_\nu \right] - \tilde{\mathcal{E}}_i[Y_{\bar{\tau}(\nu)}^i | \mathcal{F}_\nu] \right| = 0, \quad a.s. \quad (8.59)$$

For any $i \in \mathcal{I}$ and $k \in \mathbb{N}$, Remark 53 (1) implies that $Y_{\tau_{\delta^{n-k}}(\nu)} \geq \delta^{n-k} Z_{\tau_{\delta^{n-k}}(\nu)}^0 + (1 - \delta^{n-k})(C_Y + 2C_H)$, a.s. Hence Proposition 26 (3) shows that

$$\begin{aligned} & \tilde{\mathcal{E}}_i[Z_{\tau_{\delta^{n-k}}(\nu)}^0 + H_{\nu, \tau_{\delta^{n-k}}(\nu)}^i | \mathcal{F}_\nu] + \langle \text{hbox} | \frac{1}{n_k - 1} (C_Y + 2C_H) \rangle \leq \tilde{\mathcal{E}}_i \left[\langle \text{hbox} | \frac{n-k}{n-k-1} \rangle Y_{\tau_{\delta^{n-k}}(\nu)} + H_{\tau_{\delta^{n-k}}(\nu)}^i \middle| \mathcal{F}_\nu \right] - H_\nu^i \\ &= \tilde{\mathcal{E}}_i \left[\langle \text{hbox} | \frac{n-k}{n-k-1} \rangle Y_{\tau_{\delta^{n-k}}(\nu)} + H_{\tau_{\delta^{n-k}}(\nu)}^i \middle| \mathcal{F}_\nu \right] - \tilde{\mathcal{E}}_i[Y_{\bar{\tau}(\nu)}^i | \mathcal{F}_\nu] + \tilde{\mathcal{E}}_i[Y_{\bar{\tau}(\nu)} + H_{\nu, \bar{\tau}(\nu)}^i | \mathcal{F}_\nu] \\ &\leq \operatorname{esssup}_{i \in \mathcal{I}} \left| \tilde{\mathcal{E}}_i \left[\langle \text{hbox} | \frac{n-k}{n-k-1} \rangle Y_{\tau_{\delta^{n-k}}(\nu)} + H_{\tau_{\delta^{n-k}}(\nu)}^i \middle| \mathcal{F}_\nu \right] - \tilde{\mathcal{E}}_i[Y_{\bar{\tau}(\nu)}^i | \mathcal{F}_\nu] \right| + \tilde{\mathcal{E}}_i[Y_{\bar{\tau}(\nu)} + H_{\nu, \bar{\tau}(\nu)}^i | \mathcal{F}_\nu] \quad (8.60) \\ &\leq \operatorname{esssup}_{i \in \mathcal{I}} \left| \tilde{\mathcal{E}}_i \left[\langle \text{hbox} | \frac{n-k}{n-k-1} \rangle Y_{\tau_{\delta^{n-k}}(\nu)} + H_{\tau_{\delta^{n-k}}(\nu)}^i \middle| \mathcal{F}_\nu \right] - \tilde{\mathcal{E}}_i[Y_{\bar{\tau}(\nu)}^i | \mathcal{F}_\nu] \right| + \operatorname{esssup}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[Y_{\bar{\tau}(\nu)} + H_{\nu, \bar{\tau}(\nu)}^i | \mathcal{F}_\nu], \quad a.s. \end{aligned}$$

Taking the esssup of the left-hand-side over \mathcal{I} , we see from (8.55) that

$$\begin{aligned} & \operatorname{esssup}_{i \in \mathcal{I}} \left| \tilde{\mathcal{E}}_i \left[\langle \text{hbox} | \frac{n-k}{n-k-1} \rangle Y_{\tau_{\delta^{n-k}}(\nu)} + H_{\tau_{\delta^{n-k}}(\nu)}^i \middle| \mathcal{F}_\nu \right] - \tilde{\mathcal{E}}_i[Y_{\bar{\tau}(\nu)}^i | \mathcal{F}_\nu] \right| + \operatorname{esssup}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[Y_{\bar{\tau}(\nu)} + H_{\nu, \bar{\tau}(\nu)}^i | \mathcal{F}_\nu] \\ &\geq J_{\delta^{n-k}}(\nu) + \langle \text{hbox} | \frac{1}{n_k - 1} (C_Y + 2C_H) \rangle = Z(\nu) + \langle \text{hbox} | \frac{1}{n_k - 1} (C_Y + 2C_H) \rangle, \quad a.s., \quad (8.61) \end{aligned}$$

As $k \rightarrow \infty$, (8.59), (4.11) and (4.15) imply that

$$Z(\nu) \leq \operatorname{esssup}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[Y_{\bar{\tau}(\nu)} + H_{\nu, \bar{\tau}(\nu)}^i | \mathcal{F}_\nu] \leq \operatorname{esssup}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[Z(\bar{\tau}(\nu)) + H_{\nu, \bar{\tau}(\nu)}^i | \mathcal{F}_\nu] \leq Z(\nu),$$

a. s.,

which shows that

$$Z(\nu) = \operatorname{esssup}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[Y_{\bar{\tau}(\nu)} + H_{\nu, \bar{\tau}(\nu)}^i | \mathcal{F}_\nu] = \operatorname{esssup}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[Z(\bar{\tau}(\nu)) + H_{\nu, \bar{\tau}(\nu)}^i | \mathcal{F}_\nu],$$

a. s. (8.62)

Now we fix $\rho \in \mathcal{S}_{\nu, \bar{\tau}(\nu)}$. For any $i \in \mathcal{I}$, Corollary 25 and (4.16) show that

$$\tilde{\mathcal{E}}_i[Z^i(\bar{\tau}(\nu)) | \mathcal{F}_\nu] = \tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_i[Z^i(\bar{\tau}(\nu)) | \mathcal{F}_\rho] | \mathcal{F}_\nu] \leq \tilde{\mathcal{E}}_i[Z^i(\rho) | \mathcal{F}_\nu], \quad a. s.$$

Then Proposition 26 (3) implies that

$$\tilde{\mathcal{E}}_i[Z(\bar{\tau}(\nu)) + H_{\nu, \bar{\tau}(\nu)}^i | \mathcal{F}_\nu] = \tilde{\mathcal{E}}_i[Z^i(\bar{\tau}(\nu)) | \mathcal{F}_\nu] - H_{\nu}^i \leq \tilde{\mathcal{E}}_i[Z^i(\rho) | \mathcal{F}_\nu] - H_{\nu}^i = \tilde{\mathcal{E}}_i[Z(\rho) + H_{\nu, \rho}^i | \mathcal{F}_\nu], \quad a. s.$$

Taking the essential supremum of both sides over \mathcal{I} , we can deduce from (4.15) that

$$\operatorname{esssup}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[Z(\bar{\tau}(\nu)) + H_{\nu, \bar{\tau}(\nu)}^i | \mathcal{F}_\nu] \leq \operatorname{esssup}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[Z(\rho) + H_{\nu, \rho}^i | \mathcal{F}_\nu] \leq Z(\nu), \quad a. s.,$$

which together with (8.62) proves (4.22).

Finally, we will prove that $\bar{\tau}(\nu) = \tau_1(\nu)$. For any $i \in \mathcal{I}$ and $k \in \mathbb{N}$, (4.17), (4.15), Proposition 26 (3) as well as Corollary 25 imply that

$$\begin{aligned} & \tilde{\mathcal{E}}_i[Z_{\tau_{\delta}^{n-k}(\nu)}^0 + H_{\nu, \tau_{\delta}^{n-k}(\nu)}^i | \mathcal{F}_\nu] = \tilde{\mathcal{E}}_i[Z(\tau_{\delta}^{n-k}(\nu)) + H_{\nu, \tau_{\delta}^{n-k}(\nu)}^i | \mathcal{F}_\nu] \\ & \geq \tilde{\mathcal{E}}_i \left[\operatorname{esssup}_{j \in \mathcal{I}} \tilde{\mathcal{E}}_j[Z(\bar{\tau}(\nu)) + H_{\tau_{\delta}^{n-k}(\nu), \bar{\tau}(\nu)}^j | \mathcal{F}_{\tau_{\delta}^{n-k}(\nu)}] + H_{\nu, \tau_{\delta}^{n-k}(\nu)}^i \middle| \mathcal{F}_\nu \right] \\ & \geq \tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_i[Z(\bar{\tau}(\nu)) + H_{\tau_{\delta}^{n-k}(\nu), \bar{\tau}(\nu)}^i | \mathcal{F}_{\tau_{\delta}^{n-k}(\nu)}] + H_{\nu, \tau_{\delta}^{n-k}(\nu)}^i | \mathcal{F}_\nu] \\ & = \tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_i[Z(\bar{\tau}(\nu)) + H_{\nu, \bar{\tau}(\nu)}^i | \mathcal{F}_{\tau_{\delta}^{n-k}(\nu)}] | \mathcal{F}_\nu] = \tilde{\mathcal{E}}_i[Z(\bar{\tau}(\nu)) + H_{\nu, \bar{\tau}(\nu)}^i | \mathcal{F}_\nu], \quad a. s., \end{aligned}$$

which together with (8.60) shows that

$$\begin{aligned} & \operatorname{esssup}_{i \in \mathcal{I}} \left| \tilde{\mathcal{E}}_i \left[\langle \text{hbox} \mid \frac{n-k}{n-k-1} Y_{\tau_{\delta}^{n-k}(\nu)} \neg + \neg H_{\tau_{\delta}^{n-k}(\nu)}^i \middle| \mathcal{F}_\nu \right] \neg \neg \tilde{\mathcal{E}}_i[Y_{\bar{\tau}(\nu)}^i | \mathcal{F}_\nu] \right| + \tilde{\mathcal{E}}_i[Y_{\bar{\tau}(\nu)} \neg + \neg H_{\nu, \bar{\tau}(\nu)}^i | \mathcal{F}_\nu] \neg \\ & \geq \tilde{\mathcal{E}}_i[Z(\bar{\tau}(\nu)) \neg + \neg H_{\nu, \bar{\tau}(\nu)}^i | \mathcal{F}_\nu] + \langle \text{hbox} \mid \frac{1}{n_k-1} (C_Y + 2 C_H) \rangle, \quad a. s. \end{aligned}$$

As $k \rightarrow \infty$, (8.59) implies that

$$\tilde{\mathcal{E}}_i[Y_{\bar{\tau}(\nu)} + H_{\nu, \bar{\tau}(\nu)}^i | \mathcal{F}_\nu] \geq \tilde{\mathcal{E}}_i[Z(\bar{\tau}(\nu)) + H_{\nu, \bar{\tau}(\nu)}^i | \mathcal{F}_\nu], \quad a. s. \quad (8.63)$$

The reverse inequality follows easily from (4.11), thus (8.63) is in fact an equality. Then the second part of Proposition 26 (1) and (4.17) imply that

$$Y_{\bar{\tau}(\nu)} = Z(\bar{\tau}(\nu)) = Z_{\bar{\tau}(\nu)}^0, \quad a. s.,$$

which shows that $\inf\{t \in [\nu, T]: Z_t^0 = Y_t\} \leq \bar{\tau}(\nu)$, a.s. For any $\delta \in (0, 1)$, since $\{t \in [\nu, T]: Z_t^0 = Y_t\} \subset \{t \in [\nu, T]: Y_t \geq \delta Z_t^0 + (1 - \delta)(C_Y + 2C_H)\}$, one can deduce that

$$\bar{\tau}(\nu) \geq \inf\{t \in [\nu, T]: Z_t^0 = Y_t\} \geq \inf\{t \in [\nu, T]: Y_t \geq \delta Z_t^0 + (1 - \delta)(C_Y + 2C_H)\} \wedge T = \tau_\delta(\nu), \quad a.s.$$

Letting $\delta \rightarrow 1$ yields that

$$\bar{\tau}(\nu) \geq \inf\{t \in [\nu, T]: Z_t^0 = Y_t\} \geq \lim_{\delta \rightarrow 1} \tau_\delta(\nu) = \bar{\tau}(\nu), \quad a.s.,$$

which implies that $\bar{\tau}(\nu) = \inf\{t \in [\nu, T]: Z_t^0 = Y_t\}$, a.s. \square

8.4 Proofs of Section 5

Definition 83. A family $\{\xi_i\}_{i \in \mathcal{I}} \subset L^0(\mathcal{F}_T)$ is said to be directed downwards if for any $i, j \in \mathcal{I}$, there exists a $k \in \mathcal{I}$ such that $\xi_k \leq \xi_i \wedge \xi_j$, a.s.

Proof of Lemma 61: In light of (, ?, Proposition VI-11-1), it suffices to show that the family $\{R^i(\nu)\}_{i \in \mathcal{I}}$ is directed downwards. To see this, we define the event $A \triangleq \{R^i(\nu) \geq R^j(\nu)\}$ and the stopping times

$$\rho \triangleq \tau^j(\nu) \mathbf{1}_A + \tau^i(\nu) \mathbf{1}_{A^c} \in \mathcal{S}_{\nu, T} \quad \langle \text{hbox|a n d} \rangle \quad \nu(A) \triangleq \nu \mathbf{1}_A + T \mathbf{1}_{A^c} \in \mathcal{S}_{\nu, T}.$$

By Definition 7.13, there exists a $k = k(i, j, \nu(A)) \in \mathcal{I}$ such that $\tilde{\mathcal{E}}_k = \mathcal{E}_{i,j}^{\nu(\neg A)}$. Similar to (8.25) it holds for any $\xi \in Dom(\mathcal{E})$ that

$$\tilde{\mathcal{E}}_k[\xi | \mathcal{F}_\nu] = \mathbf{1}_A \tilde{\mathcal{E}}_j[\xi | \mathcal{F}_\nu] + \mathbf{1}_{A^c} \tilde{\mathcal{E}}_i[\xi | \mathcal{F}_\nu], \quad a.s. \quad (8.64)$$

Moreover, (4.5) implies that

$$\begin{aligned} & H_{\nu, \rho}^k \neg = \neg H_{\nu(A) \wedge \nu, \nu(A) \wedge \rho}^i \neg + \neg H_{\nu(A) \vee \nu, \nu(A) \vee \rho}^j \neg \\ & = \neg \mathbf{1}_{A^c} H_{\nu, \tau^i(\nu)}^i \neg + \neg \mathbf{1}_A H_{\nu, \tau^j(\nu)}^j \neg, \quad a.s. \\ \langle \text{error|compound hbox} \rangle & H_{\nu, \tau^k(\nu)}^k \neg = \neg H_{\nu(A) \wedge \nu, \nu(A) \wedge \tau^k(\nu)}^i \neg + \neg \\ & H_{\nu(A) \vee \nu, \nu(A) \vee \tau^k(\nu)}^j \neg = \neg \mathbf{1}_{A^c} H_{\nu, \tau^k(\nu)}^i \neg + \neg \\ & \mathbf{1}_A H_{\nu, \tau^k(\nu)}^j \neg, \quad a.s. \end{aligned}$$

Using (8.64) twice and applying Proposition 26 (2) to $\tilde{\mathcal{E}}_i$ and $\tilde{\mathcal{E}}_j$, we can deduce from (5.7) that

$$\begin{aligned} R^k(\nu) & \geq \tilde{\mathcal{E}}_k[Y_\rho + H_{\nu, \rho}^k | \mathcal{F}_\nu] = \mathbf{1}_A \tilde{\mathcal{E}}_j[Y_\rho + H_{\nu, \rho}^k | \mathcal{F}_\nu] + \mathbf{1}_{A^c} \tilde{\mathcal{E}}_i[Y_\rho + H_{\nu, \rho}^k | \mathcal{F}_\nu] \\ & = \tilde{\mathcal{E}}_j[\mathbf{1}_A Y_{\tau^j(\nu)} + \mathbf{1}_A H_{\nu, \tau^j(\nu)}^j | \mathcal{F}_\nu] + \tilde{\mathcal{E}}_i[\mathbf{1}_{A^c} Y_{\tau^i(\nu)} + \mathbf{1}_{A^c} H_{\nu, \tau^i(\nu)}^i | \mathcal{F}_\nu] \\ & = \mathbf{1}_A \tilde{\mathcal{E}}_j[Y_{\tau^j(\nu)} + H_{\nu, \tau^j(\nu)}^j | \mathcal{F}_\nu] + \mathbf{1}_{A^c} \tilde{\mathcal{E}}_i[Y_{\tau^i(\nu)} + H_{\nu, \tau^i(\nu)}^i | \mathcal{F}_\nu] = \\ & \mathbf{1}_A R^j(\nu) + \mathbf{1}_{A^c} R^i(\nu) \\ & \geq \mathbf{1}_A \tilde{\mathcal{E}}_j[Y_{\tau^k(\nu)} + H_{\nu, \tau^k(\nu)}^j | \mathcal{F}_\nu] + \mathbf{1}_{A^c} \tilde{\mathcal{E}}_i[Y_{\tau^k(\nu)} + H_{\nu, \tau^k(\nu)}^i | \mathcal{F}_\nu] \\ & = \tilde{\mathcal{E}}_j[\mathbf{1}_A Y_{\tau^k(\nu)} + \mathbf{1}_A H_{\nu, \tau^k(\nu)}^j | \mathcal{F}_\nu] + \tilde{\mathcal{E}}_i[\mathbf{1}_{A^c} Y_{\tau^k(\nu)} + \mathbf{1}_{A^c} H_{\nu, \tau^k(\nu)}^i | \mathcal{F}_\nu] \\ & = \mathbf{1}_A \tilde{\mathcal{E}}_j[Y_{\tau^k(\nu)} + H_{\nu, \tau^k(\nu)}^j | \mathcal{F}_\nu] + \mathbf{1}_{A^c} \tilde{\mathcal{E}}_i[Y_{\tau^k(\nu)} + H_{\nu, \tau^k(\nu)}^i | \mathcal{F}_\nu] \\ & = \tilde{\mathcal{E}}_k[Y_{\tau^k(\nu)} + H_{\nu, \tau^k(\nu)}^k | \mathcal{F}_\nu] = R^k(\nu), \quad a.s., \end{aligned}$$

which shows that $R^k(\nu) = \mathbf{1}_A R^j(\nu) + \mathbf{1}_{A^c} R^i(\nu) = R^i(\nu) \wedge R^j(\nu)$, a.s. In light of the basic properties of the essential infimum (e.g., (5.2), Proposition VI-11-1)), we can find a sequence $\{i_n\}_{n \in \mathbb{N}}$ in \mathcal{I} such that (5.8) holds. \square

Proof of Lemma 62: As in the proof of Lemma 61, it suffices to show that the family $\{\tau^i(\nu)\}_{i \in \mathcal{I}}$ is directed downwards. To see this, we define the stopping time $\sigma \triangleq \tau^i(\nu) \wedge \tau^j(\nu) \in \mathcal{S}_{\nu, T}$, the event $A \triangleq \{R_{\sigma}^{i,0} \geq R_{\sigma}^{j,0}\} \in \mathcal{F}_{\sigma}$ as well as the stopping time $\sigma(A) \triangleq \sigma \mathbf{1}_A + T \mathbf{1}_{A^c} \in \mathcal{S}_{\sigma, T}$. By Definition 7.13, there exists a $k = k(i, j, \sigma(A)) \in \mathcal{I}$ such that $\tilde{\mathcal{E}}_k = \mathcal{E}_{i,j}^{\sigma(\neg A)}$. Fix $t \in [0, T]$, similar to (8.25), it holds for any $\xi \in \mathcal{D}om(\mathcal{E})$ that

$$\tilde{\mathcal{E}}_k[\xi | \mathcal{F}_{\sigma \vee t}] = \mathbf{1}_A \tilde{\mathcal{E}}_j[\xi | \mathcal{F}_{\sigma \vee t}] + \mathbf{1}_{A^c} \tilde{\mathcal{E}}_i[\xi | \mathcal{F}_{\sigma \vee t}], \quad a.s. \quad (8.65)$$

Moreover, we can deduce from (4.5) that for any $\rho \in \mathcal{S}_{\sigma \vee t, T}$

$$H_{\sigma \vee t, \rho}^k \neg \neg = \neg H_{\sigma(A) \wedge (\sigma \vee t), \sigma(A) \wedge \rho}^i \neg \neg + \neg H_{\sigma(A) \vee (\sigma \vee t), \sigma(A) \vee \rho}^j \neg \neg = \neg \mathbf{1}_A H_{\sigma \vee t, \rho}^i \neg \neg + \neg \mathbf{1}_A H_{\sigma \vee t, \rho}^j \neg \neg, \quad a.s.,$$

which together with (8.65) and Proposition 26 (2) imply that

$$\begin{aligned} \tilde{\mathcal{E}}_k[Y_{\rho} \neg \neg + \neg H_{\sigma \vee t, \rho}^k | \mathcal{F}_{\sigma \vee t}] &= \mathbf{1}_A \tilde{\mathcal{E}}_j[Y_{\rho} \neg \neg + \neg H_{\sigma \vee t, \rho}^k | \mathcal{F}_{\sigma \vee t}] + \mathbf{1}_{A^c} \tilde{\mathcal{E}}_i[Y_{\rho} \neg \neg + \neg H_{\sigma \vee t, \rho}^k | \mathcal{F}_{\sigma \vee t}] \\ &= \tilde{\mathcal{E}}_j[\mathbf{1}_A Y_{\rho} \neg \neg + \mathbf{1}_A H_{\sigma \vee t, \rho}^j | \mathcal{F}_{\sigma \vee t}] + \tilde{\mathcal{E}}_i[\mathbf{1}_{A^c} Y_{\rho} \neg \neg + \mathbf{1}_{A^c} \neg H_{\sigma \vee t, \rho}^i | \mathcal{F}_{\sigma \vee t}] \\ &= \mathbf{1}_A \tilde{\mathcal{E}}_j[Y_{\rho} \neg \neg + \neg H_{\sigma \vee t, \rho}^j | \mathcal{F}_{\sigma \vee t}] + \mathbf{1}_{A^c} \tilde{\mathcal{E}}_i[Y_{\rho} \neg \neg + \neg H_{\sigma \vee t, \rho}^i | \mathcal{F}_{\sigma \vee t}], \quad a.s. \end{aligned}$$

Then applying Proposition 26 (3), Lemma 80 (2) as well as (5.6), we obtain

$$\begin{aligned} R_{\sigma \vee t}^{k,0} &= R^k(\sigma \vee t) = \text{esssup}_{\rho \in \mathcal{S}_{\sigma \vee t, T}} \tilde{\mathcal{E}}_k[Y_{\rho} + H_{\sigma \vee t, \rho}^k | \mathcal{F}_{\sigma \vee t}] \\ &= \mathbf{1}_A \text{esssup}_{\rho \in \mathcal{S}_{\sigma \vee t, T}} \tilde{\mathcal{E}}_j[Y_{\rho} \neg \neg + \neg H_{\sigma \vee t, \rho}^j | \mathcal{F}_{\sigma \vee t}] + \mathbf{1}_{A^c} \text{esssup}_{\rho \in \mathcal{S}_{\sigma \vee t, T}} \tilde{\mathcal{E}}_i[Y_{\rho} \neg \neg + \neg H_{\sigma \vee t, \rho}^i | \mathcal{F}_{\sigma \vee t}] \\ &= \mathbf{1}_A R^j(\sigma \vee t) + \mathbf{1}_{A^c} R^i(\sigma \vee t) = \mathbf{1}_A R_{\sigma \vee t}^{j,0} + \mathbf{1}_{A^c} R_{\sigma \vee t}^{i,0}, \quad a.s. \end{aligned} \quad (8.66)$$

Since $R^{i,0}$, $R^{j,0}$ and $R^{k,0}$ are all RCLL processes, it holds except on a null set N that

$$R_{\sigma \vee t}^{k,0} = \mathbf{1}_A R_{\sigma \vee t}^{j,0} + \mathbf{1}_{A^c} R_{\sigma \vee t}^{i,0}, \quad \forall t \in [0, T],$$

which further implies that

$$\begin{aligned} \tau^k(\nu) \neg \neg &= \neg \neg \inf \{t \in [\nu, T]: R_t^{k,0} = Y_t\} \leq \inf \{t \in [\sigma, T]: R_t^{k,0} = Y_t\} \\ &= \neg \neg \mathbf{1}_A \inf \{t \in [\sigma, T]: R_t^{j,0} = Y_t\} + \mathbf{1}_{A^c} \inf \{t \in [\sigma, T]: R_t^{i,0} = Y_t\}, \quad a.s. \end{aligned} \quad (8.67)$$

Since $R_{\tau^i(\nu)}^{i,0} = Y_{\tau^i(\nu)}$, $R_{\tau^j(\nu)}^{j,0} = Y_{\tau^j(\nu)}$, a.s. and since $\sigma = \tau^i(\nu) \wedge \tau^j(\nu)$, it holds a.s. that Y_σ is equal either to $R_\sigma^{i,0}$ or to $R_\sigma^{j,0}$. Then the definition of the set A shows that $R_\sigma^{j,0} = Y_\sigma$ a.s. on A and that $R_\sigma^{i,0} = Y_\sigma$ a.s. on A^c , both of which further implies that

$$\mathbf{1}_A \inf \{t \in [\sigma, T]: R_t^{j,0} = Y_t\} = \sigma \mathbf{1}_A \quad \langle \text{hbox} \mid a \ n \ d \rangle \quad \mathbf{1}_{A^c} \inf \{t \in [\sigma, T]: R_t^{i,0} = Y_t\} = \sigma \mathbf{1}_{A^c}, \quad a. s.$$

Hence, we see from (8.66) that $\tau^k(\nu) \leq \sigma = \tau^i(\nu) \wedge \tau^j(\nu)$, a.s. Thanks to the basic properties of the essential infimum (e.g., (, ?, Proposition VI-11-1)), we can find a sequence $\{i_n\}_{n \in \mathbb{N}}$ in \mathcal{I} such that

$$\tau(\nu) = \operatorname{ess\,inf}_{i \in \mathcal{I}} \tau^i(\nu) = \lim_{n \rightarrow \infty} \neg \neg \downarrow \tau^{i_n}(\nu), \quad a. s.$$

The limit $\lim_{n \rightarrow \infty} \neg \neg \downarrow \tau^{i_n}(\nu)$ is also a stopping time, thus we have $\tau(\nu) \in \mathcal{S}_{\nu, T}$. \square

Proof of Theorem 68: In light of Lemma 62, there exists a sequence $\{j_n\}_{n \in \mathbb{N}}$ in \mathcal{I} such that

$$\tau(\nu) = \lim_{n \rightarrow \infty} \neg \neg \downarrow \tau^{j_n}(\nu), \quad a. s.$$

Since the family of processes $\{Y^i\}_{i \in \mathcal{I}}$ is “ \mathcal{E} -uniformly-right-continuous”, we can find a subsequence of $\{j_n\}_{n \in \mathbb{N}}$ (we still denote it by $\{j_n\}_{n \in \mathbb{N}}$) such that

$$\lim_{n \rightarrow \infty} \operatorname{ess\,sup}_{i \in \mathcal{I}} \left| \tilde{\mathcal{E}}_i[Y_{\tau^{j_n}(\nu)}^i | \mathcal{F}_{\tau(\nu)}] - Y_{\tau(\nu)}^i \right| = 0, \quad a. s. \quad (8.68)$$

Fix $i \in \mathcal{I}$ and $n \in \mathbb{N}$, we know from Definition 7.13 that there exists a $k_n = k(i, j_n, \tau(\nu)) \in \mathcal{I}$ such that $\tilde{\mathcal{E}}_{k_n} = \mathcal{E}_{i, j_n}^{\tau(\nu)}$.

$$\tilde{\mathcal{E}}_{k_n} = \mathcal{E}_{i, j_n}^{\tau(\nu)} \quad \langle \text{hbox} \mid a \ n \ d \rangle \quad h_t^{k_n} = \mathbf{1}_{\{t < \tau(\nu)\}} h_t^i + \mathbf{1}_{\{t \geq \tau(\nu)\}} h_t^{j_n}, \quad a. s. \quad (8.69)$$

(fi) For any $t \in [0, T]$, Lemma 80 implies that $R_{\tau(\nu) \vee t}^{k_n, 0} = R_{\tau(\nu) \vee t}^{j_n, 0}$, a.s. Since $R^{k_n, 0}$ and $R^{j_n, 0}$ are both RCLL processes, it holds except on a null set N that

$$R_{\tau(\nu) \vee t}^{k_n, 0} = R_{\tau(\nu) \vee t}^{j_n, 0}, \quad \forall t \in [0, T],$$

which together with the fact that $\tau(\nu) \leq \tau^{k_n}(\nu) \wedge \tau^{j_n}(\nu)$, a.s. implies that

$$\begin{aligned} \tau^{k_n}(\nu) &= \inf \{t \in [\nu, T]: R_t^{k_n, 0} = Y_t\} = \inf \{t \in [\tau(\nu), T]: R_t^{k_n, 0} = Y_t\} \\ &= \inf \{t \in [\tau(\nu), T]: R_t^{j_n, 0} = Y_t\} = \inf \{t \in [\nu, T]: R_t^{j_n, 0} = Y_t\} = \tau^{j_n}(\nu), \\ &\quad a. s. \end{aligned} \quad (8.70)$$

Then (7), (8.69) and (75) show that

$$\begin{aligned}
R^{k_n}(\nu) + H_\nu^i &= R^{k_n}(\nu) + H_\nu^{k_n} = \tilde{\mathcal{E}}_{k_n}[Y_{\tau^{k_n}(\nu)} + H_{\nu, \tau^{k_n}(\nu)}^{k_n} | \mathcal{F}_\nu] + H_\nu^{k_n} \\
\neg\neg\neg &= \neg\neg\neg \tilde{\mathcal{E}}_{k_n}[Y_{\tau^{k_n}(\nu)} | \mathcal{F}_\nu] = \mathcal{E}_{i, j_n}^{(\nu)}[Y_{\tau^{j_n}(\nu)} | \mathcal{F}_\nu] = \tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_{j_n}[Y_{\tau^{j_n}(\nu)} | \mathcal{F}_{\mathcal{I}(\nu)}] | \mathcal{F}_\nu] \\
\neg\neg\neg &= \neg\neg\neg \tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_{j_n} \neg[Y_{\tau^{j_n}(\nu)} \neg + \neg H_{\mathcal{I}(\nu), \tau^{j_n}(\nu)}^{j_n} \neg + \neg H_{\mathcal{I}(\nu)}^i | \mathcal{F}_{\mathcal{I}(\nu)}] | \mathcal{F}_\nu] \neg = \\
&\quad \tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_{j_n} \neg[Y_{\tau^{j_n}(\nu)} | \mathcal{F}_{\mathcal{I}(\nu)}] - H_{\mathcal{I}(\nu)}^{j_n} \neg + \neg H_{\mathcal{I}(\nu)}^i | \mathcal{F}_\nu] \\
\neg\neg\neg &\leq \neg\neg\neg \tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_{j_n}[Y_{\tau^{j_n}(\nu)} | \mathcal{F}_{\mathcal{I}(\nu)}] \neg - \neg Y_{\mathcal{I}(\nu)}^{j_n} | \neg + \neg Y_{\mathcal{I}(\nu)}^i | \mathcal{F}_\nu] \leq \\
&\quad \tilde{\mathcal{E}}_i \left[\operatorname{esssup}_{l \in \mathcal{I}} \left| \tilde{\mathcal{E}}_l[Y_{\tau^{j_n}(\nu)}^l | \mathcal{F}_{\mathcal{I}(\nu)}] \neg - \neg Y_{\mathcal{I}(\nu)}^l | \neg + \neg Y_{\mathcal{I}(\nu)}^i | \mathcal{F}_\nu \right| \right], \\
&\quad a. s. \tag{8.71}
\end{aligned}$$

For any $l \in \mathcal{I}$, Proposition 26 (3), (7), (7) and (5.3) imply that

$$\begin{aligned}
|\tilde{\mathcal{E}}_l[Y_{\tau^{j_n}(\nu)}^l | \mathcal{F}_{\mathcal{I}(\nu)}] - Y_{\mathcal{I}(\nu)}^l| &= |\tilde{\mathcal{E}}_l[Y_{\tau^{j_n}(\nu)} + H_{\mathcal{I}(\nu), \tau^{j_n}(\nu)}^l - C | \mathcal{F}_{\mathcal{I}(\nu)}] - (Y_{\mathcal{I}(\nu)} - C_Y) + C_H| \\
&\leq |\tilde{\mathcal{E}}_l[Y_{\tau^{j_n}(\nu)} + H_{\mathcal{I}(\nu), \tau^{j_n}(\nu)}^l - C | \mathcal{F}_{\mathcal{I}(\nu)}]| + |Y_{\mathcal{I}(\nu)} - C_Y| + |C_H| \\
&= \tilde{\mathcal{E}}_l[Y_{\tau^{j_n}(\nu)} + H_{\mathcal{I}(\nu), \tau^{j_n}(\nu)}^l - C | \mathcal{F}_{\mathcal{I}(\nu)}] + (Y_{\mathcal{I}(\nu)} - C_Y) - C_H \\
&= \tilde{\mathcal{E}}_l[Y_{\tau^{j_n}(\nu)}^l + H_{\mathcal{I}(\nu), \tau^{j_n}(\nu)}^l | \mathcal{F}_{\mathcal{I}(\nu)}] + Y_{\mathcal{I}(\nu)} - 2C \leq \\
&\quad 2R^l(\mathcal{I}(\nu)) - 2C, \quad a. s.
\end{aligned}$$

Taking the essential supremum over $l \in \mathcal{I}$, we can deduce from (7) and (5.3) that

$$\begin{aligned}
C &\leq \operatorname{esssup}_{l \in \mathcal{I}} \left| \tilde{\mathcal{E}}_l[Y_{\tau^{j_n}(\nu)}^l | \mathcal{F}_{\mathcal{I}(\nu)}] - Y_{\mathcal{I}(\nu)}^l \right| + Y_{\mathcal{I}(\nu)}^i \leq 3R^l(\mathcal{I}(\nu)) - 2C + H_{\mathcal{I}(\nu)}^i, \\
&\quad a. s.,
\end{aligned}$$

where $3R^l(\mathcal{I}(\nu)) - 2C_* + H_{\mathcal{I}(\nu)}^i \in \operatorname{Dom}(\mathcal{E})$ thanks to Proposition 82 (1), (S1') and (D2). Applying the Dominated Convergence Theorem (Proposition 29) and Proposition 26 (3), we can deduce from (8.70) and (8.68) that

$$\begin{aligned}
\bar{V}(\nu) &= \operatorname{essinf}_{j \in \mathcal{I}} R^j(\nu) \leq \liminf_{n \rightarrow \infty} R^{k_n}(\nu) \leq \neg \lim_{n \rightarrow \infty} \tilde{\mathcal{E}}_i \left[\operatorname{esssup}_{l \in \mathcal{I}} \left| \tilde{\mathcal{E}}_l[Y_{\tau^{j_n}(\nu)}^l | \mathcal{F}_{\mathcal{I}(\nu)}] \neg - \neg Y_{\mathcal{I}(\nu)}^l | \neg + \neg Y_{\mathcal{I}(\nu)}^i | \mathcal{F}_\nu \right| \neg - \neg H_\nu^i \right] \\
&= \tilde{\mathcal{E}}_i[Y_{\mathcal{I}(\nu)}^i | \mathcal{F}_\nu] \neg - \neg H_\nu^i = \tilde{\mathcal{E}}_i[Y_{\mathcal{I}(\nu)} \neg + \neg H_{\nu, \mathcal{I}(\nu)}^i | \mathcal{F}_\nu], \quad a. s.
\end{aligned}$$

Taking the essential infimum of the right-hand-side over $i \in \mathcal{I}$ yields that

$$\begin{aligned}
\bar{V}(\nu) &\leq \operatorname{essinf}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[Y_{\mathcal{I}(\nu)} + \neg H_{\nu, \mathcal{I}(\nu)}^i | \mathcal{F}_\nu] \leq \operatorname{esssup}_{\rho \in \mathcal{S}_{\nu, \mathcal{I}}} \left(\operatorname{essinf}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[Y_\rho + \neg H_{\nu, \rho}^i | \mathcal{F}_\nu] \right) = \underline{V}(\nu) \leq \bar{V}(\nu), \quad a. s.
\end{aligned}$$

Hence, we have

$$V(\nu) = \operatorname{essinf}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[Y_{\mathcal{I}(\nu)} + \neg H_{\nu, \mathcal{I}(\nu)}^i | \mathcal{F}_\nu] = \bar{V}(\nu) = \operatorname{essinf}_{i \in \mathcal{I}} R^i(\nu) \geq Y_\nu, \quad a. s.,$$

where the last inequality is due to (5.3). \square

Proof of Proposition 68: By Lemma 62, there exists a sequence $\{i_n\}_{n \in \mathbb{N}}$ in \mathcal{I} such that

$$\sigma \stackrel{\Delta}{=} \tau(\nu) = \lim_{n \rightarrow \infty} \neg \neg \downarrow \tau^{i_n}(\nu), \quad a. s.$$

For any $n \in \mathbb{N}$, since $\sigma \leq \tau^{i_n}(\nu)$, a.s., we have

$$\tau^{i_n}(\nu) = \inf \{t \in [\nu, T]: R_t^{i_n, 0} = Y_t\} = \inf \{t \in [\sigma, T]: R_t^{i_n, 0} = Y_t\} = \tau^{i_n}(\sigma), \quad a. s.$$

Then (5.9) and (5.7) imply that

$$\begin{aligned} V(\sigma) &= \bar{V}(\sigma) \leq R^{i_n}(\sigma) = \tilde{\mathcal{E}}_{i_n}[Y_{\tau^{i_n}(\sigma)} + H_{\sigma, \tau^{i_n}(\sigma)}^{i_n} | \mathcal{F}_\sigma] = \tilde{\mathcal{E}}_{i_n}[Y_{\tau^{i_n}(\nu)} + H_{\sigma, \tau^{i_n}(\nu)}^{i_n} | \mathcal{F}_\sigma] \\ &= \tilde{\mathcal{E}}_{i_n}[Y_{\tau^{i_n}(\nu)} | \mathcal{F}_\sigma] - H_\sigma^{i_n} = \tilde{\mathcal{E}}_{i_n}[Y_{\tau^{i_n}(\nu)} | \mathcal{F}_\sigma] - Y_\sigma^{i_n} + Y_\sigma \leq \\ &\quad \operatorname{esssup}_{i \in \mathcal{I}} \left| \tilde{\mathcal{E}}_i[Y_{\tau^{i_n}(\nu)} | \mathcal{F}_\sigma] - Y_\sigma^i \right| + Y_\sigma, \quad a. s. \end{aligned}$$

As $n \rightarrow \infty$, the “ \mathcal{E} -uniform-right-continuity” of $\{Y^i\}_{i \in \mathcal{I}}$ implies that $V(\sigma) \leq Y_\sigma$, a.s., while the reverse inequality is obvious from (5.9). \square

Proof of Proposition 67: In light of Lemma 61 and (5.9), there exists a sequence $\{j_n\}_{n \in \mathbb{N}}$ in \mathcal{I} such that

$$V(\nu) = \bar{V}(\nu) = \lim_{n \rightarrow \infty} \neg \neg \downarrow R^{j_n}(\nu), \quad a. s.$$

For any $n \in \mathbb{N}$, Definition 7.13 assures a $k_n = k(i, j_n, \nu) \in \mathcal{I}$ such that $\tilde{\mathcal{E}}_{k_n} = \mathcal{E}_{i, j_n}^\nu$.

$$\tilde{\mathcal{E}}_{k_n} = \mathcal{E}_{i, j_n}^\nu \quad \langle \text{hbox} | a \text{ n } d \rangle \quad h_t^{k_n} = \mathbf{1}_{\{t < \nu\}} h_t^i + \mathbf{1}_{\{t \geq \nu\}} h_t^{j_n}, \quad a. s. \quad (8.72)$$

$\langle \text{fi} \rangle$ Applying Proposition 26 (5) to $\tilde{\mathcal{E}}_i$, we can deduce from (75) and (5.5) that

$$\begin{aligned} \tilde{\mathcal{E}}_{k_n}[V(\rho) + \neg H_{\nu, \rho}^{j_n} | \mathcal{F}_\nu] &\leq \tilde{\mathcal{E}}_{k_n}[R^{j_n}(\rho) + H_{\nu, \rho}^{j_n} | \mathcal{F}_\nu] = \mathcal{E}_{i, j_n}^\nu[R^{j_n}(\rho) + H_{\nu, \rho}^{j_n} | \mathcal{F}_\nu] \\ &= \tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_{j_n}[R^{j_n}(\rho) + H_{\nu, \rho}^{j_n} | \mathcal{F}_\nu] | \mathcal{F}_\nu] \\ &\neg \neg \neg = \neg \neg \neg \tilde{\mathcal{E}}_{j_n}[R^{j_n}(\rho) + H_{\nu, \rho}^{j_n} | \mathcal{F}_\nu] \leq R^{j_n}(\nu), \quad a. s. \end{aligned}$$

Then Proposition 26 (3) and (?) imply that

$$\begin{aligned} \operatorname{essinf}_{k \in \mathcal{I}} \tilde{\mathcal{E}}_k[V^k(\rho) | \mathcal{F}_\nu] \neg \neg \neg &\leq \neg \neg \neg \tilde{\mathcal{E}}_{k_n}[V^{k_n}(\rho) | \mathcal{F}_\nu] = \tilde{\mathcal{E}}_{k_n}[V(\rho) + \neg H_{\nu, \rho}^{j_n} | \mathcal{F}_\nu] + \neg \\ &H_\nu^i \leq R^{j_n}(\nu) + H_\nu^i, \quad a. s. \end{aligned}$$

As $n \rightarrow \infty$, (5.10) follows:

$$\operatorname{essinf}_{k \in \mathcal{I}} \tilde{\mathcal{E}}_k[V^k(\rho) | \mathcal{F}_\nu] \leq \lim_{n \rightarrow \infty} \neg \neg \downarrow R^{j_n}(\nu) + H_\nu^i = V(\nu) + H_\nu^i = V^i(\nu), \quad a. s.$$

Now we assume that $\nu \leq \rho \leq \tau(\nu)$, a.s. Applying Lemma 61 and (5.9) once again, we can find another sequence $\{j'_n\}_{n \in \mathbb{N}}$ in \mathcal{I} such that

$$V(\rho) = \bar{V}(\rho) = \lim_{n \rightarrow \infty} \neg \neg \downarrow R^{j'_n}(\rho), \quad a. s.$$

For any $n \in \mathbb{N}$, Definition 7.13 assures a $k'_n = k(i, j'_n, \rho) \in \mathcal{I}$ such that $\tilde{\mathcal{E}}_{k'_n} = \mathcal{E}_{i, j'_n}^\rho$. Since $\rho \leq \tau(\nu) \leq \tau^{k'_n}(\nu)$, a.s., using (5.7) with $i = k'_n$ and applying Proposition 26 (5) to $\tilde{\mathcal{E}}_{j'_n}$, we can deduce from (4.5), (75) as well as Lemma 80 that

$$\begin{aligned} V^i(\nu) \neg \neg &= \neg \neg V(\nu) + H_\nu^i = V(\nu) + H_\nu^{k'_n} \leq R^{k'_n}(\nu) + H_\nu^{k'_n} = \tilde{\mathcal{E}}_{k'_n}[R^{k'_n}(\rho) + \\ &H_\rho^{k'_n} | \mathcal{F}_\nu] = \mathcal{E}_{i, j'_n}^\rho[R^{k'_n}(\rho) + H_\rho^{k'_n} | \mathcal{F}_\nu] \\ \neg \neg &= \neg \neg \tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_{j'_n}[R^{k'_n}(\rho) + H_\rho^{k'_n} | \mathcal{F}_\rho] | \mathcal{F}_\nu] = \tilde{\mathcal{E}}_i[R^{k'_n}(\rho) + H_\rho^{k'_n} | \mathcal{F}_\nu] = \\ &\tilde{\mathcal{E}}_i[R^{j'_n}(\rho) + H_\rho^i | \mathcal{F}_\nu], \quad a. s. \end{aligned} \quad (8.73)$$

Then (?) and (5.3) imply that

$$C \leq Y_\rho^i = Y_\rho + H_\rho^i \leq R^{j'_n}(\rho) + H_\rho^i \leq R^{j'_1}(\rho) + H_\rho^i, \quad a. s.,$$

where $R^{j'_1}(\rho) + H_\rho^i \in Dom(\mathcal{E})$ thanks to Proposition 82 (1), (S1') and (D2). As $n \rightarrow \infty$ in (8.73), the Dominated Convergence Theorem (Proposition 29) imply that

$$V^i(\nu) \leq \lim_{n \rightarrow \infty} \tilde{\mathcal{E}}_i[R^{j'_n}(\rho) + H_\rho^i | \mathcal{F}_\nu] = \tilde{\mathcal{E}}_i[V(\rho) + H_\rho^i | \mathcal{F}_\nu] = \tilde{\mathcal{E}}_i[V^i(\rho) | \mathcal{F}_\nu], \quad a. s.,$$

which proves (5.11).

It remains to show that $\{V^i(\tau(0) \wedge t)\}_{t \in [0, T]}$ is an $\tilde{\mathcal{E}}_i$ -submartingale: To see this, we fix $0 \leq s < t \leq T$ and set $\nu \stackrel{\Delta}{=} \tau(0) \wedge s$, $\rho \stackrel{\Delta}{=} \tau(0) \wedge t$. It is clear that $\nu \leq \rho \leq \tau(0) \leq \tau(\nu)$, a.s., hence (5.11), Corollary 25 and Proposition 26 (5) show that

$$\begin{aligned} V^i(\tau(0) \wedge s) &= V^i(\nu) \leq \tilde{\mathcal{E}}_i[V^i(\rho) | \mathcal{F}_\nu] = \tilde{\mathcal{E}}_i[V^i(\tau(0) \wedge t) | \mathcal{F}_{\tau(0) \wedge s}] = \\ &\tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_i[V^i(\tau(0) \wedge t) | \mathcal{F}_{\tau(0)}] | \mathcal{F}_s] \\ &= \tilde{\mathcal{E}}_i[V^i(\tau(0) \wedge t) | \mathcal{F}_s], \quad a. s., \end{aligned}$$

which implies that $\{V^i(\tau(0) \wedge t)\}_{t \in [0, T]}$ is an $\tilde{\mathcal{E}}_i$ -submartingale. \square

Proof of Theorem 68: Proof of (1).

Step 1: For any $\rho, \nu \in \mathcal{S}_{0, T}$, we define

$$\Psi^\rho(\nu) \stackrel{\Delta}{=} \text{essinf}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[Y_\rho + H_{\rho \wedge \nu}^i | \mathcal{F}_{\rho \wedge \nu}] + H_{\rho \wedge \nu}^{i'} \in \mathcal{F}_{\rho \wedge \nu}.$$

It follows from (?), (?), and Proposition 26 (5) that

$$\begin{aligned} C_Y + 2C_H &= \text{essinf}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[C_Y + C_H | \mathcal{F}_{\rho \wedge \nu}] + C_H \\ &\leq \Psi^\rho(\nu) \leq \tilde{\mathcal{E}}_{i'}[Y_\rho + H_{\rho \wedge \nu}^{i'} | \mathcal{F}_{\rho \wedge \nu}] + H_{\rho \wedge \nu}^{i'} \leq R^{i'}(\rho \wedge \nu) + H_{\rho \wedge \nu}^{i'}, \\ &a. s., \end{aligned} \quad (8.74)$$

where $R^{i'}(\rho \wedge \nu) + H_{\rho \wedge \nu}^{i'} \in \text{Dom}(\mathcal{E})$ thanks to Proposition 82 (1), (S1') and (D2). Then Lemma 80 implies that $\Psi^\rho(\nu) \in \text{Dom}(\mathcal{E})$. Applying Proposition 26 (2)-(3) and Lemma 80, we can alternatively rewrite $\Psi^\rho(\nu)$ as follows:

$$\begin{aligned} \Psi^\rho(\nu) - \neg H_{\rho \wedge \nu}^{i'} \quad \neg \neg \neg &= \neg \neg \neg \text{essinf}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[\mathbf{11}_{\{\rho \leq \nu\}} Y_{\rho \wedge \nu} \neg + \neg \mathbf{11}_{\{\rho > \nu\}} \neg (Y_{\rho} \neg + \neg \\ &\quad H_{\nu, \rho}^i) | \mathcal{F}_{\rho \wedge \nu}] \\ \neg \neg \neg &= \neg \neg \neg \text{essinf}_{i \in \mathcal{I}} \neg (\mathbf{11}_{\{\rho \leq \nu\}} Y_{\rho \wedge \nu} \neg + \neg \mathbf{11}_{\{\rho > \nu\}} \tilde{\mathcal{E}}_i[Y_{\rho} \neg + \neg \\ &\quad H_{\nu, \rho}^i | \mathcal{F}_{\nu}]) \neg = \mathbf{11}_{\{\rho \leq \nu\}} Y_{\rho} \neg + \neg \mathbf{11}_{\{\rho > \nu\}} \text{essinf}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[Y_{\rho} \neg + \neg \\ &\quad \neg H_{\nu, \rho}^i | \mathcal{F}_{\nu}], \quad a. s. \end{aligned}$$

Let $\sigma \in \mathcal{S}_{0,T}$. Lemma 80 (2) and Proposition 26 (2) once again imply that

$$\begin{aligned} \mathbf{11}_{\{\nu=\sigma\}} \Psi^\rho(\nu) &= \mathbf{11}_{\{\rho \leq \nu=\sigma\}} Y_{\rho} \neg + \mathbf{11}_{\{\rho > \nu=\sigma\}} \text{essinf}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[Y_{\rho} + H_{\nu, \rho}^i | \mathcal{F}_{\nu}] + \\ &\quad \mathbf{11}_{\{\nu=\sigma\}} H_{\rho \wedge \nu}^{i'} \\ &= \mathbf{11}_{\{\rho \leq \nu=\sigma\}} Y_{\rho} \neg + \mathbf{11}_{\{\rho > \nu\}} \text{essinf}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[\mathbf{11}_{\{\nu=\sigma\}} (Y_{\rho} + \\ &\quad H_{\sigma, \rho}^i | \mathcal{F}_{\nu}) + \mathbf{11}_{\{\nu=\sigma\}} H_{\rho \wedge \sigma}^{i'} \\ &= \mathbf{11}_{\{\rho \leq \nu=\sigma\}} Y_{\rho} \neg + \mathbf{11}_{\{\rho > \nu\}} \text{essinf}_{i \in \mathcal{I}} \mathbf{11}_{\{\nu=\sigma\}} \tilde{\mathcal{E}}_i[Y_{\rho} + H_{\sigma, \rho}^i | \mathcal{F}_{\sigma}] + \\ &\quad \mathbf{11}_{\{\nu=\sigma\}} H_{\rho \wedge \sigma}^{i'} \\ &= \mathbf{11}_{\{\rho \leq \sigma=\nu\}} Y_{\rho} \neg + \mathbf{11}_{\{\rho > \sigma=\nu\}} \text{essinf}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[Y_{\rho} + H_{\sigma, \rho}^i | \mathcal{F}_{\sigma}] + \\ &\quad \mathbf{11}_{\{\nu=\sigma\}} H_{\rho \wedge \sigma}^{i'} = \mathbf{11}_{\{\nu=\sigma\}} \Psi^\rho(\sigma), \quad a. s. \end{aligned} \tag{8.75}$$

Step 2: Fix $\rho \in \mathcal{S}_{0,T}$. For any $\nu \in \mathcal{S}_{0,T}$ and $\sigma \in \mathcal{S}_{\nu,T}$, letting $(\nu, \mathcal{I}', \mathcal{U}) = (\rho \wedge \sigma, \mathcal{I}, \{\rho\})$ and $X(\rho) = Y_{\rho}$ in Lemma 80, we can find a sequence $\{j_n\}_{n \in \mathbb{N}}$ in \mathcal{I} such that

$$\text{essinf}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[Y_{\rho} + H_{\rho \wedge \sigma, \rho}^i | \mathcal{F}_{\rho \wedge \sigma}] = \lim_{n \rightarrow \infty} \neg \neg \downarrow \tilde{\mathcal{E}}_{j_n}[Y_{\rho} + H_{\rho \wedge \sigma, \rho}^{j_n} | \mathcal{F}_{\rho \wedge \sigma}], \quad a. s.$$

Definition 7.13 assures the existence of a $k_n = k(i', j_n, \rho \wedge \sigma) \in \mathcal{I}$ such that $\tilde{\mathcal{E}}_{k_n} = \mathcal{E}_{i' \neg, j_n}^{\rho \wedge \sigma}$. Applying Proposition 26 (3) to $\tilde{\mathcal{E}}_{k_n}$, we can deduce from (?) and (75) that

$$\begin{aligned} \Psi^\rho(\nu) &\leq \tilde{\mathcal{E}}_{k_n}[Y_{\rho} + H_{\rho \wedge \nu, \rho}^{k_n} | \mathcal{F}_{\rho \wedge \nu}] + H_{\rho \wedge \nu}^{i'} = \tilde{\mathcal{E}}_{k_n}[Y_{\rho} + H_{\rho \wedge \nu, \rho}^{k_n} | \mathcal{F}_{\rho \wedge \nu}] + H_{\rho \wedge \nu}^{k_n} = \\ &\quad \tilde{\mathcal{E}}_{k_n}[Y_{\rho}^{k_n} | \mathcal{F}_{\rho \wedge \nu}] \\ &= \mathcal{E}_{i' \neg, j_n}^{\rho \wedge \sigma}[Y_{\rho}^{k_n} | \mathcal{F}_{\rho \wedge \nu}] = \tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_{j_n}[Y_{\rho}^{k_n} | \mathcal{F}_{\rho \wedge \sigma}] | \mathcal{F}_{\rho \wedge \nu}], \quad a. s. \end{aligned}$$

For any $n \in \mathbb{N}$, Proposition 26 (3) & (5), (?) as well as (?) imply that

$$\begin{aligned} C &= \tilde{\mathcal{E}}_{j_n}[C | \mathcal{F}_{\rho \wedge \sigma}] \leq \tilde{\mathcal{E}}_{j_n}[Y_{\rho}^{k_n} | \mathcal{F}_{\rho \wedge \sigma}] = \tilde{\mathcal{E}}_{j_n}[Y_{\rho} + H_{\rho \wedge \sigma, \rho}^{j_n} | \mathcal{F}_{\rho \wedge \sigma}] + H_{\rho \wedge \sigma}^{i'} \\ &\leq \tilde{\mathcal{E}}_{j_1}[Y_{\rho} + H_{\rho \wedge \sigma, \rho}^{j_1} | \mathcal{F}_{\rho \wedge \sigma}] + H_{\rho \wedge \sigma}^{i'} \leq R^{j_1}(\rho \wedge \sigma) + H_{\rho \wedge \sigma}^{i'}, \quad a. s., \end{aligned}$$

where $R^{j_1}(\rho \wedge \sigma) + H_{\rho \wedge \sigma}^{i'}$ thanks to Proposition 82 (1), (S1') and (D2). Then the Dominated Convergence Theorem (Proposition 29), Corollary 25 and Proposition 26 (5) show that

$$\begin{aligned}
\Psi^\rho(\nu) &\leq \lim_{n \rightarrow \infty} \neg \neg \downarrow \tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_{j_n}[Y_\rho^{k_n} | \mathcal{F}_{\rho \wedge \sigma}] | \mathcal{F}_{\rho \wedge \nu}] = \lim_{n \rightarrow \infty} \neg \neg \downarrow \tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_{j_n}[Y_\rho + \\
&\quad H_{\rho \wedge \sigma, \rho}^{j_n} | \mathcal{F}_{\rho \wedge \sigma}] + H_{\rho \wedge \sigma}^{i'} | \mathcal{F}_{\rho \wedge \nu}] \\
&= \tilde{\mathcal{E}}_i\left[\lim_{n \rightarrow \infty} \neg \neg \downarrow \tilde{\mathcal{E}}_{j_n}[Y_\rho + H_{\rho \wedge \sigma, \rho}^{j_n} | \mathcal{F}_{\rho \wedge \sigma}] + H_{\rho \wedge \sigma}^{i'} | \mathcal{F}_{\rho \wedge \nu}\right] = \\
&\quad \tilde{\mathcal{E}}_i\left[\text{essinf}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[Y_\rho + H_{\rho \wedge \sigma, \rho}^{i'} | \mathcal{F}_{\rho \wedge \sigma}] + H_{\rho \wedge \sigma}^{i'} | \mathcal{F}_{\rho \wedge \nu}\right] \\
&= \tilde{\mathcal{E}}_i[\Psi^\rho(\sigma) | \mathcal{F}_{\rho \wedge \nu}] = \tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_i[\Psi^\rho(\sigma) | \mathcal{F}_\rho] | \mathcal{F}_\nu] = \tilde{\mathcal{E}}_i[\Psi^\rho(\sigma) | \mathcal{F}_\nu], \quad a. s.,
\end{aligned} \tag{8.76}$$

which implies that $\{\Psi^\rho(t)\}_{t \in [0, T]}$ is an $\tilde{\mathcal{E}}_i$ -submartingale. Hence, $\{-\Psi^\rho(t)\}_{t \in [0, T]}$ is an \mathcal{E}' -supermartingale by assumption (8.6). Since \mathcal{E}' satisfies (H0), (H1), (?) and since $\text{Dom}(\mathcal{E}') \in \tilde{\mathcal{D}}_T$ (which results from $\text{Dom}(\mathcal{E}) \in \tilde{\mathcal{D}}_T$ and (8.6)), we know from Theorem 20 that $\Psi_t^{\rho, +} \triangleq \liminf_{n \rightarrow \infty} \Psi^\rho(q_n^+(t))$, $t \in [0, T]$ is an RCLL process and that

$$P\left(\Psi_t^{\rho, +} = \lim_{n \rightarrow \infty} \Psi^\rho(q_n^+(t)) \mid \text{hbox} | \text{for any } t \in [0, T]\right) = 1. \tag{8.77}$$

Step 3: For any $\nu \in \mathcal{S}_{0, T}$ and $n \in \mathbb{N}$, $q_n^+(\nu)$ takes values in a finite set $\mathcal{D}_T^n \triangleq ([0, T] \cap \{k 2^{-n}\}_{k \in \mathbb{Z}}) \cup \{T\}$. Given an $\alpha \in \mathcal{D}_T^n$, it holds for any $m \geq n$ that $q_m^+(\alpha) = \alpha$ since $\mathcal{D}_T^n \subset \mathcal{D}_T^m$. It follows from (8.77) that

$$\Psi_\alpha^{\rho, +} = \lim_{m \rightarrow \infty} \Psi^\rho(q_m^+(\alpha)) = \Psi^\rho(\alpha), \quad a. s.$$

Then one can deduce from (8.74) that

$$\begin{aligned}
\Psi_{q_n^+(\nu)}^{\rho, +} &= \sum_{\alpha \in \mathcal{D}_T^n} \mathbf{1}_{\{q_n^+(\nu) = \alpha\}} \Psi_\alpha^{\rho, +} = \sum_{\alpha \in \mathcal{D}_T^n} \mathbf{1}_{\{q_n^+(\nu) = \alpha\}} \Psi^\rho(\alpha) = \\
&\sum_{\alpha \in \mathcal{D}_T^n} \mathbf{1}_{\{q_n^+(\nu) = \alpha\}} \Psi^\rho(q_n^+(\nu)) = \Psi^\rho(q_n^+(\nu)), \quad a. s.
\end{aligned}$$

Thus the right-continuity of the process $\Psi^{\rho, +}$ implies that

$$\Psi_\nu^{\rho, +} = \lim_{n \rightarrow \infty} \Psi_{q_n^+(\nu)}^{\rho, +} = \lim_{n \rightarrow \infty} \Psi^\rho(q_n^+(\nu)), \quad a. s. \tag{8.78}$$

We have assumed that $\text{esssup}_{t \in \mathcal{D}_T} \tilde{\mathcal{E}}_j[Y_\rho^j | \mathcal{F}_t] \in \text{Dom}(\mathcal{E})$ for some $j = j(\rho) \in \mathcal{I}$. It holds a.s. that

$$\tilde{\mathcal{E}}_j[Y_\rho^j | \mathcal{F}_t] \leq \text{esssup}_{s \in \mathcal{D}_T} \tilde{\mathcal{E}}_j[Y_\rho^j | \mathcal{F}_s], \quad \forall t \in \mathcal{D}_T.$$

Since $\tilde{\mathcal{E}}_j[Y_\rho^j | \mathcal{F}]$ is an RCLL process, it holds except on a null set N that

$$\begin{aligned}
&\tilde{\mathcal{E}}_j[Y_\rho^j | \mathcal{F}_t] \leq \text{esssup}_{s \in \mathcal{D}_T} \tilde{\mathcal{E}}_j[Y_\rho^j | \mathcal{F}_s], \quad \forall t \in [0, T], \quad \text{hbox} | \text{t h u s} \quad \tilde{\mathcal{E}}_j[Y_\rho^j | \mathcal{F}_{\rho \wedge q_n^+(\nu)}] \leq \\
&\text{esssup}_{s \in \mathcal{D}_T} \tilde{\mathcal{E}}_j[Y_\rho^j | \mathcal{F}_s], \quad \forall n \in \mathbb{N}.
\end{aligned}$$

Then one can deduce from (8.74), (?) and Proposition 26 (3) that

$$\begin{aligned} C_Y + 2C_H \neg\neg &\leq \neg\neg\Psi^\rho(q_n^+(\nu)) \leq \tilde{\mathcal{E}}_j[Y_\rho + H_{\rho \wedge q_n^+(\nu), \rho}^j | \mathcal{F}_{\rho \wedge q_n^+(\nu)}] + \neg\zeta^{i'} \neg= \\ &\tilde{\mathcal{E}}_j[Y_\rho^j - H_{\rho \wedge q_n^+(\nu)}^j | \mathcal{F}_{\rho \wedge q_n^+(\nu)}] + \neg\zeta^{i'} \\ \neg\neg &\leq \neg\neg\tilde{\mathcal{E}}_j[Y_\rho^j - C_H | \mathcal{F}_{\rho \wedge q_n^+(\nu)}] + \neg\zeta^{i'} = \tilde{\mathcal{E}}_j[Y_\rho^j | \mathcal{F}_{\rho \wedge q_n^+(\nu)}] - C_H + \\ &\neg\zeta^{i'} \neg \leq \text{esssup}_{s \in \mathcal{D}_T} \tilde{\mathcal{E}}_j[Y_\rho^j | \mathcal{F}_s] - C_H + \neg\zeta^{i'}, \quad a. s., \end{aligned}$$

where the right hand side belongs to $Dom(\mathcal{E})$ thanks to (D2) and the assumption that $\zeta^{i'} \in Dom(\mathcal{E})$. Hence the Dominated Convergence Theorem (Proposition 29), (8.78), (8.76) as well as Proposition 26 (5) imply that $\Psi_\nu^{\rho,+} = \lim_{n \rightarrow \infty} \Psi^\rho(q_n^+(\nu)) \in Dom(\mathcal{E})$ and that

$$\Psi^\rho(\nu) \leq \lim_{n \rightarrow \infty} \tilde{\mathcal{E}}_i[\Psi^\rho(q_n^+(\nu)) | \mathcal{F}_\nu] = \tilde{\mathcal{E}}_i[\Psi_\nu^{\rho,+} | \mathcal{F}_\nu] = \Psi_\nu^{\rho,+}, \quad a. s., \quad (8.79)$$

where in the last equality we used the fact that $\Psi_\nu^{\rho,+} = \lim_{n \rightarrow \infty} \Psi^\rho(q_n^+(\nu)) \in \mathcal{F}_\nu$, thanks to the right-continuity of the filtration F .

Step 4: Given $\nu \in \mathcal{S}_{0,T}$, we set

$$\gamma \stackrel{\Delta}{=} \tau(0) \wedge \nu, \quad \gamma_n \stackrel{\Delta}{=} \tau(0) \wedge q_n^+(\nu), \quad \forall n \in \mathbb{N}$$

and let $\rho \in \mathcal{S}_{\gamma,T}$. Since $\lim_{n \rightarrow \infty} \neg\uparrow \mathbf{1}_{\{\tau(0) > q_n^+(\nu)\}} = \mathbf{1}_{\{\tau(0) > \nu\}}$ and since

$$\{\tau(0) > \nu\} \subset \{q_n^+(\nu) = q_n^+(\tau(0) \wedge \nu)\}, \quad \{\tau(0) > q_n^+(\nu)\} \subset \{q_n^+(\nu) = \tau(0) \wedge q_n^+(\nu)\}, \quad \forall n \in \mathbb{N},$$

one can deduce from (8.79), (8.78) and (8.74) that

$$\begin{aligned} \mathbf{1}_{\{\tau(0) > \nu\}} \Psi^\rho(\gamma) &\leq \mathbf{1}_{\{\tau(0) > \nu\}} \Psi_\gamma^{\rho,+} = \mathbf{1}_{\{\tau(0) > \nu\}} \lim_{n \rightarrow \infty} \Psi^\rho(q_n^+(\gamma)) = \\ &\lim_{n \rightarrow \infty} \mathbf{1}_{\{\tau(0) > \nu\}} \Psi^\rho(q_n^+(\tau(0) \wedge \nu)) \\ &= \lim_{n \rightarrow \infty} \mathbf{1}_{\{\tau(0) > \nu\}} \Psi^\rho(q_n^+(\nu)) = \\ &\lim_{n \rightarrow \infty} \mathbf{1}_{\{\tau(0) > q_n^+(\nu)\}} \Psi^\rho(q_n^+(\nu)) \\ &= \lim_{n \rightarrow \infty} \mathbf{1}_{\{\tau(0) > q_n^+(\nu)\}} \Psi^\rho(\tau(0) \wedge q_n^+(\nu)) = \\ &\mathbf{1}_{\{\tau(0) > \nu\}} \lim_{n \rightarrow \infty} \Psi^\rho(\gamma_n), \quad a. s. \end{aligned} \quad (8.80)$$

For any $n \in \mathbb{N}$, we see from (5.9) that

$$\begin{aligned} V(\gamma_n) \neg\neg\neg &= \neg\neg\neg V(\gamma_n) = \text{esssup}_{\sigma \in \mathcal{S}_{\gamma_n,T}} \left(\text{essinf}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[Y_\sigma \neg\neg\neg H_{\gamma_n, \sigma}^i | \mathcal{F}_{\gamma_n}] \right) \geq \\ &\text{essinf}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[Y_{\rho \vee \gamma_n} \neg\neg\neg H_{\gamma_n, \rho \vee \gamma_n}^i | \mathcal{F}_{\gamma_n}], \quad a. s. \end{aligned} \quad (8.81)$$

Since $\{\tau(0) \leq \nu\} \subset \{\gamma_n = \gamma = \tau(0)\}$, Proposition 26 (2) and (3) imply that for any $i \in \mathcal{I}$

$$\begin{aligned} \mathbf{1}_{\{\tau(0) \leq \nu\}} \tilde{\mathcal{E}}_i[Y_\rho \neg\neg\neg H_{\rho \wedge \gamma_n, \rho}^i | \mathcal{F}_{\gamma_n}] &= \tilde{\mathcal{E}}_i[\mathbf{1}_{\{\tau(0) \leq \nu\}} (Y_\rho \neg\neg\neg H_{\rho \wedge \gamma, \rho}^i) | \mathcal{F}_{\gamma_n}] = \\ \mathbf{1}_{\{\tau(0) \leq \nu\}} \tilde{\mathcal{E}}_i[Y_\rho \neg\neg\neg H_{\rho \wedge \gamma, \rho}^i | \mathcal{F}_\gamma], \quad a. s., \end{aligned}$$

and that

$$\begin{aligned}
& \tilde{\mathcal{E}}_i[Y_{\rho \vee \gamma_n} \neg + \neg H_{\gamma_n, \rho \vee \gamma_n}^i | \mathcal{F}_{\gamma_n}] = \tilde{\mathcal{E}}_i[\mathbf{1}_{\{\rho \leq \gamma_n\}} Y_{\gamma_n} \neg + \neg \mathbf{1}_{\{\rho > \gamma_n\}} (Y_{\rho} \neg + \neg \\
& \quad H_{\rho \wedge \gamma_n, \rho}^i) | \mathcal{F}_{\gamma_n}] \\
& = \neg \mathbf{1}_{\{\rho \leq \gamma_n\}} Y_{\gamma_n} \neg + \neg \mathbf{1}_{\{\rho > \gamma_n\}} \tilde{\mathcal{E}}_i[Y_{\rho} \neg + \neg H_{\rho \wedge \gamma_n, \rho}^i | \mathcal{F}_{\gamma_n}] \\
& = \neg \mathbf{1}_{\{\rho \leq \gamma_n\}} Y_{\gamma_n} \neg + \neg \mathbf{1}_{\{\rho > \gamma_n, \mathcal{I}(0) > \nu\}} \tilde{\mathcal{E}}_i[Y_{\rho} \neg + \neg H_{\rho \wedge \gamma_n, \rho}^i | \mathcal{F}_{\rho \wedge \gamma_n}] + \\
& \quad \neg \mathbf{1}_{\{\rho > \gamma_n, \mathcal{I}(0) \leq \nu\}} \tilde{\mathcal{E}}_i[Y_{\rho} \neg + \neg H_{\rho \wedge \gamma, \rho}^i | \mathcal{F}_{\rho \wedge \gamma}], \quad a. s.
\end{aligned}$$

Then it follows from (8.81) and Lemma 80 that

$$\begin{aligned}
V(\gamma_n) \neg + \neg & \geq \neg \mathbf{1}_{\{\rho \leq \gamma_n\}} Y_{\gamma_n} \neg + \neg \mathbf{1}_{\{\rho > \gamma_n, \mathcal{I}(0) > \nu\}} \operatorname{essinf}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[Y_{\rho} \neg + \neg \\
& \quad H_{\rho \wedge \gamma_n, \rho}^i | \mathcal{F}_{\rho \wedge \gamma_n}] + \neg \mathbf{1}_{\{\rho > \gamma_n, \mathcal{I}(0) \leq \nu\}} \operatorname{essinf}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[Y_{\rho} \neg + \neg \\
& \quad H_{\rho \wedge \gamma, \rho}^i | \mathcal{F}_{\rho \wedge \gamma}] \\
& \neg + \neg = \neg \mathbf{1}_{\{\rho \leq \gamma_n\}} Y_{\gamma_n} \neg + \neg \mathbf{1}_{\{\rho > \gamma_n, \mathcal{I}(0) > \nu\}} (\Psi^\rho(\gamma_n) \neg + \neg H_{\rho \wedge \gamma_n}^{i'}) + \\
& \quad \neg \mathbf{1}_{\{\rho > \gamma_n, \mathcal{I}(0) \leq \nu\}} (\Psi^\rho(\gamma) \neg + \neg H_{\rho \wedge \gamma}^{i'}), \quad a. s. \tag{8.82}
\end{aligned}$$

Then (8.82), (8.83) and Proposition 26 (5) show that

$$\begin{aligned}
\Psi^\rho(\gamma_n) & = \operatorname{essinf}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[Y_{\rho} + H_{\rho \wedge \gamma_n, \rho}^i | \mathcal{F}_{\rho \wedge \gamma_n}] + H_{\rho \wedge \gamma_n}^{i'} \geq \operatorname{essinf}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[C_Y + C_H | \mathcal{F}_{\rho \wedge \gamma_n}] + \\
C_H & = C_Y + 2C_H, \quad a. s.
\end{aligned}$$

(f) As $n \rightarrow \infty$, the right-continuity of processes Y and $H^{i'}$, (8.79), Lemma 80 as well as Proposition 26 (2) & (3) show that

$$\begin{aligned}
\liminf_{n \rightarrow \infty} V(\gamma_n) & \geq \mathbf{1}_{\{\rho = \gamma\}} Y_{\gamma} \neg + \neg \mathbf{1}_{\{\rho > \gamma, \mathcal{I}(0) > \nu\}} \left(\lim_{n \rightarrow \infty} \Psi^\rho(\gamma_n) \neg + \neg H_{\rho \wedge \gamma}^{i'} \right) + \\
& \quad \neg \mathbf{1}_{\{\rho > \gamma, \mathcal{I}(0) \leq \nu\}} (\Psi^\rho(\gamma) \neg + \neg H_{\rho \wedge \gamma}^{i'}) \\
& \geq \mathbf{1}_{\{\rho = \gamma\}} Y_{\gamma} + \mathbf{1}_{\{\rho > \gamma\}} (\Psi^\rho(\gamma) - \neg H_{\rho \wedge \gamma}^{i'}) = \mathbf{1}_{\{\rho = \gamma\}} Y_{\gamma} + \\
& \quad \mathbf{1}_{\{\rho > \gamma\}} \operatorname{essinf}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[Y_{\rho} + \neg H_{\rho \wedge \gamma, \rho}^i | \mathcal{F}_{\rho \wedge \gamma}] \\
& = \operatorname{essinf}_{i \in \mathcal{I}} (\mathbf{1}_{\{\rho = \gamma\}} Y_{\gamma} + \mathbf{1}_{\{\rho > \gamma\}} \tilde{\mathcal{E}}_i[Y_{\rho} + H_{\gamma, \rho}^i | \mathcal{F}_{\gamma}]) = \\
& \quad \operatorname{essinf}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[\mathbf{1}_{\{\rho = \gamma\}} Y_{\gamma} + \mathbf{1}_{\{\rho > \gamma\}} (Y_{\rho} + H_{\gamma, \rho}^i) | \mathcal{F}_{\gamma}] \\
& = \operatorname{essinf}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[Y_{\rho} + H_{\gamma, \rho}^i | \mathcal{F}_{\gamma}], \quad a. s.
\end{aligned}$$

Taking the essential supremum of the right-hand-side over $\rho \in \mathcal{S}_{\gamma, T}$, we obtain

$$\liminf_{n \rightarrow \infty} V(\gamma_n) \geq \operatorname{esssup}_{\rho \in \mathcal{S}_{\gamma, T}} \left(\operatorname{essinf}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[Y_{\rho} + H_{\gamma, \rho}^i | \mathcal{F}_{\gamma}] \right) = V(\gamma) = V(\gamma), \quad a. s. \tag{8.83}$$

On the other hand, for any $i \in \mathcal{I}$ and $n \in \mathbb{N}$ we have that $V(\gamma_n) = \bar{V}(\gamma_n) = \operatorname{essinf}_{l \in \mathcal{I}} R^l(\gamma_n) \leq R^i(\gamma_n)$, a.s. Then (5.6) and the right continuity of the process $R^{i,0}$ imply that

$$\limsup_{n \rightarrow \infty} V(\gamma_n) \leq \lim_{n \rightarrow \infty} R^i(\gamma_n) = \lim_{n \rightarrow \infty} R_{\gamma_n}^{i,0} = R_{\gamma}^{i,0} = R^i(\gamma), \quad a. s.$$

Taking the essential infimum of $R^i(\gamma)$ over $i \in \mathcal{I}$ yields that

$$\limsup_{n \rightarrow \infty} V(\gamma_n) \leq \operatorname{essinf}_{i \in \mathcal{I}} R^i(\gamma) = \bar{V}(\gamma) = V(\gamma), \quad a.s.$$

This inequality together with (8.83) shows that $\lim_{n \rightarrow \infty} V(\gamma_n) = V(\gamma)$, a.s., which further implies that for any $\nu \in \mathcal{S}_{0,T}$ and $i \in \mathcal{I}$

$$\begin{aligned} \lim_{n \rightarrow \infty} V^{i'}(\mathcal{I}(0) \wedge q_n^+(\nu)) &= \lim_{n \rightarrow \infty} \neg \neg (V(\mathcal{I}(0) \wedge q_n^+(\nu)) + H_{\mathcal{I}(0) \wedge q_n^+(\nu)}^i) \\ &= V(\mathcal{I}(0) \wedge \nu) + H_{\mathcal{I}(0) \wedge \nu}^i = V^i(\mathcal{I}(0) \wedge \nu), \\ &\quad a.s. \end{aligned} \tag{8.84}$$

Step 5: Proposition 67 shows that the stopped process $\{V^{i'}(\mathcal{I}(0) \wedge t)\}_{t \in [0,T]}$ is an $\tilde{\mathcal{E}}_i$ -submartingale, thus $\{-V^{i'}(\mathcal{I}(0) \wedge t)\}_{t \in [0,T]}$ is an \mathcal{E}' -supermartingale by (8.6). Then Theorem 20 implies that $V_t^{i',+} \stackrel{\Delta}{=} \liminf_{n \rightarrow \infty} V^{i'}(\mathcal{I}(0) \wedge q_n^+(t))$, $t \in [0, T]$ is an RCLL process and that

$$P\left(V_t^{i',+} = \lim_{n \rightarrow \infty} V^{i'}(\mathcal{I}(0) \wedge q_n^+(t)) \mid \mathcal{F}_t\right) = 1.$$

For any $\sigma, \zeta \in \mathcal{S}_{0,T}$, Lemma 80 and (5.4) show that

$$\begin{aligned} \mathbf{1}_{\{\sigma=\zeta\}} V(\sigma) &= \neg \mathbf{1}_{\{\sigma=\zeta\}} \bar{V}(\sigma) = \operatorname{essinf}_{j \in \mathcal{I}} (\mathbf{1}_{\{\sigma=\zeta\}} R^j(\sigma)) \neg = \\ \operatorname{essinf}_{j \in \mathcal{I}} (\mathbf{1}_{\{\sigma=\zeta\}} R^j(\zeta)) \neg &= \neg \mathbf{1}_{\{\sigma=\zeta\}} \bar{V}(\zeta) = \neg \mathbf{1}_{\{\sigma=\zeta\}} V(\zeta), \quad a.s., \end{aligned}$$

which implies that

$$\begin{aligned} \mathbf{1}_{\{\sigma=\zeta\}} V^{i'}(\sigma) &= \mathbf{1}_{\{\sigma=\zeta\}} V(\sigma) + \mathbf{1}_{\{\sigma=\zeta\}} H_\sigma^{i'} = \mathbf{1}_{\{\sigma=\zeta\}} V(\zeta) + \mathbf{1}_{\{\sigma=\zeta\}} H_\zeta^{i'} \\ &= \mathbf{1}_{\{\sigma=\zeta\}} V^{i'}(\zeta), \quad a.s. \end{aligned} \tag{8.85}$$

Let $\sigma \in \mathcal{S}_{0,T}^F$ take values in a finite set $\{t_1 < \dots < t_m\}$. For any $\alpha \in \{1 \dots m\}$ and $n \in \mathbb{N}$, one can deduce from (8.85) that

$$\begin{aligned} \mathbf{1}_{\{\mathcal{I}(0) \wedge q_n^+(\sigma) = \mathcal{I}(0) \wedge q_n^+(t_\alpha)\}} V^{i'}(\mathcal{I}(0) \wedge q_n^+(\sigma)) &= \\ \mathbf{1}_{\{\mathcal{I}(0) \wedge q_n^+(\sigma) = \mathcal{I}(0) \wedge q_n^+(t_\alpha)\}} V^{i'}(\mathcal{I}(0) \wedge q_n^+(t_\alpha)), \quad a.s. \end{aligned}$$

(fi) since $\{\sigma = t_\alpha\} \subset \{\mathcal{I}(0) \wedge q_n^+(\sigma) = \mathcal{I}(0) \wedge q_n^+(t_\alpha)\}$, one can deduce from (8.85) that

$$\mathbf{1}_{\{\sigma=t_\alpha\}} V^{i'}(\mathcal{I}(0) \wedge q_n^+(\sigma)) = \mathbf{1}_{\{\sigma=t_\alpha\}} V^{i'}(\mathcal{I}(0) \wedge q_n^+(t_\alpha)), \quad a.s.$$

As $n \rightarrow \infty$, (8.83) shows that

$$\begin{aligned} \mathbf{1}_{\{\sigma=t_\alpha\}} V_\sigma^{i',+} &= \mathbf{1}_{\{\sigma=t_\alpha\}} V_{t_\alpha}^{i',+} = \lim_{n \rightarrow \infty} \mathbf{1}_{\{\sigma=t_\alpha\}} V^{i'}(\mathcal{I}(0) \wedge q_n^+(t_\alpha)) = \\ &\quad \lim_{n \rightarrow \infty} \mathbf{1}_{\{\sigma=t_\alpha\}} V^{i'}(\mathcal{I}(0) \wedge q_n^+(\sigma)) \\ &= \mathbf{1}_{\{\sigma=t_\alpha\}} V^{i'}(\mathcal{I}(0) \wedge \sigma), \quad a.s. \end{aligned}$$

Summing the above expression over α , we obtain $V_{\sigma}^{i',+} = V^{i'}(\tau(0) \wedge \sigma)$, a.s. Then the right-continuity of the process $V^{i',+}$ and (8.83) imply that

$$V_{\nu}^{i',+} = \lim_{n \rightarrow \infty} V_{q_n^+(\nu)}^{i',+} = \lim_{n \rightarrow \infty} V^{i'}(\tau(0) \wedge q_n^+(\nu)) = V^{i'}(\tau(0) \wedge \nu), \quad a.s. \quad (8.86)$$

In particular, $V^{i',+}$ is an RCLL modification of the stopped process $\{V^{i'}(\tau(0) \wedge t)\}_{t \in [0, T]}$. Therefore,

$V^0 \triangleq \{V_t^{i',+} - H_{\tau(0) \wedge t}^{i'}\}_{t \in [0, T]}$ is an RCLL modification of the stopped value process $\{V(\tau(0) \wedge t)\}_{t \in [0, T]}$. For any $\nu \in \mathcal{S}_{0, T}$, (8.86) implies that

$$V_{\nu}^0 = V_{\nu}^{i',+} - H_{\tau(0) \wedge \nu}^{i'} = V^{i'}(\tau(0) \wedge \nu) - H_{\tau(0) \wedge \nu}^{i'} = V(\tau(0) \wedge \nu), \quad a.s., \quad \text{\textcolor{violet}{\texttt{proving(5.13)}}}$$

Proof of (2). (5.13) and Proposition 68 imply that $V_{\tau(0)}^0 = V(\tau(0)) = Y_{\tau(0)}$, a.s. Hence, we can deduce from the right-continuity of processes V^0 and Y that τ_V in (5.14) is a stopping time belonging to $\mathcal{S}_{0, \tau(0)}$ and that

$$Y_{\tau_V} = V_{\tau_V}^0 = V(\tau_V), \quad a.s.,$$

where the second equality is due to (5.13). Then it follows from (5.11) that for any $i \in \mathcal{I}$

$$V(0) = V^i(0) \leq \tilde{\mathcal{E}}_i[V^i(\tau_V)] = \tilde{\mathcal{E}}_i[Y_{\tau_V}^i] = \mathcal{E}_i[Y_{\tau_V}^i].$$

In light of Lemma 61 and (5.9), we can find a sequence $\{j_n\}_{n \in \mathbb{N}}$ in \mathcal{I} such that

$$V(\tau_V) = \bar{V}(\tau_V) = \lim_{n \rightarrow \infty} \neg \neg \downarrow R^{j_n}(\tau_V), \quad a.s.$$

Fix $i \in \mathcal{I}$. For any $n \in \mathbb{N}$, Definition 7.13 assures that there exists a $k_n = k(i, j_n, \tau_V) \in \mathcal{I}$ such that $\tilde{\mathcal{E}}_{k_n} = \mathcal{E}_{i, j_n}^{\tau_V}$. Since $\tau_V \leq \tau(0)$, a.s., we can deduce from (5.7), Lemma 80, (5.13), (5.14) as well as Proposition 26 (5) that

$$\begin{aligned} V(0) &= \bar{V}(0) = \inf_{l \in \mathcal{I}} R^l(0) \leq R^{k_n}(0) = \tilde{\mathcal{E}}_{k_n}[R^{k_n}(\tau_V) + H_{\tau_V}^{k_n}] = \tilde{\mathcal{E}}_{k_n}[R^{j_n}(\tau_V) + H_{\tau_V}^i] \\ &= \mathcal{E}_{i, j_n}^{\tau_V}[R^{j_n}(\tau_V) + H_{\tau_V}^i] = \tilde{\mathcal{E}}_i[\mathcal{E}_{j_n}[R^{j_n}(\tau_V) + H_{\tau_V}^i | \mathcal{F}_{\tau_V}]] = \tilde{\mathcal{E}}_i[R^{j_n}(\tau_V) + H_{\tau_V}^i] \\ &= \tilde{\mathcal{E}}_i[V(\tau_V) + H_{\tau_V}^i] = \tilde{\mathcal{E}}_i[Y_{\tau_V} + H_{\tau_V}^i] \\ &= \tilde{\mathcal{E}}_i[Y_{\tau_V}^i] = \mathcal{E}_i[Y_{\tau_V}^i]. \end{aligned} \quad (8.87)$$

Moreover, (5.13) and (5.3) imply that

$$C \leq Y_{\tau_V}^i = Y_{\tau_V} + H_{\tau_V}^i \leq R^{j_n}(\tau_V) + H_{\tau_V}^i \leq R^{j_1}(\tau_V) + H_{\tau_V}^i, \quad a.s.,$$

where $R^{j_1}(\tau_V) + H_{\tau_V}^i \in \text{Dom}(\mathcal{E})$ thanks to Proposition 82 (1), (S1') and (D2). Letting $n \rightarrow \infty$ in (8.86) and using the Dominated Convergence Theorem (Proposition 29) and (5.13) we obtain that

$$\begin{aligned} V(0) &\leq \lim_{n \rightarrow \infty} \neg \neg \downarrow \tilde{\mathcal{E}}_i[R^{j_n}(\tau_V) + H_{\tau_V}^i] = \tilde{\mathcal{E}}_i\left[\lim_{n \rightarrow \infty} \neg \neg \downarrow R^{j_n}(\tau_V) + H_{\tau_V}^i\right] = \\ &= \tilde{\mathcal{E}}_i[V(\tau_V) + H_{\tau_V}^i] = \tilde{\mathcal{E}}_i[Y_{\tau_V} + H_{\tau_V}^i] \\ &= \tilde{\mathcal{E}}_i[Y_{\tau_V}^i] = \mathcal{E}_i[Y_{\tau_V}^i]. \end{aligned}$$

(fi) Taking the infimum of the right-hand-side over $i \in \mathcal{I}$ yields that

$$V(0) \leq \inf_{i \in \mathcal{I}} \mathcal{E}_i[Y_{\tau_V}^i] \leq \sup_{\rho \in \mathcal{S}_{0,T}} \left(\inf_{i \in \mathcal{I}} \mathcal{E}_i[Y_\rho^i] \right) = V(0) = V(0),$$

which implies that $\inf_{i \in \mathcal{I}} \mathcal{E}_i[Y_{\tau_V}^i] = \sup_{\rho \in \mathcal{S}_{0,T}} \inf_{i \in \mathcal{I}} \mathcal{E}_i[Y_\rho^i]$. \square

8.5 Proofs of Section 6

Proof of Proposition 82: Fix $t \in [0, T]$. For any $\xi \in Dom(\mathcal{E})$ and $i \in \mathcal{I}$, the definition of $Dom(\mathcal{E})$ assures that there exists a $c(\xi) \in \mathbb{R}$ such that $c(\xi) \leq \xi$, a.s. Then Proposition 26 (5) shows that

$$c(\xi) = \tilde{\mathcal{E}}_i[c(\xi)|\mathcal{F}_t] \leq \tilde{\mathcal{E}}_i[\xi|\mathcal{F}_t], \quad a. s. \quad (8.88)$$

Taking the essential infimum of the right-hand-side over $i \in \mathcal{I}$, we obtain for an arbitrary $i' \in \mathcal{I}$ that

$$c(\xi) \leq \mathcal{E}[\xi|\mathcal{F}_t] \leq \tilde{\mathcal{E}}_{i'}[\xi|\mathcal{F}_t], \quad a. s.$$

Since $\tilde{\mathcal{E}}_{i'}[\xi|\mathcal{F}_t] \in Dom^\#(\mathcal{E}_{i'}) = Dom(\mathcal{E})$, Lemma 80 implies that $\mathcal{E}[\xi|\mathcal{F}_t] \in Dom(\mathcal{E})$, thus $\mathcal{E}[\cdot|\mathcal{F}_t]$ is a mapping from $Dom(\mathcal{E})$ to $Dom_t(\mathcal{E}) = Dom(\mathcal{E}) \cap L^0(\mathcal{F}_t)$.

A simple application of Lemma 80 shows that \mathcal{E} satisfies (A3), (A4) and (?). Hence, it only remains to show (A2) for \mathcal{E} . Fix $0 \leq s < t \leq T$. Letting $(\nu, \mathcal{I}', \mathcal{U}) = (t, \mathcal{I}, \{T\})$ and taking $X(T) = \xi$ in Lemma 80, we can find a sequence $\{i_n\}_{n \in \mathbb{N}}$ in \mathcal{I} such that

$$\mathcal{E}[\xi|\mathcal{F}_t] = \text{essinf}_{i \in \mathcal{I}} \tilde{\mathcal{E}}_i[\xi|\mathcal{F}_t] = \lim_{n \rightarrow \infty} \neg \neg \downarrow \tilde{\mathcal{E}}_{i_n}[\xi|\mathcal{F}_t], \quad a. s. \quad (8.89)$$

Now fix $j \in \mathcal{I}$. For any $n \in \mathbb{N}$, it follows from Definition 7.13 that there exists a $k_n = k(j, i_n, t) \in \mathcal{I}$ such that $\tilde{\mathcal{E}}_{k_n} = \mathcal{E}_{j, i_n}^t$. Applying (75) yields that

$$\mathcal{E}[\xi|\mathcal{F}_s] \leq \tilde{\mathcal{E}}_{k_n}[\xi|\mathcal{F}_s] = \mathcal{E}_{j, i_n}^t[\xi|\mathcal{F}_s] = \tilde{\mathcal{E}}_j[\tilde{\mathcal{E}}_{i_n}[\xi|\mathcal{F}_t]|\mathcal{F}_s], \quad a. s. \quad (8.90)$$

For any $n \in \mathbb{N}$, we see from (8.88) and (8.89) that

$$c(\xi) = \tilde{\mathcal{E}}_{i_n}[c(\xi)|\mathcal{F}_t] \leq \tilde{\mathcal{E}}_{i_n}[\xi|\mathcal{F}_t] \leq \tilde{\mathcal{E}}_{i_1}[\xi|\mathcal{F}_t], \quad a. s.,$$

where $\tilde{\mathcal{E}}_{i_1}[\xi|\mathcal{F}_t] \in Dom^\#(\mathcal{E}_{i_1}) = Dom(\mathcal{E})$. The Dominated Convergence Theorem (Proposition 29) and (8.90) then imply that

$$\tilde{\mathcal{E}}_j[\mathcal{E}[\xi|\mathcal{F}_t]|\mathcal{F}_s] = \lim_{n \rightarrow \infty} \tilde{\mathcal{E}}_j[\tilde{\mathcal{E}}_{i_n}[\xi|\mathcal{F}_t]|\mathcal{F}_s] \geq \mathcal{E}[\xi|\mathcal{F}_s], \quad a. s.$$

Taking the essential infimum of the left-hand-side over $j \in \mathcal{I}$, we obtain

$$\mathcal{E}[\mathcal{E}[\xi|\mathcal{F}_t]|\mathcal{F}_s] \geq \mathcal{E}[\xi|\mathcal{F}_s], \quad a. s. \quad (8.91)$$

On the other hand, for any $i \in \mathcal{I}$ and $\rho \in \mathcal{S}_{t,T}$, applying Corollary 25, we obtain

$$\tilde{\mathcal{E}}_i[\xi|\mathcal{F}_s] = \tilde{\mathcal{E}}_i[\tilde{\mathcal{E}}_i[\xi|\mathcal{F}_t]|\mathcal{F}_s] \geq \tilde{\mathcal{E}}_i[\mathcal{E}[\xi|\mathcal{F}_t]|\mathcal{F}_s] \geq \mathcal{E}[\mathcal{E}[\xi|\mathcal{F}_t]|\mathcal{F}_s], \quad a.s.$$

Taking the essential infimum of the left-hand-side over $i \in \mathcal{I}$ yields that $\mathcal{E}[\xi|\mathcal{F}_s] \geq \mathcal{E}[\mathcal{E}[\xi|\mathcal{F}_t]|\mathcal{F}_s]$, a.s., which together with (8.91) proves (A2) for \mathcal{E} . \square

8.6 Proofs of Section 7

Proof of Proposition 72: By (7.2), it holds $dt \times dP - a.s.$ that for any $z \in \mathbb{R}^d$

$$|g(t, z)| = |g(t, z) - g(t, 0)| \leq K_g |z|, \quad \langle \text{hbox} | t h u s \rangle \quad \tilde{g}(t, z) \stackrel{\Delta}{=} -K_g |z| \leq g(t, z).$$

Clearly, \tilde{g} is a generator satisfying (7.2). It is also positively homogeneous in z , i.e.

$$\tilde{g}(t, \alpha z) = -K_g |\alpha z| = -\alpha K_g |z| = \alpha \tilde{g}(t, z), \quad \forall \alpha \geq 0, \forall z \in \mathbb{R}^d.$$

Then Example 10 of [\(citet|Peng-97\)](#) (or Proposition 8 of [\(citet|Eman\)](#)) and (7.6) imply that for any $n \in \mathbb{N}$ and any $A \in \mathcal{F}_T$ with $P(A) > 0$

$$n\mathcal{E}_{\tilde{g}}[11_A] = \mathcal{E}_{\tilde{g}}[n11_A] \leq \mathcal{E}_g[n11_A]. \quad (8.92)$$

Since $\mathcal{E}_{\tilde{g}}[11_A] > 0$ (which follows from the second part of (A1)), letting $n \rightarrow \infty$ in (8.92) yields (H0).

Next, we consider a sequence $\{\xi_n\}_{n \in \mathbb{N}} \subset L^2(\mathcal{F}_T)$ with $\sup_{n \in \mathbb{N}} \|\xi_n\| \in L^2(\mathcal{F}_T)$. If ξ_n converges a.s., it is clear that $\xi \stackrel{\Delta}{=} \lim_{n \rightarrow \infty} \xi_n \in L^2(\mathcal{F}_T)$. Applying Lemma 71 with $\mu = K_g$, we obtain

$$\begin{aligned} |\mathcal{E}_g[\xi_n] - \mathcal{E}_g[\xi]| &\leq \mathcal{E}_{g_\mu}[|\xi_n - \xi|] = \left| \mathcal{E}_{g_\mu}[|\xi_n - \xi|] \right|_{L^2(\mathcal{F}_T)} \leq \left| \sup_{t \in [0, T]} \mathcal{E}_{g_\mu}[|\xi_n - \xi| | \mathcal{F}_t] \right|_{L^2(\mathcal{F}_T)} \\ &\leq C e^{(K_g + K_g^2)T} \|\xi_n - \xi\|_{L^2(\mathcal{F}_T)}, \end{aligned}$$

where we used the fact that $K_{g_\mu} = \mu$ in the last inequality. As $n \rightarrow \infty$, thanks to the Dominated Convergence Theorem of the linear expectation E , we have that $\|\xi_n - \xi\|_{L^2(\mathcal{F}_T)}^2 = E \|\xi_n - \xi\|^2 \rightarrow 0$; thus $\lim_{n \rightarrow \infty} \mathcal{E}_g[\xi_n] = \mathcal{E}_g[\xi]$. Then (H1) and (H2) follow.

For any $\nu \in \mathcal{S}_{0,T}$ and $\xi \in L^{2,+}(\mathcal{F}_T) \stackrel{\Delta}{=} \{\xi \in L^2(\mathcal{F}_T) : \xi \geq 0, a.s.\}$, Lemma 71

(1) shows that $\sup_{t \in [0, T]} \left| \mathcal{E}_g[\xi | \mathcal{F}_t] \right| \in L^{2,+}(\mathcal{F}_T)$, consequently $\mathcal{E}_g[\xi | \mathcal{F}_\nu] \in L^{2,+}(\mathcal{F}_T)$.

Since $X \stackrel{\Delta}{=} \mathcal{E}_g[\xi | \mathcal{F}]$ is a continuous process, $X_\nu^{\xi,+} = X_\nu^\xi = \mathcal{E}_g[\xi | \mathcal{F}_\nu] \in L^{2,+}(\mathcal{F}_T)$, which proves (H3). \square

Proof of Proposition 75: Fix $\nu \in \mathcal{S}_{0,T}$. It is easy to check that the generator g^ν satisfies (7.2) with Lipschitz coefficient $K_1 \vee K_2$. For any $\xi \in L^2(\mathcal{F}_T)$, we set $\eta \triangleq \Gamma_\nu^{\xi, g_2} \in \mathcal{F}_\nu$ and define

$$\tilde{\Theta}_t \triangleq \mathbf{1}_{\{\nu \leq t\}} \Theta_t^{\xi, g_2} + \mathbf{1}_{\{\nu > t\}} \Theta_t^{\eta, g_1}, \quad \forall t \in [0, T].$$

It follows that

$$g^\nu(t, \tilde{\Theta}_t) = \mathbf{1}_{\{\nu \leq t\}} g_2(t, \tilde{\Theta}_t) + \mathbf{1}_{\{\nu > t\}} g_1(t, \tilde{\Theta}_t) = \mathbf{1}_{\{\nu \leq t\}} g_2(t, \Theta_t^{\xi, g_2}) + \mathbf{1}_{\{\nu > t\}} g_1(t, \Theta_t^{\eta, g_1}), \quad \forall t \in [0, T].$$

For any $t \in [0, T]$, since $\{\nu \leq t\} \in \mathcal{F}_t$, one can deduce that

$$\begin{aligned} & \mathbf{1}_{\{\nu \leq t\}} \left(\xi + \int_t^T \neg g^\nu(s, \tilde{\Theta}_s) ds - \int_t^T \neg \tilde{\Theta}_s dB_s \right) = \mathbf{1}_{\{\nu \leq t\}} \left(\xi + \int_t^T \neg \mathbf{1}_{\{\nu \leq t\}} g^\nu(s, \right. \\ & \quad \left. \tilde{\Theta}_s) ds - \int_t^T \neg \mathbf{1}_{\{\nu \leq t\}} \tilde{\Theta}_s dB_s \right) \\ &= \mathbf{1}_{\{\nu \leq t\}} \left(\xi + \int_t^T \neg \mathbf{1}_{\{\nu \leq t\}} g_2(s, \Theta_s^{\xi, g_2}) ds - \int_t^T \neg \mathbf{1}_{\{\nu \leq t\}} \Theta_s^{\xi, g_2} dB_s \right) \\ &= \mathbf{1}_{\{\nu \leq t\}} \left(\xi + \int_t^T \neg g_2(s, \Theta_s^{\xi, g_2}) ds - \int_t^T \neg \Theta_s^{\xi, g_2} dB_s \right) = \mathbf{1}_{\{\nu \leq t\}} \Gamma_t^{\xi, g_2}, \\ & \quad \text{a. s.} \end{aligned} \tag{8.93}$$

The continuity of processes $\int_t^T \neg g^\nu(s, \tilde{\Theta}_s) ds$, $\int_t^T \neg \tilde{\Theta}_s dB_s$ and Γ_t^{ξ, g_2} then implies that except on a null set N

$$\mathbf{1}_{\{\nu \leq t\}} \left(\xi + \int_t^T \neg g^\nu(s, \tilde{\Theta}_s) ds - \int_t^T \neg \tilde{\Theta}_s dB_s \right) = \mathbf{1}_{\{\nu \leq t\}} \Gamma_t^{\xi, g_2}, \quad \forall t \in [0, T].$$

Taking $t = \nu(\omega)$ for any $\omega \in N^c$ yields that

$$\xi + \int_\nu^T g^\nu(s, \tilde{\Theta}_s) ds - \int_\nu^T \tilde{\Theta}_s dB_s = \Gamma_\nu^{\xi, g_2} = \eta, \quad \text{a. s.} \tag{8.94}$$

Now fix $t \in [0, T]$. We can deduce from (8.94) that

$$\begin{aligned} & \mathbf{1}_{\{\nu > t\}} \left(\xi + \int_t^T g^\nu(s, \tilde{\Theta}_s) ds - \int_t^T \tilde{\Theta}_s dB_s \right) \\ & \quad \neg \neg = \neg \neg \mathbf{1}_{\{\nu > t\}} \left(\eta + \int_t^\nu g^\nu(s, \tilde{\Theta}_s) ds - \int_t^\nu \tilde{\Theta}_s dB_s \right) \\ & \quad \neg \neg = \neg \neg \mathbf{1}_{\{\nu > t\}} \left(\eta + \int_t^\nu g_1(s, \Theta_s^{\eta, g_1}) ds - \int_t^\nu \Theta_s^{\eta, g_1} dB_s \right), \\ & \quad \text{a. s.} \end{aligned} \tag{8.95}$$

Moreover, Proposition 26 (5) implies that

$$\begin{aligned}\mathcal{E}_{g_1}[\eta|\mathcal{F}_{t\wedge\nu}] &= \eta + \int_{t\wedge\nu}^T g_1(s, \Theta_s^{\eta, g_1}) ds - \int_{t\wedge\nu}^T \Theta_s^{\eta, g_1} dB_s = \mathcal{E}_{g_1}[\eta|\mathcal{F}_\nu] + \\ &\quad \int_{t\wedge\nu}^\nu g_1(s, \Theta_s^{\eta, g_1}) ds - \int_{t\wedge\nu}^\nu \Theta_s^{\eta, g_1} dB_s \\ &= \eta + \int_{t\wedge\nu}^\nu g_1(s, \Theta_s^{\eta, g_1}) ds - \int_{t\wedge\nu}^\nu \Theta_s^{\eta, g_1} dB_s, \quad a.s.\end{aligned}$$

Multiplying both sides with $\mathbf{1}_{\{\nu>t\}}$ and using (8.94), we obtain

$$\begin{aligned}\mathbf{1}_{\{\nu>t\}} \left(\xi + \int_t^T g^\nu(s, \tilde{\Theta}_s) ds - \int_t^T \tilde{\Theta}_s dB_s \right) &= \mathbf{1}_{\{\nu>t\}} \mathcal{E}_{g_1}[\eta|\mathcal{F}_t] = \\ \mathbf{1}_{\{\nu>t\}} \mathcal{E}_{g_1}[\Gamma_\nu^{\xi, g_2} | \mathcal{F}_t], \quad a.s.,\end{aligned}$$

which in conjunction with (8.93) shows that for any $t \in [0, T]$

$$\begin{aligned}\xi + \int_t^T g^\nu(s, \tilde{\Theta}_s) ds - \int_t^T \tilde{\Theta}_s dB_s &= \mathbf{1}_{\{\nu\leq t\}} \Gamma_t^{\xi, g_2} + \mathbf{1}_{\{\nu>t\}} \mathcal{E}_{g_1}[\Gamma_\nu^{\xi, g_2} | \mathcal{F}_t] \\ &= \mathbf{1}_{\{\nu\leq t\}} \mathcal{E}_{g_2}[\xi | \mathcal{F}_t] + \\ &\quad \mathbf{1}_{\{\nu>t\}} \mathcal{E}_{g_1}[\mathcal{E}_{g_2}[\xi | \mathcal{F}_\nu] | \mathcal{F}_t] = \mathcal{E}_{g_1, g_2}^\nu[\xi | \mathcal{F}_t], \\ &\quad a.s.\end{aligned}$$

Since $\int_t^T g^\nu(s, \tilde{\Theta}_s) ds$, $\int_t^T \tilde{\Theta}_s dB_s$ and $\mathcal{E}_{g_1, g_2}^\nu[\xi | \mathcal{F}_\cdot]$ are all continuous processes, it holds except a null N' that

$$\mathcal{E}_{g_1, g_2}^\nu[\xi | \mathcal{F}_t] = \xi + \int_t^T g^\nu(s, \tilde{\Theta}_s) ds - \int_t^T \tilde{\Theta}_s dB_s, \quad \forall t \in [0, T].$$

One can easily show that $(\mathcal{E}_{g_1, g_2}^\nu[\xi | \mathcal{F}_\cdot], \tilde{\Theta}) \in \mathbb{C}_{\mathcal{F}}^2([0, T]) \times \mathcal{H}_{\mathcal{F}}^2([0, T]; \mathbb{R}^d)$. Thus the pair is the unique solution to the BSDE (ξ, g^ν) , namely $\mathcal{E}_{g^\nu}[\xi | \mathcal{F}_t] = \mathcal{E}_{g_1, g_2}^\nu[\xi | \mathcal{F}_t]$ for any $t \in [0, T]$. \square

Proof of Theorem 82: We first note that for any $g \in \mathcal{G}'$, (7.7) implies that for every \mathcal{E}_g -submartingale X , $-X$ is an \mathcal{E}_g -supermartingale although g^- is concave (which means that \mathcal{E}_{g^-} may not belong to \mathcal{E}'). Hence, condition (?) is satisfied.

Fix $g \in \mathcal{G}'$. Clearly $H_0^g = 0$. For any $s, t \in \mathcal{D}_T$ with $s < t$, we can deduce from (h1) and (h2) that

$$\begin{aligned}C_{\mathcal{H}}^{\triangleleft, c'} T &\leq - \int_s^t -c' ds \leq - \int_s^t -h_r^g dr = H_{s, t}^g \leq \int_s^t h'(r) dr \leq \int_0^T h'(r) dr, \\ a.s.,\end{aligned} \tag{8.96}$$

which implies that

$$C_{\mathcal{H}}^{\triangleleft, c'} \leq \operatorname{essinf}_{s, t \in \mathcal{D}_T; s < t} H_{s, t}^g \leq \operatorname{esssup}_{s, t \in \mathcal{D}_T; s < t} H_{s, t}^g \leq \int_0^T h'(r) dr, \quad a.s.,$$

thus (S2) holds. Since $\int_0^T h'(r) dr \in L^2(\mathcal{F}_T)$, it follows that

$$\begin{aligned} \text{esssup}_{s,t \in \mathcal{D}_T; s < t} H_{s,t}^g \in L^{2,\#}(\mathcal{F}_T) &\stackrel{\Delta}{=} \{\xi \in L^2(\mathcal{F}_T) : \xi \geq c, \text{ a.s. } \langle \text{hbox} | \text{f o r s o m e} \rangle c \in \mathbb{R}\} = \\ &Dom(\mathcal{E}'). \end{aligned}$$

We can also deduce from (8.96) that except on a null set N

$$C_{\mathcal{H}'} \leq H_{s,t}^g \leq \int_0^T h'(r) dr, \quad \forall 0 \leq s < t \leq T.$$

Hence, for any $\nu, \rho \in \mathcal{S}_{0,T}$ with $\nu \leq \rho$, a.s., we have

$$C_{\mathcal{H}'} \leq H_{\nu,\rho}^g \leq \int_0^T h'(r) dr, \quad \text{a.s.},$$

which implies that $H_{\nu,\rho}^g \in L^{2,\#}(\mathcal{F}_T) = Dom(\mathcal{E}')$; so we got (S1'). Moreover, (S3) directly follows from (\tilde{h} 3).

Next, we check that the process Y satisfies (Y1) and (4.6). By (?) and (Y3), it holds a.s. that $C_Y \leq Y_t \leq \zeta_Y'$ for any $t \in \mathcal{D}_T$. The right-continuity of the process Y then implies that except on a null set \tilde{N}

$$C_Y \leq Y_t \leq \zeta_Y', \quad \forall t \in [0, T], \quad \langle \text{hbox} | \text{t h u s} \rangle C_Y \leq Y_\rho \leq \zeta_Y', \quad \forall \rho \in \mathcal{S}_{0,T}. \quad (8.97)$$

Since $\zeta_Y' \in L^2(\mathcal{F}_T)$, it follows that $Y_\rho \in L^{2,\#}(\mathcal{F}_T) = Dom(\mathcal{E}')$ for any $\rho \in \mathcal{S}_{0,T}$, thus (Y1) holds. Moreover, for any $g \in \mathcal{G}'$, $\rho \in \mathcal{S}_{0,T}$ and $t \in \mathcal{D}_T$, Proposition 8 (2), (8.97) and Lemma 71 (2) show that

$$\begin{aligned} C_Y + c' T &= \mathcal{E}_g[C_Y + c' T | \mathcal{F}_t] \leq \mathcal{E}_g\left[Y_\rho + \int_0^\rho c' ds \middle| \mathcal{F}_t\right] \leq \mathcal{E}_g[Y_\rho^g | \mathcal{F}_t] \leq \\ &\quad \left| \mathcal{E}_g[Y_\rho^g | \mathcal{F}_t] \right| \\ &= \left| \mathcal{E}_g[Y_\rho^g | \mathcal{F}_t] - \mathcal{E}_g[0 | \mathcal{F}_t] \right| \leq \mathcal{E}_{g_M}[|Y_\rho^g| | \mathcal{F}_t] \leq \mathcal{E}_{g_M}\left[\left|Y_\rho\right| + \int_0^\rho |h_s^g| ds \middle| \mathcal{F}_t\right] \\ &\leq \sup_{t \in [0, T]} \mathcal{E}_{g_M}[\zeta_Y' \vee (-C_Y) + \int_0^T h'(s) \vee (-c') ds \middle| \mathcal{F}_t], \quad \text{a.s.} \end{aligned}$$

Taking essential supremum of $\mathcal{E}_g[Y_\rho^g | \mathcal{F}_t]$ over $(g, \rho, t) \in \mathcal{G}' \times \mathcal{S}_{0,T} \times \mathcal{D}_T$, we can deduce from (A4) that

$$\begin{aligned} C_Y + c' T &\leq \text{esssup}_{(g, \rho, t) \in \mathcal{G}' \times \mathcal{S}_{0,T} \times \mathcal{D}_T} \mathcal{E}_g[Y_\rho^g | \mathcal{F}_t] \leq \sup_{t \in [0, T]} \mathcal{E}_{g_M}\left[\zeta_Y' + \int_0^T h'(s) ds \middle| \mathcal{F}_t\right] - C_Y - c' T, \quad \text{a.s.} \end{aligned} \quad (8.98)$$

Lemma 71 (1) implies that

$$\left| \sup_{t \in [0, T]} \mathcal{E}_{g_M} \left[\zeta'_{Y \neg + \neg} \int_0^T \neg h'(s) ds \middle| \mathcal{F}_t \right] \right|_{L^2(\mathcal{F}_T)} \leq C e^{(M+M^2)T} \left| \zeta'_{Y \neg + \neg} \int_0^T \neg h'(s) ds \right|_{L^2(\mathcal{F}_T)} < \infty.$$

Hence, we see from (8.98) that $\text{esssup}_{(g, \rho, t) \in \mathcal{G}' \times \mathcal{S}_{0, T} \times \mathcal{D}_T} \mathcal{E}_g[Y_\rho^g | \mathcal{F}_t] \in L^{2, \#}(\mathcal{F}_T) = \text{Dom}(\mathcal{E}')$, which is exactly (4.6).

Now we show that the family of processes $\{Y_t^g, t \in [0, T]\}_{g \in \mathcal{G}'}$ is both " \mathcal{E}' -uniformly-left-continuous" and " \mathcal{E}' -uniformly-right-continuous". For any $\nu, \rho \in \mathcal{S}_{0, T}$ with $\nu \leq \rho$, a.s., let $\{\rho_n\}_{n \in \mathbb{N}} \subset \mathcal{S}_{\nu, T}$ be a sequence increasing a.s. to ρ . For any $g \in \mathcal{G}'$, Lemma 71 (2) implies that

$$\begin{aligned} & \left| \mathcal{E}_g \left[\langle \text{hbox} | \frac{n}{n-1} \rangle Y_{\rho_n} + H_{\rho_n}^g \middle| \mathcal{F}_\nu \right] - \mathcal{E}_g[Y_\rho^g | \mathcal{F}_\nu] \right| \leq \mathcal{E}_{g_M} \left[\left| \langle \text{hbox} | \frac{n}{n-1} \rangle Y_{\rho_n} - Y_\rho - \int_{\rho_n}^\rho h^g(s) ds \right| \middle| \mathcal{F}_\nu \right] \\ & \leq \mathcal{E}_{g_M} \left[\left| \langle \text{hbox} | \frac{n}{n-1} \rangle Y_{\rho_n} - Y_\rho + \int_{\rho_n}^\rho \tilde{h}'(s) ds \right| \middle| \mathcal{F}_\nu \right], \quad a.s., \end{aligned}$$

where $g_M(z) \triangleq M|z|$, $z \in \mathbb{R}^d$ and $\tilde{h}'(t) \triangleq h'(t) - c'$, $t \in [0, T]$. Taking essential supremum of the left hand side over $g \in \mathcal{G}'$ yields that

$$\text{esssup}_{g \in \mathcal{G}'} \left| \mathcal{E}_g \left[\langle \text{hbox} | \frac{n}{n-1} \rangle Y_{\rho_n} + H_{\rho_n}^g \middle| \mathcal{F}_\nu \right] - \mathcal{E}_g[Y_\rho^g | \mathcal{F}_\nu] \right| \leq \mathcal{E}_{g_M} \left[\left| \langle \text{hbox} | \frac{n}{n-1} \rangle Y_{\rho_n} - Y_\rho + \int_{\rho_n}^\rho \tilde{h}'(s) ds \right| \middle| \mathcal{F}_\nu \right], \quad a.s. \quad (8.99)$$

Moreover, Lemma 71 (1) implies that

$$\begin{aligned} & \left| \mathcal{E}_{g_M} \neg \left[\left| \langle \text{hbox} | \frac{n}{n-1} \rangle Y_{\rho_n} \neg \neg Y_\rho \neg + \neg \int_{\rho_n}^\rho \neg \tilde{h}'(s) ds \right| \middle| \mathcal{F}_\nu \right] \right|_{L^2(\mathcal{F}_T)} \neg \leq \left| \sup_{t \in [0, T]} \mathcal{E}_{g_M} \neg \left[\left| \langle \text{hbox} | \frac{n}{n-1} \rangle Y_{\rho_n} \neg \neg Y_\rho \neg + \neg \int_{\rho_n}^\rho \neg \tilde{h}'(s) ds \right| \middle| \mathcal{F}_t \right] \right|_{L^2(\mathcal{F}_T)} \\ & \leq C e^{(M+M^2)T} \left| \langle \text{hbox} | \frac{n}{n-1} \rangle Y_{\rho_n} \neg \neg Y_\rho + \int_{\rho_n}^\rho \tilde{h}'(s) ds \right|_{L^2(\mathcal{F}_T)}. \end{aligned} \quad (8.100)$$

Since

$$\begin{aligned} & \left| \langle \text{hbox} | \frac{n}{n-1} \rangle Y_{\rho_n} \neg \neg Y_\rho \right| \leq \left| \langle \text{hbox} | \frac{n}{n-1} \rangle Y_{\rho_n} - Y_\rho \right| + \left| \langle \text{hbox} | \frac{1}{n-1} \rangle Y_\rho \right| \leq 2 \left| Y_{\rho_n} - Y_\rho \right| + \left| \langle \text{hbox} | \frac{1}{n-1} \rangle Y_\rho \right|, \quad \forall n \geq 2, \end{aligned}$$

the continuity of Y implies that $\lim_{n \rightarrow \infty} \left(\left| \langle \text{hbox} | \frac{n}{n-1} \rangle Y_{\rho_n} - Y_\rho \right| + \int_{\rho_n}^\rho \tilde{h}'(s) ds \right) = 0$, a.s. It also holds for any $n \geq 2$ that

$$\left| \langle \text{hbox} | \frac{n}{n-1} \rangle Y_{\rho_n} - Y_\rho \right| + \int_{\rho_n}^\rho \tilde{h}'(s) ds \leq 3(\zeta_Y - C_Y) + \int_0^T h'(s) ds - c'T, \\ \text{a.s.,}$$

where the right-hand sides belongs to $L^2(\mathcal{F}_T)$. Thus the Dominated Convergence Theorem implies that

$$\langle \text{hbox} | \text{thesequence} \rangle \left\{ \left| \langle \text{hbox} | \frac{n}{n-1} \rangle Y_{\rho_n} - Y_\rho \right| + \int_{\rho_n}^\rho \tilde{h}'(s) ds \right\}_{n \in \mathbb{N}} \langle \text{hbox} | \text{convergesto0in} L^2(\mathcal{F}_T), \rangle$$

which together with (8.99) and (8.100) implies that

$$\langle \text{hbox} | \text{thesequence} \rangle \left\{ \text{esssup}_{g \in \mathcal{G}'} \left| \mathcal{E}_g \left[\langle \text{hbox} | \frac{n}{n-1} \rangle Y_{\rho_n} + H_{\rho_n}^g | \mathcal{F}_\nu \right] - \mathcal{E}_g[Y_\rho^g | \mathcal{F}_\nu] \right| \right\}_{n \in \mathbb{N}} \langle \text{hbox} | \text{alsoconvergesto0in} L^2(\mathcal{F}_T), \rangle$$

Then we can find a subsequence $\{n_k\}_{k \in \mathbb{N}}$ of \mathbb{N} such that

$$\lim_{n \rightarrow \infty} \text{esssup}_{g \in \mathcal{G}'} \left| \mathcal{E}_g \left[\langle \text{hbox} | \frac{n_k}{n_k-1} \rangle Y_{\rho_{n_k}} + H_{\rho_{n_k}}^g | \mathcal{F}_\nu \right] - \mathcal{E}_g[Y_\rho^g | \mathcal{F}_\nu] \right| = 0, \quad \text{a.s.}$$

Therefore, the family of process $\{Y^g\}_{g \in \mathcal{G}'}$ is “ \mathcal{E}^ν -uniformly-left-continuous”. The “ \mathcal{E}^ν -uniform-right-continuity” of $\{Y^g\}_{g \in \mathcal{G}'}$ can be shown similarly. \square

Proof of Theorem 82: For any $U \in \mathfrak{U}$, Theorem 82 and Proposition 46 imply that $Z^{U,0} = \{Z_t^0 + \int_0^t h_s^U ds\}_{t \in [0,T]}$ is an \mathcal{E}_{g_U} -supermartingale. In light of the Doob-Meyer Decomposition of g -expectation (see e.g. (, ?, Theorem 3.3), or (, ?, Theorem 3.9)), there exists an RCLL increasing process Δ^U null at 0 and a process $\Theta^U \in \mathcal{H}_{\mathbb{F}}^2([0, T]; \mathbb{R}^d)$ such that

$$Z_t^{U,0} = Z_T^{U,0} + \int_t^T g_U(s, \Theta_s^U) ds + \Delta_T^U - \Delta_t^U - \int_t^T \Theta_s^U dB_s, \quad t \in [0, T]. \quad (8.101)$$

In what follows we will show that

$$U(t, \omega) \stackrel{\Delta}{=} u(t, \omega, \Theta_t^{U^0}(\omega)), \quad (t, \omega) \in [0, T] \times \Omega$$

is an optimal control desired, where $U^0 \equiv 0$ denotes the null control.

Recall that $\bar{\tau}(0) = \inf\{t \in [0, T] \mid Z_t^0 = Y_t\}$. Taking $t = \bar{\tau}(0)$ and $t = \bar{\tau}(0) \wedge t$ respectively in (8.101) and subtracting the former from the latter yields that

$$Z_{\bar{\tau}(0) \wedge t}^{U,0} = Z_{\bar{\tau}(0)}^{U,0} + \int_{\bar{\tau}(0) \wedge t}^{\bar{\tau}(0)} g_U(s, \Theta_s^U) ds + \Delta_{\bar{\tau}(0)}^U - \Delta_{\bar{\tau}(0) \wedge t}^U - \int_{\bar{\tau}(0) \wedge t}^{\bar{\tau}(0)} \Theta_s^U dB_s, \quad t \in [0, T], \quad (8.102)$$

which is equivalent to

$$\begin{aligned} Z_{\bar{\tau}(0) \wedge t}^0 &= Z_{\bar{\tau}(0)}^0 + \int_{\bar{\tau}(0) \wedge t}^{\bar{\tau}(0)} H(s, \Theta_s^U, U_s) ds + \Delta_{\bar{\tau}(0)}^U - \Delta_{\bar{\tau}(0) \wedge t}^U - \int_{\bar{\tau}(0) \wedge t}^{\bar{\tau}(0)} \Theta_s^U dB_s, \\ t &\in [0, T]. \end{aligned} \quad (8.103)$$

In particular, taking $U = U^0$, we obtain

$$\begin{aligned} Z_{\bar{\tau}(0) \wedge t}^0 &= Z_{\bar{\tau}(0)}^0 + \int_{\bar{\tau}(0) \wedge t}^{\bar{\tau}(0)} H(s, \Theta_s^{U^0}, U_s^0) ds + \Delta_{\bar{\tau}(0)}^{U^0} - \Delta_{\bar{\tau}(0) \wedge t}^{U^0} - \int_{\bar{\tau}(0) \wedge t}^{\bar{\tau}(0)} \Theta_s^{U^0} dB_s, \\ t &\in [0, T]. \end{aligned} \quad (8.104)$$

Comparing the martingale parts of (8.103) and (8.104), we see that for any $U \in \mathfrak{U}$,

$$\Theta_t^U = \Theta_t^{U^0}, \quad dt \times dP - a.s. \quad (8.105)$$

on the stochastic interval

$$0, \bar{\tau}(0)$$

$\triangleq \{(t, \omega) \in [0, T] \times \Omega : 0 \leq t \leq \bar{\tau}(0)\}$. Plugging this back into (8.103) yields that

$$\begin{aligned} Z_{\bar{\tau}(0) \wedge t}^0 &= Z_{\bar{\tau}(0)}^0 + \int_{\bar{\tau}(0) \wedge t}^{\bar{\tau}(0)} H(s, \Theta_s^{U^0}, U_s^0) ds + \Delta_{\bar{\tau}(0)}^U - \Delta_{\bar{\tau}(0) \wedge t}^U - \int_{\bar{\tau}(0) \wedge t}^{\bar{\tau}(0)} \Theta_s^{U^0} dB_s, \\ t &\in [0, T]. \end{aligned} \quad (8.106)$$

which implies that

$$\begin{aligned} \int_{\bar{\tau}(0) \wedge t}^{\bar{\tau}(0)} H(t, \omega, \Theta_s^{U^0}, U_s^0) ds + \Delta_{\bar{\tau}(0)}^U - \Delta_{\bar{\tau}(0) \wedge t}^U &= \int_{\bar{\tau}(0) \wedge t}^{\bar{\tau}(0)} H(t, \omega, \Theta_s^{U^0}, U_s^0) ds + \\ \Delta_{\bar{\tau}(0)}^U - \Delta_{\bar{\tau}(0) \wedge t}^U & \end{aligned}$$

(fi)

Let us define $g_{K_o}(z) \triangleq K_o|z|$, $z \in \mathbb{R}^d$. Note that it is not necessary that $g_{K_o} = g_U$ for some $U \in \mathfrak{U}$. For any $U \in \mathfrak{U}$, we set $\Gamma_t \triangleq \mathcal{E}_{g_U}[Z_{\bar{\tau}(0)}^{U,0} | \mathcal{F}_t]$ and $\hat{\Gamma}_t \triangleq \mathcal{E}_{g_{K_o}}[-\Delta_{\bar{\tau}(0)}^{U*} | \mathcal{F}_t]$, $t \in [0, T]$, which are the solutions to the BSDE($Z_{\bar{\tau}(0)}^{U,0}, g_U$) and BSDE($-\Delta_{\bar{\tau}(0)}^{U*}, g_{K_o}$) respectively, i.e.,

$$\begin{aligned} \Gamma_t &= Z_{\bar{\tau}(0)}^{U,0} + \int_t^T g_U(s, \Theta_s) ds - \int_t^T \Theta_s dB_s, \quad t \in [0, T], \\ \hat{\Gamma}_t &= -\Delta_{\bar{\tau}(0)}^U + \int_t^T K_o \left| \hat{\Theta}_s \right| ds - \int_t^T \hat{\Theta}_s dB_s, \quad t \in [0, T], \end{aligned}$$

where $\Theta, \hat{\Theta} \in \mathcal{H}_{\mathbb{F}}^2([0, T]; \mathbb{R}^d)$. Applying Proposition 26 (5) and Corollary 25, we obtain that for any $t \in [0, T]$

$$\begin{aligned} \Gamma_{\bar{\tau}(0)} - \Gamma_{\bar{\tau}(0) \wedge t} &= \mathcal{E}_{g_U}[Z_{\bar{\tau}(0)}^{U,0} | \mathcal{F}_{\bar{\tau}(0)}] - \mathcal{E}_{g_U}[Z_{\bar{\tau}(0)}^{U,0} | \mathcal{F}_{\bar{\tau}(0) \wedge t}] = Z_{\bar{\tau}(0)}^{U,0} - \\ &\quad \mathcal{E}_{g_U}[\mathcal{E}_{g_U}[Z_{\bar{\tau}(0)}^{U,0} | \mathcal{F}_{\bar{\tau}(0)}] | \mathcal{F}_t] \\ &= Z_{\bar{\tau}(0)}^{U,0} - \mathcal{E}_{g_U}[Z_{\bar{\tau}(0)}^{U,0} | \mathcal{F}_t] = Z_{\bar{\tau}(0)}^{U,0} - \Gamma_t, \quad a.s. \end{aligned}$$

Then the continuity of processes Γ and $Z^{U,0}$ imply that

$$\begin{aligned}
\Gamma_t - Z_{\bar{\tau}(0) \wedge t}^{U,0} &= Z_{\bar{\tau}(0)}^{U,0} - Z_{\bar{\tau}(0) \wedge t}^{U,0} + \Gamma_{\bar{\tau}(0) \wedge t} - \Gamma_{\bar{\tau}(0)} = Z_{\bar{\tau}(0)}^{U,0} - Z_{\bar{\tau}(0) \wedge t}^{U,0} + \int_{\bar{\tau}(0) \wedge t}^{\bar{\tau}(0)} g_U(s, \\
&\quad \Theta_s) ds - \int_{\bar{\tau}(0) \wedge t}^{\bar{\tau}(0)} \Theta_s dB_s \\
&= -\Gamma_{\bar{\tau}(0)} Z_{\bar{\tau}(0)}^0 - Z_{\bar{\tau}(0) \wedge t}^0 + \int_{\bar{\tau}(0) \wedge t}^{\bar{\tau}(0)} H(s, \Theta_s, U_s) ds - \int_{\bar{\tau}(0) \wedge t}^{\bar{\tau}(0)} \Theta_s dB_s \\
&= -\Gamma_{\bar{\tau}(0)} \Delta_{\bar{\tau}(0)}^U + \Delta_{\bar{\tau}(0) \wedge t}^U + \int_{\bar{\tau}(0) \wedge t}^{\bar{\tau}(0)} [H(s, \Theta_s, U_s) - H(s, \Theta_s^{U^0}, U_s)] ds - \\
&\quad \int_{\bar{\tau}(0) \wedge t}^{\bar{\tau}(0)} (\Theta_s - \Theta_s^{U^0}) dB_s, \quad t \in [0, T],
\end{aligned}$$

where we used (8.106) with $U = U^*$ in the last inequality. Since it holds $dt \times dP$ -a.s. that

$$\begin{aligned}
H(t, \Theta_t, U_t) - H(t, \Theta_t^{U^0}, U_t) &= H(t, \Theta_t, U_t) - H(t, \Theta_t^{U^0}, u(t, \Theta_t^{U^0})) \leq H(t, \Theta_t, U_t) - H(t, \\
&\quad \Theta_t^{U^0}, U_t) \\
&= |g^o(t, \Theta_t, U_t) - g^o(t, \Theta_t^{U^0}, U_t)| \leq |g^o(t, \Theta_t, U_t) - g^o(t, \Theta_t^{U^0}, U_t)| \leq K_o |\Theta_t - \\
&\quad \Theta_t^{U^0}|,
\end{aligned}$$

the comparison Theorem for BSDEs (see e.g. ([10], Theorem 35.3)) implies that

$$\hat{\Gamma}_t \geq \Gamma_t - Z_{\bar{\tau}(0) \wedge t}^{U,0} - \Delta_{\bar{\tau}(0) \wedge t}^U, \quad t \in [0, T].$$

In particular, when $t=0$, we can deduce from (4.17) that

$$\mathcal{E}_{g_{K_o}}[-\Delta_{\bar{\tau}(0)}^U] \geq \mathcal{E}_{g_U}[Z^U(\bar{\tau}(0))] - Z(0).$$

Taking supremum of the right hand side over $U \in \mathcal{U}$ and applying Theorem 57 with $\nu=0$, we obtain

$$0 \geq \mathcal{E}_{g_{K_o}}[-\Delta_{\bar{\tau}(0)}^U] \geq \sup_{U \in \mathcal{U}} \mathcal{E}_{g_U}[Z^U(\bar{\tau}(0))] - Z(0) = 0,$$

thus $\mathcal{E}_{g_{K_o}}[-\Delta_{\bar{\tau}(0)}^{U^*}] = 0$. The strict monotonicity of g -expectation (see e.g. ([10], Proposition 2.2(iii))) then implies that $\Delta_{\bar{\tau}(0)}^{U^*} = 0$, a.s. Plugging it back to (8.102) and using (8.105), we obtain

$$\begin{aligned}
Z_{\bar{\tau}(0) \wedge t}^{U^*,0} &= Z_{\bar{\tau}(0)}^{U^*,0} + \int_{\bar{\tau}(0) \wedge t}^{\bar{\tau}(0)} g_U(s, \Theta_s^{U^0}) ds - \int_{\bar{\tau}(0) \wedge t}^{\bar{\tau}(0)} \Theta_s^{U^0} dB_s \\
&= Z_{\bar{\tau}(0)}^{U^*,0} + \int_t^T g_U(s, \mathbf{1}_{\{s \leq \bar{\tau}(0)\}} \Theta_s^{U^0}) ds - \int_t^T \mathbf{1}_{\{s \leq \bar{\tau}(0)\}} \Theta_s^{U^0} dB_s, \\
&\quad t \in [0, T],
\end{aligned} \tag{8.107}$$

which implies that $\mathcal{E}_{g_U^*}[Z_{\bar{\tau}(0)}^{U^*,0}|\mathcal{F}_t] = Z_{\bar{\tau}(0)\wedge t}^{U^*,0}, \forall t \in [0, T]$. Namely, $\{Z_{\bar{\tau}(0)\wedge t}^{U^*,0}\}_{t \in [0, T]}$ is a g_U^* -martingale. Eventually, letting $t = 0$ in (8.106), we can deduce from (4.17) and Theorem 57 that

$$Z(0) = Z_0^{U^*,0} = \mathcal{E}_{g_U}[Z_{\bar{\tau}(0)}^{U^*,0}] = \mathcal{E}_{g_U}\left[Z(\bar{\tau}(0)) + \int_0^{\bar{\tau}(0)} h_s^U ds\right] = \mathcal{E}_{g_U}\left[Y_{\bar{\tau}(0)} + \int_0^{\bar{\tau}(0)} h_s^U ds\right]. \quad \langle \text{hbox} | \quad \square \rangle$$

Proof of Proposition 82: Because of its linearity in z , the primary generator

$$g^o(t, \omega, z, u) \stackrel{\Delta}{=} \sigma^{-1}(t, X(\omega)) f(t, X(\omega), u), z \quad \forall (t, \omega, z, u) \in [0, T] \times \Omega \times \mathbb{R}^d \times S \quad (8.108)$$

satisfies $(g^o 2)$ and $(g^o 4)$. Then $(g^o 1)$ follows from the continuity of the process $\{X(t)\}_{t \in [0, T]}$ as well as the measurability of the volatility σ and of the function f . Moreover, (7.15) and (7.21) imply that for a.e. $t \in [0, T]$

$$\begin{aligned} |g^o(t, \omega, z_1, u) - g^o(t, \omega, z_2, u)| &= |\sigma^{-1}(t, X(\omega)) f(t, X(\omega), u), \\ &\quad z_1 - z_2| \leq |\sigma^{-1}(t, X(\omega))| \cdot |f(t, \\ &\quad X(\omega), u)| \cdot |z_1 - z_2| \\ &\leq K^2 |z_1 - z_2|, \quad \forall z_1, z_2 \in \mathbb{R}^d, \\ &\quad \forall (\omega, u) \in \Omega \times S, \end{aligned}$$

which shows that g^o satisfies $(g^o 4)$ with $K_o = K^2$. Clearly, $\tilde{\mathcal{U}} = \mathcal{H}_{\mathbf{F}}^0([0, T]; S)$ is closed under the pasting in the sense of (?). Hence, we know from last section that $\{\mathcal{E}_{g_U}\}_{U \in \tilde{\mathcal{U}}}$ is a stable class of g -expectations, where g_U is defined in (?).

Fix $U \in \tilde{\mathcal{U}}$. For any $\xi \in L^2(\mathcal{F})$, we see from (?) that

$$\begin{aligned} \mathcal{E}_{g_U}[\xi|\mathcal{F}_t] &= \xi + \int_t^T g_U(s, \Theta_s) ds - \int_t^T \Theta_s dB_s \\ &= \xi + \int_t^T \sigma^{-1}(s, X) f(s, X, U_s), \Theta_s ds - \int_t^T \Theta_s dB_s = \xi - \\ &\quad \int_t^T \Theta_s dB_s^U, \quad t \in [0, T], \end{aligned}$$

where $B_t^U \stackrel{\Delta}{=} B_t - \int_0^t \sigma^{-1}(s, X) f(s, X, U_s) ds, t \in [0, T]$ is a Brownian Motion with respect to P_U . For any $t \in [0, T]$, taking $E_U[\cdot|\mathcal{F}_t]$ on both sides above yields that

$$\mathcal{E}_{g_U}[\xi|\mathcal{F}_t] = E_U[\mathcal{E}_{g_U}[\xi|\mathcal{F}_t]|\mathcal{F}_t] = E_U[\xi|\mathcal{F}_t] - E_U\left[\int_t^T \Theta_s dB_s^U \middle| \mathcal{F}_t\right] = E_U[\xi|\mathcal{F}_t], \quad a. s. \quad (8.109)$$

Hence the g -expectation \mathcal{E}_{g_U} coincides with the linear expectation E_U on $L^2(\mathcal{F}_T)$.

Clearly, the process $Y \triangleq \{\varphi(X(t))\}_{t \in [0, T]}$ satisfies (Y3) since φ is bounded from below by $-K$. We see from (7.20) that for any $t \in [0, T]$

$$Y_t = \varphi(X(t)) \leq K|X(t)| \leq K\|X\|_T.$$

Taking essential supremum of Y_t over $t \in \mathcal{D}_T$ yields that

$$\zeta'_Y \triangleq \left(\operatorname{esssup}_{t \in \mathcal{D}_T} Y_t \right)^+ \leq K\|X\|_T, \quad a.s. \quad (8.110)$$

For any $t \in [0, T]$, the Burkholder-Davis-Gundy inequality, $(\sigma 1)$, (7.15) as well as Fubini Theorem imply that

$$\begin{aligned} E[(\|X\|_t)^2] &= E\left[\sup_{s \in [0, t]} |X(s)|^2\right] \leq 2x^2 + 2E\left\{\sup_{s \in [0, t]} \left|\int_0^s \sigma(r, X) dB_r\right|^2\right\} \leq \\ &2x^2 + 2CE \int_0^t |\sigma(s, X)|^2 ds \\ &\leq 2x^2 + 4C \int_0^t |\sigma(s, \vec{0})|^2 ds + 4CE \int_0^t |\sigma(s, X) - \sigma(s, \vec{0})|^2 ds \\ &\leq 2x^2 + 4C \int_0^T |\sigma(s, \vec{0})|^2 ds + 4Cn^2 K^2 \int_0^t E[(\|X\|_s)^2] ds. \end{aligned}$$

Then applying Gronwall's inequality yields that

$$E[(\|X\|_T)^2] \leq \left(2x^2 + 4C \int_0^T |\sigma(s, \vec{0})|^2 ds\right) e^{4Cn^2 K^2 T} < \infty, \quad (8.111)$$

which together with (8.110) shows that $\zeta'_Y \in L^2(\mathcal{F}_T)$, proving (?).

Next, we define a function $h^o(t, \omega, u) \triangleq h(t, X(\omega), u)$, $\forall (t, \omega, u) \in [0, T] \times \Omega \times S$. The continuity of the process $\{X(t)\}_{t \in [0, T]}$ and the measurability of the function h imply that h^o is $\mathcal{P} \otimes \mathfrak{S} / \mathcal{B}(\mathbb{R})$ -measurable. We see from (7.21) that h^o satisfies $(\hat{h}1)$. It also follows from (7.21) that for a.e. $t \in [0, T]$ and for any $\omega \in \Omega$,

$$h_t^U(\omega) \triangleq h^o(t, \omega, U_t(\omega)) = h(t, X(\omega), U_t(\omega)) \leq K\|X(\omega)\|_T, \quad \forall U \in \tilde{\mathfrak{U}}.$$

Taking essential supremum of $h_t^U(\omega)$ over $U \in \tilde{\mathfrak{U}}$ with respect to the product measure space $([0, T] \times \Omega, \mathcal{P}, \lambda \times P)$ yields that

$$\hat{h}(t, \omega) \triangleq \left(\operatorname{esssup}_{U \in \tilde{\mathfrak{U}}} h_t^U(\omega) \right)^+ \leq K\|X(\omega)\|_T, \quad dt \times dP - a.s.,$$

which leads to that $\int_0^T \hat{h}(t, \omega) dt \leq KT\|X(\omega)\|_T^*$, a.s. Hence, (8.111) implies that $\int_0^T \hat{h}(t, \omega) dt \in L^2(\mathcal{F}_T)$, proving $(\hat{h}2)$ for h^o .

We can apply the optimal stopping theory developed in Section 4 to the triple $(\{\mathcal{E}_{g_U}\}_{U \in \tilde{\mathcal{U}}}, \{h^U\}_{U \in \tilde{\mathcal{U}}}, Y)$ and use (8.109) to obtain (7.17). In addition, if there exists a measurable mapping $u^*: [0, T] \times \Omega \times \mathbb{R}^d \mapsto S$ satisfying (7.18), then (?) indicates that for any $(t, \omega, z) \in [0, T] \times \Omega \times \mathbb{R}^d$

$$\begin{aligned} \sup_{u \in S} (g^o(t, \omega, z, u) + h^o(t, \omega, u)) &= \sup_{u \in S} \tilde{H}(t, X(\omega), z, u) = \tilde{H}(t, X(\omega), z, u(t, \\ &\quad X(\omega), z)) \\ &= \neg\neg\neg g^o(t, \omega, z, u(t, X(\omega), z)) + h^o(t, \omega, \\ &\quad u(t, X(\omega), z)), \end{aligned}$$

which shows that (7.14) holds for the mapping $\tilde{u}^*(t, \omega, z) = u^*(t, X(\omega), z)$, $(t, \omega, z) \in [0, T] \times \Omega \times \mathbb{R}^d$. Therefore, an application of Theorem 82 yields (7.19) for some $U^* \in \tilde{\mathcal{U}}$. \square

Proof of Proposition 79: (7.24) directly follows from (, ?, Theorem 5).

To see (7.25), we set $\Delta\Gamma \stackrel{\Delta}{=} \Gamma^{\xi_1, \hat{g}} - \Gamma^{\xi_2, \hat{g}}$ and $\Delta\Theta \stackrel{\Delta}{=} \Theta^{\xi_1, \hat{g}} - \Theta^{\xi_2, \hat{g}}$, then (7.22)(i) implies that

$$\begin{aligned} d\Delta\Gamma_t &= -(\hat{g}(t, \Theta_t^{\xi_1, \hat{g}}) - \hat{g}(t, \Theta_t^{\xi_2, \hat{g}})) dt + \Delta\Theta_t dB_t = -\int_0^1 \frac{\partial \hat{g}}{\partial z}(t, \lambda \Delta\Theta_t + \\ &\quad \Theta_t^{\xi_2, \hat{g}}) \Delta\Theta_t d\lambda dt + \Delta\Theta_t dB_t \\ &= \Delta\Theta_t (-a_t dt + dB_t), \quad t \in [0, T], \end{aligned}$$

where $a_t \stackrel{\Delta}{=} \int_0^1 \frac{\partial \hat{g}}{\partial z}(\lambda \Delta\Theta_t + \Theta_t^{\xi_2, \hat{g}}) d\lambda$, $t \in [0, T]$. Since $M_{\mathbf{F}}([0, T]; \mathbb{R}^d) \subset M_{\mathbf{F}}^2([0, T]; \mathbb{R}^d) = \mathcal{H}_{\mathbf{F}}^2([0, T]; \mathbb{R}^d)$, one can deduce from (7.22)(ii) that

$$\begin{aligned} E \int_0^T |a_t|^2 dt &\leq E \int_0^T \neg \int_0^1 \left| \frac{\partial \hat{g}}{\partial z}(\lambda \Delta\Theta_t + \Theta_t^{\xi_2, \hat{g}}) \right|^2 d\lambda dt \leq 2\kappa^2 T + 2\kappa^2 E \int_0^T \neg \\ &\quad \int_0^1 |\lambda \Theta_t^{\xi_1, \hat{g}} + (1 - \lambda) \Theta_t^{\xi_2, \hat{g}}|^2 d\lambda dt \\ &\leq 2\kappa^2 T + \frac{4}{3} \kappa^2 E \int_0^T (|\Theta_t^{\xi_1, \hat{g}}|^2 + |\Theta_t^{\xi_2, \hat{g}}|^2) dt < \infty. \end{aligned}$$

Moreover, Doob's martingale inequality shows that

$$E \left[\sup_{t \in [0, T]} \left| \int_0^t a_s dB_s \right|^2 \right] \leq 4E \left[\left| \int_0^T a_s dB_s \right|^2 \right] \neg = 4E \int_0^T |a_t|^2 dt < \infty. \quad (8.112)$$

Thus, we can define process $Q_t \stackrel{\Delta}{=} \exp \left\{ \neg -\frac{1}{2} \int_0^t |a_s|^2 ds + \int_0^t a_s dB_s \right\}$, $t \in [0, T]$ as well as stopping times

$$\nu_n \stackrel{\Delta}{=} \inf \{ t \in [\nu, T] : Q_t \vee |\Delta\Gamma_t| > n \} \wedge T, \quad \forall n \in \mathbb{N}.$$

It is clear that $\lim_{n \rightarrow \infty} \neg \uparrow \nu_n = T$, a.s., and (8.112) assures that there exists a null set N such that for any $\omega \in N^c$, $T = \nu_m(\omega)$ for some $m = m(\omega) \in \mathbb{N}$.

For any $n \in \mathbb{N}$, integrating by parts on $[\nu, \nu_n]$ yields that

$$\begin{aligned} Q_{\nu_n} \Delta \Gamma_{\nu_n} &= Q_{\nu} \Delta \Gamma_{\nu} - \int_{\nu}^{\nu_n} Q_t \Delta \Theta_t a_t dt + \int_{\nu}^{\nu_n} Q_t \Delta \Theta_t dB_t + \int_{\nu}^{\nu_n} \Delta \Gamma_t Q_t a_t dB_t + \int_{\nu}^{\nu_n} Q_t \Delta \Theta_t a_t dt \\ &= \int_{\nu}^{\nu_n} (Q_t \Delta \Theta_t + \Delta \Gamma_t Q_t a_t) dB_t. \end{aligned}$$

which implies that $E[Q_{\nu_n} \Delta \Gamma_{\nu_n}] = 0$. Thus we can find a null set N_n such that $\Delta \Gamma_{\nu_n(\omega)}(\omega) = 0$, $\forall \omega \in N_n^c$. Eventually, for any $\omega \in \left\{ N \cup \left(\bigcup_{n \in \mathbb{N}} N_n \right) \right\}^c$, we have

$$\xi^1(\omega) = \Gamma_T^{\xi^1, \hat{g}}(\omega) = \lim_{n \rightarrow \infty} \Gamma_{\nu_n(\omega)}^{\xi^1, \hat{g}}(\omega) = \lim_{n \rightarrow \infty} \Gamma_{\nu_n(\omega)}^{\xi^2, \hat{g}}(\omega) = \Gamma_T^{\xi^2, \hat{g}}(\omega) = \xi^2(\omega). \quad \langle \text{hbox} \mid \square \rangle$$

Proof of Proposition 81: Let $\{A_n\}_{n \in \mathbb{N}}$ be any sequence in \mathcal{F}_T such that $\lim_{n \rightarrow \infty} \mathbb{1}_{A_n} = 0$, a.s. For any $\xi, \eta \in L^{\neg e, +}(\mathcal{F}_T) \stackrel{\Delta}{=} \{\xi \in L^{\neg e}(\mathcal{F}_T) : \xi \geq 0, \text{ a. s.}\}$, since

$$E[e^{\lambda|\xi|}] < \infty \quad \langle \text{hbox} \mid \text{and since} \rangle \quad \sup_{n \in \mathbb{N}} E[e^{\lambda|\xi + \mathbb{1}_{A_n} \eta|}] \leq E[e^{\lambda|\xi|} e^{\lambda|\eta|}] \leq \frac{1}{2} E[e^{2\lambda|\xi|}] + \frac{1}{2} E[e^{2\lambda|\eta|}] < \infty$$

holds for each $\lambda > 0$, Lemma 80 implies that

$$0 = \lim_{n \rightarrow \infty} E \left[\sup_{t \in [0, T]} \left| \mathcal{E}_{\hat{g}}[\xi + \mathbb{1}_{A_n} \eta | \mathcal{F}_t] - \mathcal{E}_{\hat{g}}[\xi | \mathcal{F}_t] \right| \right] \geq \lim_{n \rightarrow \infty} \left| \mathcal{E}_{\hat{g}}[\xi + \mathbb{1}_{A_n} \eta] - \mathcal{E}_{\hat{g}}[\xi] \right| \geq 0,$$

thus $\mathcal{E}_{\hat{g}}$ satisfies (H2). Similarly, we can show that (H1) also holds for $\mathcal{E}_{\hat{g}}$.

Moreover, for any $\nu \in \mathcal{S}_{0, T}$ and $\xi \in L^{\neg e, +}(\mathcal{F}_T)$, since the process $\Gamma^{\xi, \hat{g}}$ belongs to $\mathbb{C}_{\mathcal{F}}^e([0, T])$, one can deduce that $\mathcal{E}_{\hat{g}}[\xi | \mathcal{F}_{\nu}] = \Gamma_{\nu}^{\xi, \hat{g}} \in L^{\neg e, +}(\mathcal{F}_T)$. Then the continuity of the process $X^{\xi} \stackrel{\Delta}{=} \mathcal{E}_{\hat{g}}[\xi | \mathcal{F}_{\cdot}]$ implies that $X_{\nu}^{\xi, +} = X_{\nu}^{\xi} = \mathcal{E}_{\hat{g}}[\xi | \mathcal{F}_{\nu}] \in L^{\neg e, +}(\mathcal{F}_T)$, which proves (H3). \square

Proof of Theorem 82: This proof is just an application of the optimal stopping theory developed in Section 4 to the singleton $\{\mathcal{E}_{\hat{g}}\}$. Hence, it suffices to check that Y satisfies (Y1), (Y2) and (4.21).

Similar to (8.97), it holds except on a null set N that

$$C_Y \leq Y_t \leq \hat{\zeta}_Y, \quad \forall t \in [0, T], \quad \langle \text{hbox} \mid \text{thus} \rangle \quad C_Y \leq Y_{\rho} \leq \hat{\zeta}_Y, \quad \forall \rho \in \mathcal{S}_{0, T}. \quad (8.113)$$

Since $\hat{\zeta}_Y \in L^{\neg e}(\mathcal{F}_T)$, it holds for any $\rho \in \mathcal{S}_{0, T}$ that

$$E[e^{\lambda|Y_{\rho}|}] \leq E[e^{\lambda(\hat{\zeta}_Y - C_Y)}] = e^{-\lambda C_Y} E[e^{\lambda \hat{\zeta}_Y}] < \infty, \quad \forall \lambda > 0, \quad (8.114)$$

which implies that $Y_\rho \in L^{-e, \#}(\mathcal{F}_T) = D o m(\{\mathcal{E}_{\hat{g}}\})$. Hence (Y1) holds.

Next, for any $\rho \in \mathcal{S}_{0,T}$ and $t \in \mathcal{D}_T$, Proposition 8 (2), (?) show that

$$C_Y = \mathcal{E}_{\hat{g}}[C_Y | \mathcal{F}_t] \leq \mathcal{E}_{\hat{g}}[Y_\rho | \mathcal{F}_t] \leq \mathcal{E}_{\hat{g}}[\hat{\zeta}_Y | \mathcal{F}_t] = \Gamma_t^{\hat{\zeta}_Y, \hat{g}} \leq \sup_{t \in [0, T]} \left| \Gamma_t^{\hat{\zeta}_Y, \hat{g}} \right|, \quad a. s.$$

Taking essential supremum of $\mathcal{E}_{\hat{g}}[Y_\rho | \mathcal{F}_t]$ over $(\rho, t) \in \mathcal{S}_{0,T} \times \mathcal{D}_T$ yields that

$$C_Y \leq \operatorname{esssup}_{(\rho, t) \in \mathcal{S}_{0,T} \times \mathcal{D}_T} \mathcal{E}_{\hat{g}}[Y_\rho | \mathcal{F}_t] \leq \sup_{t \in [0, T]} \left| \Gamma_t^{\hat{\zeta}_Y, \hat{g}} \right|, \quad a. s.$$

Since $\Gamma^{\hat{\zeta}_Y, \hat{g}} \in \mathbb{C}_{\mathcal{F}}^e([0, T])$, or equivalently $\sup_{t \in [0, T]} \left| \Gamma_t^{\hat{\zeta}_Y, \hat{g}} \right| \in L^{-e}(\mathcal{F}_T)$, we can deduce that $\operatorname{esssup}_{(\rho, t) \in \mathcal{S}_{0,T} \times \mathcal{D}_T} \mathcal{E}_{\hat{g}}[Y_\rho | \mathcal{F}_t] \in L^{-e, \#}(\mathcal{F}_T) = D o m(\{\mathcal{E}_{\hat{g}}\})$, which together with Remark 73 (2) proves (Y2).

Moreover, for any $\nu, \rho \in \mathcal{S}_{0,T}$ with $\nu \leq \rho$, a.s. and any sequence $\{\rho_n\}_{n \in \mathbb{N}} \subset \mathcal{S}_{\nu, T}$ increasing a.s. to ρ , the continuity of the process Y implies that $\frac{n}{n-1} Y_{\rho_n}$ converges to Y_ρ a.s. By (?), one can deduce that

$$\sup_{n \in \mathbb{N}} E \left[\exp \left\{ \lambda \left| \langle \text{hbox} | \frac{n}{n-1} \rangle Y_{\rho_n} \right| \right\} \right] \leq \sup_{n \in \mathbb{N}} E[e^{2\lambda |Y_{\rho_n}|}] \leq E[e^{2\lambda(\hat{\zeta}_Y - C_Y)}] = e^{-2\lambda C_Y} E[e^{2\lambda \hat{\zeta}_Y}] < \infty, \quad \forall \lambda > 0,$$

which together with (8.114) allows us to apply Lemma 80:

$$0 = \lim_{n \rightarrow \infty} E \left[\sup_{t \in [0, T]} \left| \mathcal{E}_{\hat{g}}[\langle \text{hbox} | \frac{n}{n-1} \rangle Y_{\rho_n} | \mathcal{F}_t] - \mathcal{E}_{\hat{g}}[Y_\rho | \mathcal{F}_t] \right| \right] \geq \lim_{n \rightarrow \infty} E \left[\left| \mathcal{E}_{\hat{g}}[\langle \text{hbox} | \frac{n}{n-1} \rangle Y_{\rho_n} | \mathcal{F}_\nu] - \mathcal{E}_{\hat{g}}[Y_\rho | \mathcal{F}_\nu] \right| \right] \geq 0,$$

thus $\lim_{n \rightarrow \infty} E \left[\left| \mathcal{E}_{\hat{g}}[\langle \text{hbox} | \frac{n}{n-1} \rangle Y_{\rho_n} | \mathcal{F}_\nu] - \mathcal{E}_{\hat{g}}[Y_\rho | \mathcal{F}_\nu] \right| \right] = 0$. Then we can find a subsequence $\{n_k\}_{k \in \mathbb{N}}$ of \mathbb{N} such that

$$\lim_{n \rightarrow \infty} \left| \mathcal{E}_{\hat{g}}[\langle \text{hbox} | \frac{n_k}{n_k-1} \rangle Y_{\rho_{n_k}} | \mathcal{F}_\nu] - \mathcal{E}_{\hat{g}}[Y_\rho | \mathcal{F}_\nu] \right| = 0, \quad a. s.,$$

proving (4.21) for Y . \square

Bibliography